

Catalogue of Type Specimens of Recent Fishes in Field Museum of Natural History

Abstract

We list 1,406 nominal taxa of Recent fishes for which there are type specimens present in Field Museum of Natural History, including 837 holotypes, eight lectotypes, three neotypes, 48 taxa represented by syntypes, and 510 taxa represented only by secondary types. A large proportion of the species was described between 1905 and 1930 by Seth E. Meek, Samuel F. Hildebrand, Carl H. Eigenmann, David S. Jordan, and their associates. About 500 primary types came to Field Museum in 1952 with the purchase of the Carnegie Museum of Natural History fish collection. The main list of types presented here is arranged alphabetically by genus; each entry includes genus, species, and subspecies (if any) as it was originally proposed; author(s) and year of publication; family where the taxon is presently classified; abbreviated literature citation to original description; current type status; catalogue number(s) for primary types; total number of syntypes and secondary types; and brief locality information. In a few cases, comments are included to indicate discrepancies in previously published spellings or catalogue numbers, citations to subsequent designations of neotypes and lectotypes, and status of the material if uncertain. Also included are a list of the same taxa arranged by family and an alphabetical index of species.

Introduction

Field Museum of Natural History is one of seven recognized international depositories of Recent fishes in North America and is ranked among the

top five in the United States and Canada with respect to its type holdings (Collette & Lachner, 1976; Nitecki, 1980). The collection includes type material for 1,406 nominal taxa, of which 896 are represented by primary types (including 837 holotypes, eight lectotypes, three neotypes, and 48 taxa represented by syntypes). This catalogue was prepared in response to Recommendation 17D(4) of the International Code of Zoological Nomenclature (ICZN) to publish information on holdings of type materials. It also stems from our sense of obligation to the international community of systematic ichthyologists to provide the best possible curation for the types that have been entrusted to this institution since it was founded in 1893.

The type specimens of Recent fishes in Field Museum's collection stem from (1) exploratory fieldwork and systematic research of curators and associates, (2) purchase of the fish collection of the Carnegie Museum, Pittsburgh, in 1952, and (3) gifts and exchanges received from researchers throughout the world. Acquisition of the Carnegie Museum collection, with about 500 taxa represented by primary types, more than doubled the type collection at Field Museum. Most of the taxa in the collection were described after 1900, and a large proportion of them between 1905 and 1930 (fig. 1). Most of the specimens in our collection from that time period were described by Seth E. Meek (Field Museum), Samuel F. Hildebrand (United States National Museum), Carl H. Eigenmann (Indiana University), David S. Jordan (Stanford University), and their associates.

This catalogue represents a synthesis of information presented in two previously published type lists (Henn, 1928; Grey, 1947) with that in our catalogues and associated recent literature. Henn (1928) reported only the primary types in the Car-




FIG. 1. Frequency histogram for number of species of Recent fishes described in each year through 1984 that are represented by type material in Field Museum of Natural History.

negie Museum collection, while Grey (1947) listed all types in Field Museum. These two lists cover about 1,000 of our 1,406 taxa. In preparing this catalogue, the published lists and subsequent catalogue entries were checked against all jars on the shelf and outstanding loan invoices. A first draft of this list was then prepared, and nearly every entry was checked against the original publication. Those few taxa for which we were not able to get a copy of the original publication were included only if the names were clearly established in the literature; references for all such taxa (i.e., five species) are indicated as "not seen."

This list includes the types of all taxa present in Field Museum's fish collection that, to our knowledge, were described by mid-1985 when the final revision of this catalogue was submitted. It seems likely that a few other species based in part on FMNH materials were published in the last few months, but that we have not received a copy of the paper(s). The fish collections of Field Museum include specimens for about 50 taxa which have been labeled "manuscript types", none of which were included in this list. We were able to trace most of these hopeful taxa to determine their present status, but a few remain unresolved. We urge colleagues who find omissions or other errors in this list to notify Division of Fishes staff at Field Museum so that records can be updated.

Format of the List

The list of types is arranged alphabetically by genus and, within each genus, by species or subspecies as it was originally proposed without subsequent relocation. Information included for each species is as follows:

GENUS—This is spelled as it appears in the original description of the species, except where an emendation is clearly called for according to the rules of the ICZN, or where the author apparently used an inappropriate emendation of a previously published generic name. In almost every such case, the spelling used here has appeared in the recent literature. When the spelling used here differs from that used in the species description, the presumed incorrect spelling is given in a comment at the end of the entry. Subgeneric assignments were included in some species descriptions, but are omitted here.

SPECIES AND SUBSPECIES—These are spelled as they appear in the original description, except that

diacritical marks and hyphens have been deleted as recommended by the ICZN (i.e., hyphens are retained only in species names such as *x-punctata*). In a few cases, the original publication included a typographic error; such names are corrected to agree with current usage, and a comment to that effect is added at the end of the entry.

AUTHOR(S) AND DATE—The surname of the author(s) of the species is given followed by the year in which the publication was issued.

LITERATURE REFERENCE—Title of the journal where the description was published is abbreviated (using BIOSIS, 1981, where possible) and followed by volume number, issue number in parentheses, and page number on which the description begins. Some book titles are abbreviated. When authorship of the publication differed from that of the species described within, author(s) of the publication precedes the title of the journal or book.

PRIMARY TYPES—Categories used here for primary types are holotype, syntype(s), lectotype, and neotype. Field Museum of Natural History (FMNH) catalogue numbers are given for all primary types, and for former Carnegie Museum (CM) types, catalogue numbers are given in parentheses after FMNH number. For syntypes, the total number of specimens is given just ahead of the catalogue number(s). Missing types are indicated as such, and omission of a catalogue number means simply that no catalogue entry was ever made for the missing lot. A question mark indicates some uncertainty about type status of our material, and the problem is discussed at the end of the entry. If a lectotype or neotype has been designated, the relevant literature reference is given. Finally, the type locality is presented in general terms. Geographic and geopolitical place names have been changed in some cases to agree with modern usage (e.g., Guyana for British Guiana). We follow Bartholomew et al. (1980) for spellings of many of the place names.

SECONDARY TYPES—Categories of secondary types used here are paratype, lectoparatype, and allotype. Allotypes are considered equivalent to paratypes, but they are catalogued separately in the collection. The total number of secondary types is given followed by brief locality information. We consider secondary types to be all materials other than primary types that were used by the author to describe the taxon. Many early workers and occasionally recent workers gave incomplete data on the secondary types, and deciding which should be considered as such is, in some cases, subjective.

We excluded any material for which there was not at least a vague reference to the collection locality in the original description. In this regard, we differ from Grey (1947) who included materials from localities not given in the original description. We also excluded specimens which were not explicitly called types in the original description *and* indicated by the author as differing in some way from the other types.

FAMILY—Assignment of each taxon to a family and spelling of family names follows Nelson (1984) for most entries, because that work is the most recent and comprehensive synthesis dealing with all of the world's fish families. We differ from Nelson (1984) and follow Robins et al. (1980) for the more widely used spellings of Cobitidae, Eleotridae, Engraulidae, and Paralepididae.

List of Types

Arranged Alphabetically by Genus

Abudefduf lorenzi Hensley and Allen, 1977

Rec. W. Aust. Mus., 6(1): 109.

Holotype: FMNH 23564. Solomon Islands: Isabel Island, Tenibuli Bay.

Paratypes: 2. Solomon and Philippine Islands: 2 localities.

POMACENTRIDAE

Acahara jusanensis Jordan and Hubbs, 1925

Mem. Carnegie Mus., 10(2): 180.

Holotype: FMNH 58722 (CM 7828). Japan: Lake Jusan near Aomori.

CYPRINIDAE

Acanthemblemaria greenfieldi Smith-Vaniz and Palacio, 1974

Proc. Acad. Nat. Sci. Phila., 125(11): 210.

Paratypes: 7. Belize: Glover's Reef.

CHAENOPSIDAE

Acanthicus canensis Meek and Hildebrand, 1913

Field Mus. Nat. Hist. Zool. Ser., 10(8): 80.

Holotype: FMNH 7581. Panama: Río Cana, Cana.

Paratypes: 30. Panama: 2 localities.

LORICARIIDAE

Acanthocharax microlepis Eigenmann, 1912

Mem. Carnegie Mus., 5: 405.

Holotype: FMNH 54475 (CM 2138). Guyana: Tumatumari.

Paratypes: 31. Guyana: 3 localities.

CHARACIDAE

Acanthophaelus bifurcus Eigenmann, 1909

Ann. Carnegie Mus., 6(1): 52.

Holotype: FMNH 53539 (CM 1088). Guyana: Christianburg.

Paratypes: 36. Guyana: 2 localities.

POECILIIDAE

Acanthophaelus melanzonus Eigenmann, 1909

Ann. Carnegie Mus., 6(1): 51.

Holotype: FMNH 52717 (CM 1086). Guyana: Georgetown Trenches.

Paratypes: 6. Same locality.

POECILIIDAE

Acanthopthalmus mariae Inger and Chin, 1962

Fieldiana Zool., 45: 118.

Holotype: FMNH 68161. Malaysia: Sabah, tributary of Kinabatangan at Deramakot.

Paratypes: 112. Malaysia: Sabah, 2 localities.

COBITIDAE

Acanthopthalmus sandakanensis Inger and Chin, 1962

Fieldiana Zool., 45: 120.

Holotype: FMNH 68158. Malaysia: Sabah, Sepilok Forest Reserve.

Paratypes: 36. Malaysia: Sabah, 4 localities.

COBITIDAE

Acanthoplesiops hiatti Schultz, 1953

Bull. U.S. Natl. Mus., 202(1): 407.

Paratype. Marshall Islands: Bikini Atoll, Boro Island.

ACANTHOCLINIDAE

Acanthurus randalli Briggs and Caldwell, 1957

Bull. Fla. State Mus. Biol. Sci., 2(4): 43.

Paratype. Florida: Bay County near Panama City.

ACANTHURIDAE

Acanthurus triostegus marquesensis Schultz and Woods, 1948

J. Wash. Acad. Sci., 38(7): 250.

Paratypes: 5. Marquesas Islands: 2 localities.

ACANTHURIDAE

Acestridium discus Haseman, 1911

Ann. Carnegie Mus., 7(3–4): 319.

Holotype: FMNH 54339 (CM 2985). Brazil: Igarapé de Cachoeira Grande, near Manaus.

Paratype. Same locality.

LORICARIIDAE

- Acestrorhynchus altus** Menezes, 1969
Arq. Zool. (São Paulo), 18(1–2): 52.
Paratypes: 10. Brazil: Mato Grosso, Descalvados.
CHARACIDAE
- Acestrorhynchus guianensis** Menezes, 1969
Arq. Zool. (São Paulo), 18(1–2): 70.
Holotype: FMNH 74359. Guyana: Botanic Garden.
Paratypes: 7. Guyana: 4 localities.
CHARACIDAE
- Acestrorhynchus nasutus** Eigenmann, 1912
Mem. Carnegie Mus., 5: 411.
Holotype: FMNH 53475 (CM 1959). Guyana: Rockstone.
CHARACIDAE
- Acheilognathus barbatus** Nichols, 1926
Am. Mus. Nov., 214: 6.
Paratype. China: Anhwei, Ningkwo.
CYPRINIDAE
- Acheilognathus gracilis** Nichols, 1926
Am. Mus. Nov., 214: 5.
Paratype. China: Hunan, Tungting Lake.
CYPRINIDAE
- Acheilognathus tabira** Jordan and Thompson, 1914
Mem. Carnegie Mus., 6(4): 220.
Holotype: FMNH 57071 (CM 6007). Japan: Lake Biwa at Matsubara.
Paratypes: 41. Same locality.
CYPRINIDAE
- Achirus fluviatilis** Meek and Hildebrand, 1928
Field Mus. Nat. Hist. Zool. Ser., 15(3): 1002.
Paratypes: 40. Panama: 4 localities.
SOLEIDAE
- Acipenser oxyrhynchus desotoi** Vladykov, 1955
J. Fish. Res. Board Can., 12(5): 755.
Paratype. Mississippi: off Gautier at mouth of Singing River.
ACIPENSERIDAE
- Adioryx poco** Woods, 1965
Not. Nat. (Phila.), 377: 1.
Paratype. Grand Cayman Island: north side of Georgetown.
HOLOCENTRIDAE
- Adontosternarchus clarkae** Mago-Leccia, Lundberg, and Baskin, 1985
Nat. Hist. Mus. Los Ang. Cty. Contrib. Sci., 358: 14.
Paratypes: 6. Colombia and Ecuador: 2 localities.
APTERONOTIDAE
- Adontosternarchus devenanzii** Mago-Leccia, Lundberg, and Baskin, 1985
Nat. Hist. Mus. Los Ang. Cty. Contrib. Sci., 358: 11.
Paratypes: 12. Venezuela: Río Orinoco, north of Isla Tres Caños.
APTERONOTIDAE
- Aequidens awani** Haseman, 1911
Ann. Carnegie Mus., 7(3–4): 335.
Holotype: FMNH 54036 (CM 2576). Brazil: São Antonio de Guaporé.
Paratype. Same locality.
CICHLIDAE
- Aequidens azurifer** Fowler, 1911
Proc. Acad. Nat. Sci. Phila., 63: 515.
Paratype. Ecuador: near Bucay.
CICHLIDAE
- Aequidens chimantanus** Inger, 1956
Fieldiana Zool., 34(37): 437.
Holotype: FMNH 45702. Venezuela: Río Abácapa, west side of Chimantá-tepui.
Paratypes: 2. Same locality.
CICHLIDAE
- Aequidens duopunctata** Haseman, 1911
Ann. Carnegie Mus., 7(3–4): 338.
Holotype: FMNH 54033 (CM 2573). Brazil: Manaus.
Paratype. Same locality.
CICHLIDAE
- Aequidens guaporensis** Haseman, 1911
Ann. Carnegie Mus., 7(3–4): 335.
Holotype: FMNH 54035 (CM 2575). Brazil: São Antonio de Guaporé.
CICHLIDAE
- Aequidens mariae** Eigenmann, 1922
Mem. Carnegie Mus., 9(1): 240.
Paratypes: 3. Colombia: Río Negro at Villaviciencia.
CICHLIDAE

Aequidens plagiozonatus Kullander, 1984

Zool. Scr., 13(2): 155.

Paratype. Brazil: Descalvados.

CICHLIDAE

Aequidens potaroensis Eigenmann, 1912

Mem. Carnegie Mus., 5: 490.

Holotype: FMNH 53892 (CM 2407). Guyana: Amatuk.

Paratypes: 79. Guyana: 14 localities.

CICHLIDAE

Ageneiosus guianensis Eigenmann, 1912

Mem. Carnegie Mus., 5: 204.

Holotype: FMNH 53247 (CM 1712). Guyana: Wismar.

AGENEIOSIDAE

Ageneiosus madeirensis Fisher, 1917

Ann. Carnegie Mus., 11(3-4): 426.

Holotype: FMNH 58143 (CM 7143). Bolivia: San Joaquín.

Paratypes: 14. Same locality.

AGENEIOSIDAE

Ageneiosus marmoratus Eigenmann, 1912

Mem. Carnegie Mus., 5: 206.

Holotype: FMNH 53245 (CM 1710). Guyana: creek below Potaro Landing.

AGENEIOSIDAE

Agmus lyriformis Eigenmann, 1912

Mem. Carnegie Mus., 5: 128.

Holotype: FMNH 53120 (CM 1554). Guyana: Gluck Island.

ASPREDINIDAE

Agoniates anchovia Eigenmann, 1914

Indiana Univ. Stud., 19: 46.

Holotype: FMNH 56401 (CM 5216). Bolivia: Villa Bella.

Paratypes: 8. Same locality.

CHARACIDAE

Agosia adobe Jordan and Evermann, 1891

Bull. U.S. Fish Comm. 1889, 9: 36.

Paratypes: 3. Utah: Sevier River near Juab.

CYPRINIDAE

Agosia yarrowi Jordan and Evermann, 1891

Bull. U.S. Fish Comm. 1889, 9: 28.

Syntypes: 21, FMNH 1878, 6688. Colorado: Tomichi Creek, Gunnison.

CYPRINIDAE

Albula virgata Jordan and Jordan, 1922

Mem. Carnegie Mus., 10(1): 6.

Holotype: FMNH 55190 (CM 3896). Hawaiian Islands: Honolulu.

ALBULIDAE

Alburnops illecebrosus Girard, 1857

Proc. Acad. Nat. Sci. Phila., 1856: 194.

Lectoparatype. Arkansas: Arkansas River, Fort Smith.

Lectotype designated by Meek (In: Jordan, 1885, Proc. U.S. Natl. Mus., 8(500): 123).

CYPRINIDAE

Alectis breviventralis Wakiya, 1924

Ann. Carnegie Mus., 15(2-3): 216.

Holotype: FMNH 59411 (CM 7753). Japan: Kii. Paratype. Same locality.

CARANGIDAE

Alectis temmincki Wakiya, 1924

Ann. Carnegie Mus., 15(2-3): 215.

Holotype: FMNH 59410 (CM 7752). Japan: Bonin Islands.

CARANGIDAE

Alepideleotris tigris Herre, 1935

Field Mus. Nat. Hist. Zool. Ser., 18(12): 413.

Holotype: FMNH 17348. Ecuador: Islas Galápagos, Isla South Seymour.

ELEOTRIDAE

Alepocephalus owstoni Tanaka, 1908

J. Coll. Sci. Imp. Univ. Tokyo, 23(7): 27.

Holotype: FMNH 89529. Japan: Sagami Sea outside Okinose.

ALEPOCEPHALIDAE

Alepocephalus umbriceps Jordan and Thompson, 1914

Mem. Carnegie Mus., 6(4): 209.

Holotype: (CM 6030) MISSING. Japan: Aomori.

ALEPOCEPHALIDAE

Alestes langi Fowler, 1935

Ann. Transvaal Mus., 16(2): 258.

Paratypes: 3. Botswana: Thamalakane River at Maun.

CHARACIDAE

Alestes thamalakanensis Fowler, 1935

Ann. Transvaal Mus., 16(2): 257.

Paratypes: 5. Botswana: Thamalakane River at Maun.

CHARACIDAE

Alfaro acutiventralis Meek, 1912

Field Mus. Nat. Hist. Zool. Ser., 10(7): 72.
Holotype: FMNH 7679. Costa Rica: Guapilis.
Allotype and paratypes: 31. Same locality.
POECILIIDAE

Algansea aphanea Barbour and Miller, 1978

Misc. Publ. Mus. Zool. Univ. Mich., 155: 21.
Paratypes: 10. Mexico: Jalisco, Río Ayutla.
CYPRINIDAE

Algansea monticola Barbour and Contreras, 1968

Proc. Biol. Soc. Wash., 81: 101.
Paratypes: 2. Mexico: Zacatecas, Río Juchipila
near Jalpa.
CYPRINIDAE

Algansea rubescens Meek, 1902

Field Mus. Nat. Hist. Zool. Ser., 3(6): 78.
Holotype: FMNH 3653. Mexico: Río Santiago
at Ocotlán.
Paratypes: 8. Same locality.
CYPRINIDAE

Allanetta crenolepis Schultz, 1953

Bull. U.S. Natl. Mus., 202(1): 302.
Paratypes: 3. Philippine Islands: Tara Island.
ATHERINIDAE

Allolepis hollandi Jordan and Hubbs, 1925

Mem. Carnegie Mus., 10(2): 323.
Holotype: FMNH 58842 (CM 7952). Sea of Ja-
pan: near Fukui.
ZOARCIDAE

Alosa ohiensis Evermann, 1902

Rept. U.S. Fish Comm. 1901, 27: 277.
Paratype. Kentucky: Ohio River at Louisville.
CLUPEIDAE

Alticus novemmaculosus Snyder, 1908

Proc. U.S. Natl. Mus., 35(1635): 107.
Paratypes: 129. Japan: Ryukyu Islands, Oki-
nawa, Naha.
BLENNIIDAE

Ambassis lafa Jordan and Seale, 1906

Bull. Bur. Fish. 1905, 25: 253.
Paratypes: 16. Samoan Islands: Apia.
CENTROPOMIDAE

Ambassis vaivasensis Jordan and Seale, 1906

Bull. Bur. Fish. 1905, 25: 254.
Paratypes: 3. Samoan Islands: Vaivase River,
Apia.
CENTROPOMIDAE

Amblygobius myersi Herre, 1935

Field Mus. Nat. Hist. Zool. Ser., 18(12): 426.
Holotype: FMNH 17352. Solomon Islands: New
Georgia.
GOBIIDAE

Amblypharyngodon grandisquamis Jordan and
Starks, 1917

Ann. Carnegie Mus., 11: 438.
Holotype: FMNH 58964 (CM 8050). Sri Lanka:
Ceylon River at Colombo.
Paratypes: 17. Same locality.
CYPRINIDAE

Ameiurus australis Meek, 1904

Field Mus. Nat. Hist. Zool. Ser., 5: 13.
Holotype: FMNH 4474. Mexico: Tamaulipas,
Forlón.
Spelled *Amiurus* in Meek (1904).
ICTALURIDAE

Ameiurus mexicanus Meek, 1904

Field Mus. Nat. Hist. Zool. Ser., 5: 15.
Holotype: FMNH 4507. Mexico: San Luis Po-
tosí, Rascón.
Paratypes: 3. Mexico: San Luis Potosí, 2 local-
ities.
Spelled *Amiurus* in Meek (1904).
ICTALURIDAE

Amia sialis Jordan and Thompson, 1914

Mem. Carnegie Mus., 6: 246.
Holotype: FMNH 57084 (CM 6021). Japan: Su-
ruga Bay.
APOGONIDAE

Ammocryptocharax elegans Weitzman and Ka-
nazawa, 1976

Proc. Biol. Soc. Wash., 89(26): 331.
Paratypes: 7. Colombia: Río Vichada basin,
Caño Muco west of Puerto Gaitán.
CHARACIDAE

Anarchias yoshiae Kanazawa, 1952

Fieldiana Zool., 34(7): 75.
Holotype: FMNH 48729. Bermuda: ship's
channel.
MURAENIDAE

Anchoa hepsetus colonensis Hildebrand, 1943

Bull. Bingham Oceanogr. Collect. Yale Univ.,
8(2): 60.
Paratypes: 12. Jamaica.
ENGRAULIDAE

Anchoa lamprotaenia Hildebrand, 1943
Bull. Bingham Oceanogr. Collect. Yale Univ.,
8(2): 62.
Paratypes: 86. Gulf of Mexico and Caribbean
Sea: 7 localities.
ENGRAULIDAE

Anchoa mitchilli diaphana Hildebrand, 1943
Bull. Bingham Oceanogr. Collect. Yale Univ.,
8(2): 91.
Paratypes: 10. Louisiana: Grande Isle.
ENGRAULIDAE

Anchovia arenicola Meek and Hildebrand, 1923
Field Mus. Nat. Hist. Zool. Ser., 15(1): 201.
Paratypes: 122. Panama: 4 localities.
ENGRAULIDAE

Anchovia brevirostra Meek and Hildebrand, 1923
Field Mus. Nat. Hist. Zool. Ser., 15(1): 198.
Paratypes: 10. Panama: Canal Zone, Balboa Bay
at Balboa.
ENGRAULIDAE

Anchovia eigenmannia Meek and Hildebrand, 1923
Field Mus. Nat. Hist. Zool. Ser., 15(1): 205.
Paratypes: 7. Panama: Taboga Island, Balboa.
ENGRAULIDAE

Anchovia elongata Meek and Hildebrand, 1923
Field Mus. Nat. Hist. Zool. Ser., 15(1): 198.
Paratypes: 3. Panama: Canal Zone, Mindi Cut.
ENGRAULIDAE

Anchovia parva Meek and Hildebrand, 1923
Field Mus. Nat. Hist. Zool. Ser., 15(1): 202.
Paratypes: 24. Panama: 2 localities.
ENGRAULIDAE

Anchoviella belizensis Thomerson and Greenfield,
1975
Copeia, 1975(1): 50.
Holotype: FMNH 78363. Belize: Sibun River.
Paratypes: 35. Same locality.
ENGRAULIDAE

Anchoviella helleri Hubbs, 1921
Proc. Biol. Soc. Wash., 34: 47.
Holotype: FMNH 3332. Mexico: Baja California,
Bahía San Felipe.
Paratypes: 2. Same locality.
ENGRAULIDAE

Ancistrus lithurgicus Eigenmann, 1912
Mem. Carnegie Mus., 5: 241.
Holotype: FMNH 53091 (CM 1524). Guyana:
Crab Falls.
Paratypes: 5. Same locality.
LORICARIIDAE

Ancistrus melas Eigenmann, 1916
Ann. Carnegie Mus., 10(1-2): 83.
Holotype: FMNH 58339 (CM 7335). Colombia:
Condoto.
Paratype. Colombia: Raspadura.
LORICARIIDAE

Ancistrus planiceps Meek and Hildebrand, 1913
Field Mus. Nat. Hist. Zool. Ser., 10(8): 79.
Holotype: FMNH 7580. Panama: Río Tuyra,
Boca de Cupe.
Paratypes: 16. Panama: 6 localities.
LORICARIIDAE

Ancistrus spinosus Meek and Hildebrand, 1916
Field Mus. Nat. Hist. Zool. Ser., 10(15): 252.
Holotype: FMNH 8942. Panama: Río Calobre.
LORICARIIDAE

Ancylosetta antillarum Gutherz, 1966
Bull. Mar. Sci., 16(3): 449.
Paratypes: 2. Caribbean Sea: 2 localities.
BOTHIDAE

Ancylosetta cycloidea Tyler, 1959
Copeia, 1959(2): 140.
Paratypes: 9. Western Atlantic Ocean: 2 localities.
BOTHIDAE

Ancylosetta kumperae Tyler, 1959
Copeia, 1959(2): 143.
Paratypes: 2. Western Atlantic Ocean: 2 localities.
BOTHIDAE

Ancylosetta microctenus Gutherz, 1966
Bull. Mar. Sci., 16(3): 455.
Paratypes: 2. Caribbean Sea.
BOTHIDAE

Anduzedoras arleo Fernández-Yépez, 1968
Bol. Inst. Oceanogr. Univ. Oriente, 7(1): 29.
Holotype: FMNH 84068. Venezuela: Río
Auatana, east of Puerto Ayacucho.
DORADIDAE

Anduzedoras copei Fernández-Yépez, 1968
Bol. Inst. Oceanogr. Univ. Oriente, 7(1): 31.
Holotype: FMNH 84069. Venezuela: Lagoon at
side of Río Capanaparo.
DORADIDAE

Anisotremus williamsi Meek, 1905
Proc. Biol. Soc. Wash., 18: 241.
Holotype: FMNH 3350. Brazil: Santos.
Paratypes: 3. Same locality.
Misspelled *Anisotrema* in Meek (1905); *wil-*
liamsoni in Grey (1947, p. 116).
HAEMULIDAE

Anoplarchus purpureus archolepis Hubbs, 1926
Pap. Mich. Acad. Sci. Arts and Lett., 7: 375.
Paratypes: 34. California: 6 localities.
Misspelled *purpurascens* in Grey (1947, p.
118).
STICHAEIDAE

Anostomus plicatus Eigenmann, 1912
Mem. Carnegie Mus., 5: 296.
Holotype: FMNH 53393 (CM 1861). Guyana:
Crab Falls.
Paratypes: 3. Guyana: 3 localities.
ANOSTOMIDAE

Antennarius bermudensis Schultz, 1957
Proc. U.S. Natl. Mus., 107(3383): 98.
Paratype: Bermuda: Harrington Sound.
ANTENNARIIDAE

Antennarius pauciradiatus Schultz, 1957
Proc. U.S. Natl. Mus., 107(3383): 100.
Paratypes: 2. Florida: south end of Biscayne Bay.
ANTENNARIIDAE

Antennarius verrucosus Bean, 1906
Proc. Biol. Soc. Wash., 19: 31.
Holotype: FMNH 4853. Bermuda: Saint
George's Island.
ANTENNARIIDAE

Anthias conspicuus Heemstra, 1973
Copeia, 1973(2): 200.
Paratypes: 12. India: Arabian Sea, off Bombay.
SERRANIDAE

Anthias mooreanus Herre, 1935
Field Mus. Nat. Hist. Zool. Ser., 18(12): 405.
Holotype: FMNH 17325. Society Islands: Moo-
rea Island.
SERRANIDAE

Anthias woodsi Anderson and Heemstra, 1980
Copeia, 1980(1): 74.
Holotype: FMNH 70709. Straits of Florida: off
Dry Tortugas.
Paratypes: 2. Straits of Florida off Dry Tortugas:
2 localities.
SERRANIDAE

Antigonía combatia Berry and Rathjen, 1958
Q. J. Fla. Acad. Sci., 21(3): 255.
Paratypes: 52. Gulf of Mexico and off Cuba: 9
localities.
CAPROIDAE

Antigonía malayana Weber, 1913
Die Fische der Siboga-Expedition, Leiden, p.
299.
Syntype: FMNH 52432 (CM 689). Arafura Sea.
CAPROIDAE

Apareiodon hasemani Eigenmann, 1916
Ann. Carnegie Mus., 10(1–2): 75.
Holotype: FMNH 57615 (CM 6587). Brazil:
Pirapora.
Paratypes: 59. Brazil: 5 localities.
PARODONTIDAE

Apareiodon itapicuruensis Eigenmann and Henn,
1916
Ann. Carnegie Mus., 10(1–2): 72.
Holotype: FMNH 56988 (CM 5804). Brazil: Rio
Itapicuru basin, Rio Paiaia.
Paratypes: 7. Brazil: 2 localities.
PARODONTIDAE

Aparrius aurocingulus Herre, 1935
Field Mus. Nat. Hist. Zool. Ser., 18(12): 425.
Holotype: FMNH 17353. Fiji Islands: Ovalau
Island.
Paratypes: 2. Same locality.
GOBIIDAE

Aphyocharax anisitsi Eigenmann and Kennedy,
1903
Proc. Acad. Nat. Sci. Phila., 59: 517.
Paratypes: 2. Paraguay: Campo Grande.
CHARACIDAE

Aphyocharax erythrurus Eigenmann, 1912
Mem. Carnegie Mus., 5: 313.
Holotype: FMNH 53579 (CM 1879). Guyana:
Rockstone sand-bank.
Paratypes: 14. Guyana: 3 localities.
CHARACIDAE

- Aphyocharax melanotus** Eigenmann, 1912
Mem. Carnegie Mus., 5: 312.
Holotype: FMNH 53578 (CM 1877). Guyana:
Rockstone sand-bank.
CHARACIDAE
- Aphyocharax paraguayensis** Eigenmann, 1915
Mem. Carnegie Mus., 7(1): 33.
Holotype: FMNH 57922 (CM 6906). Brazil:
Cáceres.
Paratype. Same locality.
CHARACIDAE
- Aphyocheiroidon hemigrammus** Eigenmann, 1915
Mem. Carnegie Mus., 7(1): 59.
Holotype: FMNH 57819 (CM 6802). Brazil: Ja-
quara.
Paratypes: 25. Brazil: 3 localities.
CHARACIDAE
- Aphyodite grammica** Eigenmann, 1912
Mem. Carnegie Mus., 5: 314.
Holotype: FMNH 53408 (CM 1882). Guyana:
Konawaruk.
CHARACIDAE
- Apistogramma caetei** Kullander, 1980
Bonn. Zool. Monogr., 14: 76.
Holotype: FMNH 54164 (CM 2732). Brazil:
Pará, Igarapé in Bragança.
Paratypes: 2. Same locality.
CICHLIDAE
- Apistogramma inconspicua** Kullander, 1982
Zool. Scr., 11(4): 307.
Paratypes: 3. Brazil: São Luiz de Cáceres.
CICHLIDAE
- Apistogramma macmasteri** Kullander, 1979
Zool. Scr., 8: 70.
Paratype. Colombia: Villavicencio.
CICHLIDAE
- Aplatophis chauliodus** Böhlke, 1956
Not. Nat. (Phila.), 289: 3.
Holotype: FMNH 61772. Puerto Rico: Maya-
güez pier.
OPHICHTHIDAE
- Aplocheilus chobensis** Fowler, 1935
Ann. Transvaal Mus., 16(2): 276.
Paratypes: 4. Botswana: Chobe River at Ka-
bulabula.
APLOCHEILIDAE
- Aplocheilus luzonensis** Herre and Ablan, 1934
Philipp. J. Sci., 54(2): 275.
Paratypes: 4. Philippine Islands: Luzon, Ilocos
Norte, Solsona.
APLOCHEILIDAE
- Apogon abo** Herre, 1935
Field Mus. Nat. Hist. Zool. Ser., 18(12): 402.
Holotype: FMNH 17307. New Guinea: Sepik
River at Koragu.
Paratypes: 3. New Guinea: 2 localities.
APOGONIDAE
- Apogon beauforti** Weber, 1908
Nova Guinea, 5(2): 246.
Syntypes: 3, FMNH 52390 (CM 646). New
Guinea: Lake Sentani.
APOGONIDAE
- Apogon leptocaulus** Gilbert, 1972
Proc. Biol. Soc. Wash., 85(36): 421.
Paratype. Belize: Glover's Reef.
APOGONIDAE
- Apogon nigrofasciatus** Lachner, 1953
Bull. U.S. Natl. Mus., 202(1): 466.
Paratypes: 3. Marshall Islands: Bikini Atoll, 2
localities.
APOGONIDAE
- Apogon snyderi** Jordan and Evermann, 1903
Bull. U.S. Fish Comm. 1902, 22: 180.
Paratypes: 2. Hawaiian Islands: Hilo.
APOGONIDAE
- Apogon wichmanni** Weber, 1908
Nova Guinea, 5(2): 248.
Syntype: FMNH 52389 (CM 645). New Guinea:
Lake Sentani.
APOGONIDAE
- Apogonichthys melampodus** Blosser, 1909
Ann. Carnegie Mus., 6(1): 296.
Holotype: FMNH 53049 (CM 1474). Lesser An-
tilles: Saint Croix.
APOGONIDAE
- Apuredoras rivasi** Fernández-Yépez, 1950
Mem. Soc. Cienc. Nat. La Salle, 10(27): 195.
Paratypes: 7. Venezuela: Río Apure near San
Fernando.
DORADIDAE

- Araiophos gracilis** Grey, 1961
Pac. Sci., 15(3): 465.
Holotype: FMNH 73901. Hawaiian Islands: off Mauna Loa lava flow.
Paratypes: 8. Same locality.
STERNOPTYCHIDAE
- Archicheir minutus** Eigenmann, 1909
Ann. Carnegie Mus., 6(1): 46.
Holotype: FMNH 52786 (CM 1186). Guyana: Christianburg.
LEBIASINIDAE
- Archosargus oviceps** Ginsburg, 1952
J. Wash. Acad. Sci., 42(3): 94.
Paratypes: 2. Texas: Harbor Island.
SPARIDAE
- Areliscus hollandi** Jordan and Metz, 1913
Mem. Carnegie Mus., 6(1): 62.
Holotype: FMNH 55627 (CM 4369). Korea: Fusan.
CYNOGLOSSIDAE
- Argentina alicae** Cohen and Atsides, 1969
U.S. Fish Wildl. Serv. Fish. Bull., 68(1): 19.
Paratypes: 5. Pacific Ocean: off Peru.
ARGENTINIDAE
- Argentina brucei** Cohen and Atsides, 1969
U.S. Fish Wildl. Serv. Fish. Bull., 68(1): 31.
Paratypes: 20. Caribbean Sea: 7 localities.
ARGENTINIDAE
- Argentina georgei** Cohen and Atsides, 1969
U.S. Fish Wildl. Serv. Fish. Bull., 68(1): 27.
Paratypes: 3. Bahama Banks and Antilles: 3 localities.
ARGENTINIDAE
- Argentina stewarti** Cohen and Atsides, 1969
U.S. Fish Wildl. Serv. Fish. Bull., 68(1): 29.
Paratypes: 4. Antilles: 3 localities.
ARGENTINIDAE
- Arius kanganamanensis** Herre, 1935
Field Mus. Nat. Hist. Zool. Ser., 18(12): 387.
Holotype: FMNH 17194. New Guinea: Sepik River at Kanganaman.
ARIIDAE
- Arius nox** Herre, 1935
Field Mus. Nat. Hist. Zool. Ser., 18(12): 388.
Holotype: FMNH 17195. New Guinea: Sepik River at Nyaurangai.
Paratypes: 5. New Guinea: 2 localities.
ARIIDAE
- Arius solidus** Herre, 1935
Field Mus. Nat. Hist. Zool. Ser., 18(12): 385.
Holotype: FMNH 17201. New Guinea: Sepik River at Timbunke.
Paratypes: 9. New Guinea: Sepik River, 3 localities.
ARIIDAE
- Arius tuyra** Meek and Hildebrand, 1923
Field Mus. Nat. Hist. Zool. Ser., 15(1): 128.
Paratypes: 5. Panama: 2 localities.
ARIIDAE
- Asiphonichthys hemigrammus** Eigenmann, 1912
Mem. Carnegie Mus., 5: 403.
Holotype: FMNH 53665 (CM 2137). Guyana: Gluck Island.
CHARACIDAE
- Aspidoras maculosus** Nijssen and Isbrucker, 1976
Bijdr. Dierkd., 46(1): 119.
Holotype: FMNH 54810 (CM 3457). Brazil: Rio Paiaia.
Paratypes: 6. Brazil: 3 localities.
CALLICHTHYIDAE
- Astroblepus cyclopus santanderensis** Eigenmann, 1917
Proc. Am. Philos. Soc., 56(7): 675.
Lectotype: FMNH 58433 (CM 7430). Colombia: Quebrada de Guapota.
Lectoparatypes: 30. Colombia: 8 localities.
Lectotype designated by Henn (1928, p. 82).
ASTROBLEPIDAE
- Astroblepus frenatus** Eigenmann, 1917
Proc. Am. Philos. Soc., 56(7): 676.
Holotype: FMNH 58384 (CM 7380). Colombia: Quebrada de San Joaquín.
ASTROBLEPIDAE
- Astroblepus grixalvii micrescens** Eigenmann, 1917
Proc. Am. Philos. Soc., 56(7): 677.
Holotype: FMNH 58376 (CM 7372). Colombia: Quebrada de Agua Larga.
Paratypes: 41. Colombia: 9 localities.
ASTROBLEPIDAE
- Astroblepus latidens** Eigenmann, 1917
Proc. Am. Philos. Soc., 56(7): 674.
Holotype: FMNH 58366 (CM 7362). Colombia: Piperel.
Paratypes: 40. Colombia: 5 localities.
ASTROBLEPIDAE

Astroblepus pholeter Collette, 1962
Proc. Biol. Soc. Wash., 75: 311.
Paratype. Ecuador: Napo Province, cave in La-
tas.
ASTROBLEPIDAE

Astroblepus rosei Eigenmann, 1922
Mem. Carnegie Mus., 9(1): 55.
Paratype. Peru: Río Jequetepeque at Lllallán.
ASTROBLEPIDAE

Astronesthes ijimai Tanaka, 1908
J. Coll. Sci. Imp. Univ. Tokyo, 23(13): 9.
Syntypes (?): 10, FMNH 89100 (CM 4610). Ja-
pan: Sagami Sea.
These differ from material listed in original
publication in being a series of 10 col-
lected on 1 June 1907 (versus 5 from
May).
ASTRONESTHIDAE

Astronotus orbiculatus Haseman, 1911
Ann. Carnegie Mus., 7(3-4): 331.
Holotype: FMNH 53992 (CM 2515). Brazil:
Santarém.
Paratype. Same locality.
CICHLIDAE

Astyanax abramoides Eigenmann, 1909
Ann. Carnegie Mus., 6(1): 21.
Holotype: FMNH 52863 (CM 1028). Guyana:
Potaro River at Tumatumari.
Paratypes: 78. Guyana: 8 localities.
CHARACIDAE

Astyanax aeneus costaricensis Meek, 1914
Field Mus. Nat. Hist. Zool. Ser., 10(10): 105.
Syntypes: 203, FMNH 6849 and 7866-7873.
Costa Rica: 8 localities.
CHARACIDAE

Astyanax albeolus Eigenmann, 1908
Bull. Mus. Comp. Zool. Harv. Coll., 52(6):
97.
Holotype: FMNH 6241. Costa Rica: Río Ma-
chuca.
CHARACIDAE

Astyanax aurocaudatus Eigenmann, 1913
Indiana Univ. Stud., 18: 26.
Holotype: FMNH 56882 (CM 5162). Colombia:
Boquia.
Paratypes: 3. Same locality.
CHARACIDAE

Astyanax bimaculatus novae Eigenmann, 1911
Ann. Carnegie Mus., 8(1): 175.
Holotype (?): FMNH 54641 (CM 3278). Brazil:
Rio Sapon at Prazeres.
Paratypes: 23. Brazil: 2 localities.
One of the 11 specimens in the paratype lot
FMNH 54641 could be the illustrated
holotype.
CHARACIDAE

Astyanax bimaculatus paraguayensis Eigenmann,
1921
Mem. Mus. Comp. Zool. Harv. Coll., 43(3):
256.
Syntypes: 36, FMNH 54643-54648 (CM 3280-
3286). Bolivia, Brazil, and Paraguay: 7 lo-
calities.
Two lots CM 3274 and 3287 with 6 specimens
are MISSING.
CHARACIDAE

Astyanax daguae Eigenmann, 1913
Indiana Univ. Stud., 18: 23.
Holotype: FMNH 56251 (CM 5052). Colombia:
Córdova.
CHARACIDAE

Astyanax essequibensis Eigenmann, 1909
Ann. Carnegie Mus., 6(1): 17.
Holotype: FMNH 53519 (CM 1018). Guyana:
Tumatumari.
Paratypes: 80. Guyana: 5 localities.
CHARACIDAE

Astyanax grandis Meek and Hildebrand, 1912
Field Mus. Nat. Hist. Zool. Ser., 10(6): 67.
Holotype: FMNH 7571. Panama: Río Juan Díaz.
Paratypes: 196. Panama: 7 localities.
CHARACIDAE

Astyanax guaporensis Eigenmann, 1911
Ann. Carnegie Mus., 8(1): 176.
Holotype: FMNH 54709 (CM 3351). Brazil: Rio
Guaporé at Maciel.
Paratype. Same locality.
CHARACIDAE

Astyanax guianensis Eigenmann, 1909
Ann. Carnegie Mus., 6(1): 16.
Holotype: FMNH 53518 (CM 1013). Guyana:
Warraputa.
Paratypes: 45. Guyana: 3 localities.
CHARACIDAE

Astyanax gymnogonys Eigenmann, 1911
Ann. Carnegie Mus., 8(1): 179.
Holotype: FMNH 54707 (CM 3350). Brazil: Rio
Iguaçu at Porto União.
Paratype. Same locality.
CHARACIDAE

Astyanax hasemani Eigenmann, 1914
Indiana Univ. Stud., 19: 10.
Holotype: FMNH 56659 (CM 5476). Brazil:
Porto Alegre.
Paratypes: 6. Same locality.
CHARACIDAE

Astyanax heterurus Eigenmann and Wilson, 1914
In: Eigenmann, Henn, and Wilson, Indiana
Univ. Stud., 19: 11.
Holotype: FMNH 56577 (CM 5392). Colombia:
Río Truando.
CHARACIDAE

Astyanax kompi Hildebrand, 1938
Field Mus. Nat. Hist. Zool. Ser., 22(4): 260.
Paratypes: 6. Panama: Volcán, Laguna Gulnar
or Grande.
CHARACIDAE

Astyanax magdalenae Eigenmann and Henn, 1916
Ann. Carnegie Mus., 10(1–2): 89.
Holotype: FMNH 57006 (CM 5822). Colombia:
Girardot.
CHARACIDAE

Astyanax marionae Eigenmann, 1911
Ann. Carnegie Mus., 8(1): 175.
Holotype: FMNH 54711 (CM 3353). Brazil: near
São Luiz de Cáceres.
Paratypes: 8. Brazil: 2 localities.
CHARACIDAE

Astyanax metae Eigenmann, 1914
Indiana Univ. Stud., 19: 11.
Holotype: FMNH 56640 (CM 5457). Colombia:
Río Negro at Villavicencio.
Paratypes: 2. Same locality.
CHARACIDAE

Astyanax microlepis Eigenmann, 1913
Indiana Univ. Stud., 18: 24.
Holotype: FMNH 56209 (CM 5001). Colombia:
Piedra de Moler.
Paratypes: 843. Colombia: 5 localities.
CHARACIDAE

Astyanax mucronatus Eigenmann, 1909
Ann. Carnegie Mus., 6(1): 19.
Holotype: FMNH 53520 (CM 1025). Guyana:
Potaro River at Tumatumari.
Paratypes: 7. Guyana: 2 localities.
CHARACIDAE

Astyanax mutator Eigenmann, 1909
Ann. Carnegie Mus., 6(1): 18.
Holotype: FMNH 52688 (CM 1023). Guyana:
Potaro River at Savannah Landing.
Paratypes: 61. Guyana: 2 localities.
CHARACIDAE

Astyanax nasutus Meek, 1907
Field Mus. Nat. Hist. Zool. Ser., 7(4): 108.
Holotype: FMNH 5909. Nicaragua: Lago Ma-
nagua at Managua.
Paratypes: 15. Nicaragua: 2 localities.
CHARACIDAE

Astyanax notemigonoides Fowler, 1911
Proc. Acad. Nat. Sci. Phila., 63: 506.
Paratype. Ecuador: near Bucay.
CHARACIDAE

Astyanax paranahybae Eigenmann, 1911
Ann. Carnegie Mus., 8(1): 177.
Holotype: FMNH 54714 (CM 3356). Brazil: Rio
Paranahyba.
CHARACIDAE

Astyanax potaroensis Eigenmann, 1909
Ann. Carnegie Mus., 6(1): 22.
Holotype: FMNH 52695 (CM 1037). Guyana:
Potaro River at Amatuk Cataract.
Paratypes: 7. Guyana: 3 localities.
CHARACIDAE

Astyanax regani Meek, 1909
Field Mus. Nat. Hist. Zool. Ser., 7(7): 207.
Holotype: FMNH 6257. Costa Rica: Las Cañas.
CHARACIDAE

Astyanax ribeirae Eigenmann, 1911
Ann. Carnegie Mus., 8(1): 177.
Holotype: FMNH 54725 (CM 3368). Brazil:
Xiririca.
Paratypes: 103. Brazil: 4 localities.
CHARACIDAE

- Astyanax robustus** Meek, 1912
Field Mus. Nat. Hist. Zool. Ser., 10(7): 69.
Holotype: FMNH 7682. Costa Rica: Virginia River.
Paratypes (?): 116. Costa Rica.
See comment in Grey (1947, pp. 122–123); status of various paratypes uncertain.
CHARACIDAE
- Astyanax ruberrimus** Eigenmann, 1913
Indiana Univ. Stud., 18: 25.
Holotype: FMNH 56122 (CM 4912). Colombia: Istmina.
Paratypes: 315. Colombia: 4 localities.
CHARACIDAE
- Astyanax scabripinnis paranae** Eigenmann, 1914
Indiana Univ. Stud., 20: 47.
Paratypes: 307. Brazil: 10 localities.
CHARACIDAE
- Astyanax scierus** Fowler, 1911
Proc. Acad. Nat. Sci. Phila., 63: 509.
Paratype: Ecuador: near Bucay.
CHARACIDAE
- Atherina evermanni** Eigenmann, 1903
Bull. U.S. Fish Comm. 1902, 22: 228.
Paratypes: 5. Cuba: 2 localities.
ATHERINIDAE
- Atherina insularum** Jordan and Evermann, 1903
Bull. U.S. Fish Comm. 1902, 22: 170.
Paratype: Hawaiian Islands: Honolulu.
ATHERINIDAE
- Atherina ovalaua** Herre, 1935
Field Mus. Nat. Hist. Zool. Ser., 18(12): 401.
Holotype: FMNH 17272. Fiji Islands: Ovalau Island.
Paratypes: 33. Fiji Islands: 2 localities.
ATHERINIDAE
- Atherinichthys nouhuysi** Weber, 1910
Notes Leiden Mus., 32: 229.
Syntypes: 2, FMNH 52500 (CM 758). New Guinea: Lorentz River, Alkmaar.
ATHERINIDAE
- Atherinops affinis littoralis** Hubbs, 1918
Bull. Am. Mus. Nat. Hist., 38(13): 426.
Holotype: FMNH 2064. California: San Diego Bay, North Island.
Paratypes: 5. California: 3 localities.
ATHERINIDAE
- Atherinops insularum cedroscensis** Hubbs, 1918
Bull. Am. Mus. Nat. Hist., 38(13): 434.
Paratypes: 2. Mexico: Baja California, Islas Cedros.
ATHERINIDAE
- Atherinops insularum guadalupae** Hubbs, 1918
Bull. Am. Mus. Nat. Hist., 38(13): 435.
Paratypes: 2. Mexico: Baja California, Isla Guadalupe.
ATHERINIDAE
- Atherion elymus aphrozoicus** Schultz, 1953
Bull. U.S. Natl. Mus., 202(1): 295.
Paratype. Philippine Islands: Tara Island.
ATHERINIDAE
- Atherion elymus freyi** Schultz, 1953
Bull. U.S. Natl. Mus., 202(1): 292.
Paratypes: 2. Guam: 2 localities.
ATHERINIDAE
- Atuona tricuspidata** Herre, 1935
Field Mus. Nat. Hist. Zool. Ser., 18(12): 429.
Holotype: FMNH 17356. Marquesas Islands: Hiva Oa Island, Autona.
Paratypes: 4. Same locality.
GOBIIDAE
- Auchenipterus brevior** Eigenmann, 1912
Mem. Carnegie Mus., 5: 202.
Holotype: FMNH 53249 (CM 1715). Guyana: Tumatumari.
Paratypes: 6. Same locality.
AUCHENIPTERIDAE
- Auchenipterus demerarae** Eigenmann, 1912
Mem. Carnegie Mus., 5: 202.
Holotype: FMNH 53248 (CM 1714). Guyana: Wismar.
Paratype. Guyana: Malali.
AUCHENIPTERIDAE
- Awaous decemlineatus** Eigenmann, 1917
Proc. Am. Philos. Soc., 56(7): 686.
Holotype: FMNH 58476 (CM 7478). Colombia: Quibdo.
Paratypes: 9. Colombia: 2 localities.
GOBIIDAE
- Aztecula mexicana** Meek, 1902
Field Mus. Nat. Hist. Zool. Ser., 3(6): 81.
Holotype: FMNH 3606. Mexico: Río Moctezuma at San Juan del Río.
Paratypes: 220. Same locality.
CYPRINIDAE

Barbatula toni posteroventralis Nichols, 1925

Am. Mus. Nov., 171: 4.

Paratypes: 3. China: Shansi, Kwei-hwan.

COBITIDAE

Barbatula yarkandensis sellaefer Nichols, 1925

Am. Mus. Nov., 171: 4.

Paratypes: 3. China: Shansi, Chin-ssu.

COBITIDAE

Barbuligobius boehlkei Lachner and McKinney, 1974

Copeia, 1974(4): 871.

Paratype. Sri Lanka: Galle.

GOBIIDAE

Barbus akeleyi Hubbs, 1918

Field Mus. Nat. Hist. Zool. Ser., 12(2): 13.

Holotype: FMNH 6106. Kenya: Athi River.

Paratypes: 12. Same locality.

CYPRINIDAE

Barbus anoplus Weber, 1897

Zool. Jahrb., 10: 151.

Syntypes: 3, FMNH 52397 (CM 653). South

Africa: Buffels River near Laingsburg.

CYPRINIDAE

Barbus athi Hubbs, 1918

Field Mus. Nat. Hist. Zool. Ser., 12(2): 10.

Holotype: FMNH 6108. Kenya: Athi River.

Paratypes: 63. Same locality.

CYPRINIDAE

Barbus bantolanensis Day, 1914

Philipp. J. Sci., 9(2): 188.

Paratypes: 12. Philippine Islands: Palawan Island, Lake Manguao.

CYPRINIDAE

Barbus bifrenatus Fowler, 1935

Ann. Transvaal Mus., 16(2): 266.

Paratypes: 10. Botswana: Tsotsoroga Pan.

CYPRINIDAE

Barbus carens Boulenger, 1912

Ann. Mus. Congo Belge Ser. Quarto Zool., Ser. 1, 2(3): 15.

Syntype: FMNH 56148 (CM 4939). Zaïre: Le-buzi River at Kuka Muno.

CYPRINIDAE

Barbus fitsimensi Fowler, 1935

Ann. Transvaal Mus., 16(2): 267.

Paratypes: 15. Botswana: 2 localities.

CYPRINIDAE

Barbus helleri Hubbs, 1918

Field Mus. Nat. Hist. Zool. Ser., 12(2): 12.

Holotype: FMNH 6105. Kenya: Athi River.

Paratypes: 103. Same locality.

CYPRINIDAE

Barbus oatesii Boulenger, 1893

Ann. Mag. Nat. Hist., Ser. 6, 12: 201.

Syntypes: 2, FMNH 5708. Burma: Nampandet.

CYPRINIDAE

Barbus tsotsorogensis Fowler, 1935

Ann. Transvaal Mus., 16(2): 265.

Paratypes: 23. Botswana: 2 localities.

CYPRINIDAE

Bathylagus tetranema Bailey and Stewart, 1984

Misc. Publ. Mus. Zool. Univ. Mich., 168:11.

Paratypes: 2. Zambia: Lake Tanganyika near Mpulungu.

BAGRIDAE

Bathylupea schroederi Dick, 1962

Breviora, 167: 1.

Paratypes: 4. Western Caribbean Sea: 3 localities.

BATHYCLUPEIDAE

Bathylagus greyae Cohen, 1958

Breviora, 98: 5.

Holotype: FMNH 49730. Western North Atlantic: near Bermuda.

BATHYLAGIDAE

Bathypterois bigelowi Mead, 1958

J. Wash. Acad. Sci., 48(11): 366.

Holotype: FMNH 64435. Caribbean Sea: off Nicaragua.

Paratypes: 42. Caribbean Sea: 12 localities.

CHLOROPHTHALMIDAE

Bathypterois dubius Vaillant, 1888

Exp. Sci. Trav. et Talis. . . Poissons, Paris, p. 124 (not seen).

Paratype. Off Mauritania: Banc d'Arguin.

CHLOROPHTHALMIDAE

Bathypterois pectoralis Garman, 1899

Mem. Mus. Comp. Zool. Harv. Coll., 24: 257.

Syntype: FMNH 33730. Pacific Ocean: off Colombia.

CHLOROPHTHALMIDAE

- Bathytyphlops marionae** Mead, 1958
J. Wash. Acad. Sci., 48(11): 370.
Holotype: FMNH 64439. Caribbean Sea.
CHLOROPHTHALMIDAE
- Batrachoides waltersi** Collette and Russo, 1981
Bull. Mar. Sci., 31(2): 208.
Paratypes: 3. El Salvador: Bahía Jiquilisco, La Venadona.
BATRACHOIDIDAE
- Beirabarus palustris** Herre, 1936
Proc. Biol. Soc. Wash., 49: 100.
Paratypes: 12. Mozambique: near Beira.
CYPRINIDAE
- Bellator ribeiroi** Miller, 1965
Q. J. Fla. Acad. Sci., 28(3): 259.
Paratypes: 2. Western Atlantic Ocean: off Colombia.
TRIGLIDAE
- Belligobio eristigma** Jordan and Hubbs, 1925
Mem. Carnegie Mus., 10(2): 173.
Holotype: FMNH 58714 (CM 7820). Japan: near Okayama.
CYPRINIDAE
- Belone svetovidovi** Knudsen and Wolff, 1970
Atl. Rep., 11: 21.
Paratypes: 7. Tunisia: Gabes.
BELONIDAE
- Bembrops greyi** Poll, 1959
Résult. Sci. Exp. Oceanogr. Belge . . . (1948–1949), 4(3B): 40.
Paratype. South Atlantic: off Angola, Ambri-zete.
PERCOPHIDAE
- Bivibranchia protractila** Eigenmann, 1912
Mem. Carnegie Mus., 5: 259.
Holotype: FMNH 53576 (CM 1873). Guyana: Bartica sand-bank.
Paratypes: 8. Guyana: 3 localities.
HEMIODONTIDAE
- Blenniulus proteus** Krejsa, 1960
Copeia, 1960(4): 326.
Paratypes: 2. Mexico: Islas Revilla Gigedo southeast of Isla San Benedictino.
BLENNIIDAE
- Blennius nicholsi** Tavalga, 1954
Copeia, 1954(2): 135.
Paratypes: 2. Florida: 2 localities.
BLENNIIDAE
- Bleptonema amazonae** Eigenmann, 1914
Indiana Univ. Stud., 20: 44.
Holotype: FMNH 56679 (CM 5497). Brazil: Santarém.
Paratypes: 2. Same locality.
CHARACIDAE
- Bleptonema paraguayensis** Eigenmann, 1914
Indiana Univ. Stud., 20: 44.
Holotype: FMNH 56681 (CM 5499). Paraguay: Corumbá.
Paratypes: 4. Same locality.
CHARACIDAE
- Bodianus stellatus** Blosser, 1909
Ann. Carnegie Mus., 6(1): 297.
Holotype: FMNH 53048 (CM 1473). Lesser Antilles: Saint Croix Island.
SERRANIDAE
- Bombonia uxorius** Herre, 1935
Field Mus. Nat. Hist. Zool. Ser., 18(12): 395.
Holotype: FMNH 17493. Indonesia: Waigiu Island.
SYNGNATHIDAE
- Boroda malua** Herre, 1935
Field Mus. Nat. Hist. Zool. Ser., 18(12): 414.
Holotype: FMNH 17349. New Guinea: Sepik River at Malu.
Paratypes: 2. Same locality.
ELEOTRIDAE
- Bostrychus zonatus** Weber, 1908
Nova Guinea, 5(2): 259.
Syntype: FMNH 52457 (CM 715). New Guinea: Merauke River.
ELEOTRIDAE
- Botia purpurea** Nichols, 1925
Am. Mus. Nov., 177: 4.
Paratype. China: Hunan, Tungting Lake.
COBITIDAE
- Brachirus bellus** Jordan and Hubbs, 1925
Mem. Carnegie Mus., 10(2): 274.
Holotype: FMNH 58783 (CM 7894). Japan: Misaki.
SCORPAENIDAE

- Brachychalcinus guianensis** Boeseman, 1952
Zool. Meded. (Leiden), 31(26): 303.
Syntypes: 6, FMNH 51535 and 59525. Guyana:
New River drainage, Itabu Creek.
CHARACIDAE
- Brachyglanis frenata** Eigenmann, 1912
Mem. Carnegie Mus., 5: 156.
Holotype: (CM 1670) MISSING. Guyana:
Amatuk.
PIMELODIDAE
- Brachyglanis melas** Eigenmann, 1912
Mem. Carnegie Mus., 5: 157.
Holotype: FMNH 53217 (CM 1672). Guyana:
Crab Falls.
Paratypes: 4. Guyana: 3 localities.
PIMELODIDAE
- Brachyglanis phalacra** Eigenmann, 1912
Mem. Carnegie Mus., 5: 157.
Holotype: FMNH 53216 (CM 1671). Guyana:
Amatuk.
PIMELODIDAE
- Brachygobius kabiliensis** Inger, 1958
Fieldiana Zool., 39(14): 110.
Holotype: FMNH 47991. Malaysia: Sabah, Ka-
bili River.
Paratypes: 6. Malaysia: Sabah, 2 localities.
GOBIIDAE
- Brachygobius sabanus** Inger, 1958
Fieldiana Zool., 39(14): 113.
Holotype: FMNH 47990. Malaysia: Sabah, La-
mag.
Paratypes: 21. Malaysia: Sabah, 3 localities.
GOBIIDAE
- Brachygobius xanthomelas** Herre, 1937
Bull. Raffles Mus., 13: 43.
Paratype. Malaysia: Johore, Mawai district.
GOBIIDAE
- Bramocharax elongatus** Meek, 1907
Field Mus. Nat. Hist. Zool. Ser., 7(4): 110.
Holotype: FMNH 5922. Nicaragua: Lago Ma-
nagua at Managua.
Paratypes: 26. Nicaragua: 3 localities.
See comment in Grey (1947, p. 126); status
of various paratypes uncertain.
CHARACIDAE
- Branchiostoma bermudae** Hubbs, 1922
Occas. Pap. Mus. Zool. Univ. Mich., 105: 9.
Paratypes: 71. Bermuda: The Flatts.
BRANCHIOSTOMATIDAE
- Branchiostoma floridae** Hubbs, 1922
Occas. Pap. Mus. Zool. Univ. Mich., 105: 7.
Paratypes: 7. Florida: Tampa Bay.
BRANCHIOSTOMATIDAE
- Bryconamericus alfredeae** Eigenmann, 1927
Mem. Mus. Comp. Zool. Harv. Coll., 43(4):
394.
Paratypes: 5. Peru: Río Urubamba at Santa Ana.
CHARACIDAE
- Bryconamericus alpha** Eigenmann, 1914
In: Eigenmann, Henn, and Wilson, Indiana
Univ. Stud., 19: 7.
Holotype: FMNH 56646 (CM 5463). Colombia:
Villavicencio.
Paratypes: 2. Same locality.
CHARACIDAE
- Bryconamericus beta** Eigenmann, 1914
In: Eigenmann, Henn, and Wilson, Indiana
Univ. Stud., 19: 7.
Holotype: FMNH 56648 (CM 5465). Colombia:
Villavicencio.
Paratypes: 4. Same locality.
CHARACIDAE
- Bryconamericus cascajalensis** Meek and Hilde-
brand, 1916
Field Mus. Nat. Hist. Zool. Ser., 10(15): 284.
Holotype: FMNH 8946. Panama: Río Cascajal
at Porto Bello.
Paratypes: 24. Same locality.
CHARACIDAE
- Bryconamericus caucanus** Eigenmann, 1913
Indiana Univ. Stud., 18: 17.
Holotype: FMNH 56229 (CM 5031). Colombia:
Piedra de Moler.
Paratypes: 106. Colombia: 6 localities.
CHARACIDAE
- Bryconamericus chocoensis** Eigenmann, 1913
Indiana Univ. Stud., 18: 14.
Holotype: FMNH 56235 (CM 5036). Colombia:
Istmina.
Paratypes: 270. Colombia: 6 localities.
CHARACIDAE

Bryconamericus cismontanus Eigenmann, 1914
In: Eigenmann, Henn, and Wilson, Indiana Univ. Stud., 19: 4.
Holotype: FMNH 56642 (CM 5459). Colombia: Villavicencio.
CHARACIDAE

Bryconamericus conventus Eigenmann, 1913
Indiana Univ. Stud., 18: 13.
Holotype: FMNH 56261 (CM 5060). Colombia: Soplaviento.
Paratypes: 57. Same locality.
CHARACIDAE

Bryconamericus decurrens Eigenmann, 1913
Indiana Univ. Stud., 18: 20.
Holotype: FMNH 56255 (CM 5055). Colombia: Soplaviento.
CHARACIDAE

Bryconamericus dentatus Eigenmann, 1913
Indiana Univ. Stud., 18: 19.
Holotype: FMNH 56253 (CM 5054). Colombia: Piedra de Moler.
Paratypes: 21. Colombia: 3 localities.
CHARACIDAE

Bryconamericus deuterodonoides Eigenmann, 1914
In: Eigenmann, Henn, and Wilson, Indiana Univ. Stud., 19: 5.
Holotype: FMNH 56644 (CM 5461). Colombia: Río Negro at Villavicencio.
Paratypes: 6. Same locality.
CHARACIDAE

Bryconamericus deuterodonoides caudovitatus Inger, 1956
Fieldiana Zool., 34(37): 435.
Holotype: FMNH 45706. Venezuela: Río Abácapa, west side of Chimantá-tepui.
Paratypes: 12. Same locality.
CHARACIDAE

Bryconamericus diquensis Eigenmann, 1913
Indiana Univ. Stud., 18: 14.
Holotype: FMNH 56272 (CM 5072). Colombia: Soplaviento.
Paratypes: 11. Same locality.
CHARACIDAE

Bryconamericus grosvenori Eigenmann, 1927
Mem. Mus. Comp. Zool. Harv. Coll., 43(4): 365.
Paratypes: 3. Peru: 2 localities.
CHARACIDAE

Bryconamericus henni Eigenmann, 1914
In: Eigenmann, Henn, and Wilson, Indiana Univ. Stud., 19: 6.
Holotype: FMNH 56594 (CM 5410). Ecuador: Vines.
Paratypes: 10. Ecuador: 2 localities.
CHARACIDAE

Bryconamericus hyphesson Eigenmann, 1909
Ann. Carnegie Mus., 6(1): 32.
Holotype: FMNH 52708 (CM 1070). Guyana: Tumatumari.
Paratypes: 3. Same locality.
CHARACIDAE

Bryconamericus magdalenensis Eigenmann, 1913
Indiana Univ. Stud., 18: 14.
Holotype: FMNH 56263 (CM 5063). Colombia: Girardot.
Paratypes: 675. Colombia: 8 localities.
CHARACIDAE

Bryconamericus novae Eigenmann and Henn, 1914
Indiana Univ. Stud., 24: 234.
Holotype: FMNH 54905 (CM 3568). Brazil: Rio Novo near Piabana.
Paratypes: 9. Same locality.
CHARACIDAE

Bryconamericus ortholepis Eigenmann, 1913
Indiana Univ. Stud., 18: 15.
Holotype: (CM 5088) MISSING. Colombia: Boca de Raspadura.
CHARACIDAE

Bryconamericus osgoodi Eigenmann and Allen, 1942
Fishes of Western South America, Lexington, Kentucky, p. 225.
Paratypes: 3. Peru: Moyobamba.
CHARACIDAE

Bryconamericus pachacuti Eigenmann, 1927
Mem. Mus. Comp. Zool. Harv. Coll., 43(4): 376.
Paratypes: 12. Peru: 2 localities.
CHARACIDAE

Bryconamericus peruanus ricae Eigenmann, 1908
Bull. Mus. Comp. Zool. Harv. Coll., 52(6): 106.
Lectotype: FMNH 59523. Costa Rica: Chitaria.
Lectoparatypes: 2. Same locality.
Lectotype designated by Grey (1947, pp. 127–128).
CHARACIDAE

Bryconamericus scopiferus Eigenmann, 1913

Indiana Univ. Stud., 18: 16.

Holotype: FMNH 56224 (CM 5026). Colombia:
Río San Juan at Itsmina.

Paratypes: 122. Colombia: 3 localities.

CHARACIDAE

Bryconamericus scopiferus guaytarae Eigenmann
and Henn, 1914

In: Eigenmann, Henn, and Wilson, Indiana
Univ. Stud., 19: 7.

Holotype: FMNH 56657 (CM 5474). Colombia:
Río Patia at mouth of Río Guáytara.

Paratypes: 122. Colombia: 3 localities.

CHARACIDAE

Bryconamericus tolimae Eigenmann, 1913

Indiana Univ. Stud., 18: 18.

Holotype: FMNH 56257 (CM 5057). Colombia:
Ibagué.

Paratypes: 7. Same locality.

CHARACIDAE

Bryconamericus zeteki Hildebrand, 1938

Field Mus. Nat. Hist. Zool. Ser., 22(4): 265.

Paratypes: 8. Panama: Pacific slope, creek in El
Valle.

CHARACIDAE

Brycon argenteus Meek and Hildebrand, 1913

Field Mus. Nat. Hist. Zool. Ser., 10(8): 84.

Holotype: FMNH 7588. Panama: Río Aruza at
Aruza.

Paratypes: 168. Panama: 11 localities.

CHARACIDAE

Brycon behrae Hildebrand, 1938

Field Mus. Nat. Hist. Zool. Ser., 22(4): 278.

Paratype. Panama: Río Chiriquí del Tire above
Caldera.

CHARACIDAE

Brycon henni Eigenmann, 1913

Indiana Univ. Stud., 18: 26.

Holotype: FMNH 56384 (CM 5152). Colombia:
Caldas.

Paratypes: 102. Colombia: 6 localities.

CHARACIDAE

Brycon obscurus Hildebrand, 1938

Field Mus. Nat. Hist. Zool. Ser., 22(4): 283.

Paratypes: 2. Panama: Pacific slope, creek in El
Valle.

CHARACIDAE

Brycon petrosus Meek and Hildebrand, 1913

Field Mus. Nat. Hist. Zool. Ser., 10(8): 84.

Holotype: FMNH 7589. Panama: Upper
Chagres.

Paratypes: 89. Panama: 10 localities.

CHARACIDAE

Brycon siebenthalae Eigenmann, 1912

Mem. Carnegie Mus., 5: 372.

Holotype: FMNH 53353 (CM 1819). Guyana:
Aruka River, Mud Creek.

CHARACIDAE

Bunocephalus amaurus Eigenmann, 1912

Mem. Carnegie Mus., 5: 126.

Holotype: FMNH 53121 (CM 1555). Guyana:
Konawaruk.

ASPREDINIDAE

Bunocephalus chamaizelus Eigenmann, 1912

Mem. Carnegie Mus., 5: 127.

Holotype: FMNH 53122 (CM 1556). Guyana:
Erukin.

Paratypes: 7. Guyana: 3 localities.

ASPREDINIDAE

Bunocephalus colombianus Eigenmann, 1912

Indiana Univ. Stud., 16: 10.

Holotype: FMNH 56038 (CM 4828). Colombia:
Raspadura.

ASPREDINIDAE

Bunocephalus depressus Haseman, 1911

Ann. Carnegie Mus., 7(3–4): 319.

Holotype: FMNH 54338 (CM 2984). Bolivia:
Río Machupo near San Joaquín.

ASPREDINIDAE

Calamopteryx jeb Cohen, 1973

Proc. Biol. Soc. Wash., 86(28): 343.

Holotype: FMNH 71603. Ecuador: Islas Galá-
pagos, Tagus Cove.

Paratypes: 24. Same locality.

OPHIDIIDAE

Callechelys bilinearis Kanazawa, 1952

Fieldiana Zool., 34(7): 72.

Holotype: FMNH 48973. Bermuda: Cooper's
Island and Saint David's.

OPHICHTHIDAE

Callionymus distethommatus Fowler, 1941

Proc. U.S. Natl. Mus., 90(3106): 18.

Paratypes: 2. Philippine Islands: Port Matalvi
off western Luzon.

CALLIONYMIDAE

Callionymus hudsoni Fowler, 1941

Proc. U.S. Natl. Mus., 90(3106): 8.

Paratype. Philippine Islands: Mantacao Island.

CALLIONYMIDAE

Callionymus tenuis Fricke, 1981

Ann. Mus. Civ. Stor. Nat. "Giacomo Doria"

Genova, 83: 387.

Holotype: FMNH 78916. Maldives Archipelago: South Nilandu Atoll.

CALLIONYMIDAE

Callionymus wilburi Herre, 1935

Field Mus. Nat. Hist. Zool. Ser., 18(12): 431.

Holotype: FMNH 17491. Indonesia: Waigiu Island.

Paratype. Same locality.

CALLIONYMIDAE

Calliurichthys astrinius Jordan and Jordan, 1922

Mem. Carnegie Mus., 10(1): 80.

Holotype: FMNH 55207 (CM 3903). Hawaiian Islands: Honolulu Market.

CALLIONYMIDAE

Calliurichthys zanectes Jordan and Jordan, 1922

Mem. Carnegie Mus., 10(1): 81.

Holotype: FMNH 55208 (CM 3904). Hawaiian Islands: Honolulu Market.

CALLIONYMIDAE

Callogobius ocellatus Herre, 1935

Field Mus. Nat. Hist. Zool. Ser., 18(12): 422.

Holotype: FMNH 17363. Fiji Islands: Ovalau Island.

GOBIIDAE

Calymmichthys xenicus Jordan and Thompson, 1914

Mem. Carnegie Mus., 6: 296.

Holotype: FMNH 57089 (CM 6027). Japan: Sagami Bay.

CALLIONYMIDAE

Caranx formosanus Jordan and Snyder, 1909

Mem. Carnegie Mus., 4(2): 38.

Holotype: FMNH 55364 (CM 412). Taiwan: Takao.

CARANGIDAE

Caranx ishikawai Wakiya, 1924

Ann. Carnegie Mus., 15(2-3): 193.

Holotype: FMNH 59390 (CM 7734). Japan: Bonin Islands.

CARANGIDAE

Caranx miyakamii Wakiya, 1924

Ann. Carnegie Mus., 15(2-3): 201.

Holotype: FMNH 59407 (CM 7742). Taiwan.

CARANGIDAE

Caranx oshimai Wakiya, 1924

Ann. Carnegie Mus., 15(2-3): 189.

Holotype: FMNH 59492 (CM 7731). Taiwan.

CARANGIDAE

Caranx rastrosus Jordan and Snyder, 1909

Mem. Carnegie Mus., 4(2): 37.

Holotype: FMNH 55363 (CM 411). Taiwan: Takao.

CARANGIDAE

Caranx tanakai Wakiya, 1924

Ann. Carnegie Mus., 15(2-3): 177.

Paratype. Japan: Kii.

CARANGIDAE

Caranx uii Wakiya, 1924

Ann. Carnegie Mus., 15(2-3): 174.

Holotype: FMNH 59494 (CM 7720). Japan: Kii.

Paratype. Same locality.

CARANGIDAE

Careproctus burkei Jordan and Thompson, 1914

Mem. Carnegie Mus., 6(4): 281.

Holotype: FMNH 57515 (CM 6457). Japan: Yokohama Market.

CYCLOPTERIDAE

Careproctus gilberti Jordan and Thompson, 1914

Mem. Carnegie Mus., 6(4): 282.

Holotype: FMNH 57514 (CM 6456). Japan: Sagami Bay, Misaki.

CYCLOPTERIDAE

Carpiodes elongatus Meek, 1904

Field Mus. Nat. Hist. Zool. Ser., 5: 28.

Holotype: FMNH 4425. Mexico: Nuevo León, Linares.

Paratypes: 13. Mexico: 3 localities.

CATOSTOMIDAE

Carpiodes labiosus Meek, 1904

Field Mus. Nat. Hist. Zool. Ser., 5: 29.

Holotype: FMNH 4492. Mexico: San Luis Potosí, Valles.

Paratype. Same locality.

CATOSTOMIDAE

- Carpiodes microstomus*** Meek, 1904
Field Mus. Nat. Hist. Zool. Ser., 5: 27.
Holotype: FMNH 3542. Mexico: Río Noavaco at Santa Rosalia.
Paratypes: 18. Mexico: 2 localities.
CATOSTOMIDAE
- Cascadura maculocephala*** Ellis, 1913
Ann. Carnegie Mus., 8(3–4): 387.
Holotype: FMNH 54878 (CM 3539). Brazil: Rio Uruguay at Uruguayana.
CALLICHTHYIDAE
- Catostomus conchos*** Meek, 1902
Field Mus. Nat. Hist. Zool. Ser., 3(6): 75.
Holotype: FMNH 3557. Mexico: Río Conchos at Jiménez.
Paratypes: 8. Same locality.
CATOSTOMIDAE
- Catostomus sonorensis*** Meek, 1902
Field Mus. Nat. Hist. Zool. Ser., 3(6): 75.
Holotype: FMNH 3564. Mexico: Río Paphigochic at Miñaca.
Paratypes: 21. Same locality.
CATOSTOMIDAE
- Centriscus capito*** Oshima, 1922
Ann. Carnegie Mus., 13(3–4): 263.
Holotype: FMNH 59149 (CM 8287). Taiwan: Toko.
CENTRISCIDAE
- Centropristes melanus*** Ginsburg, 1952
J. Wash. Acad. Sci., 42(3): 84.
Holotype: FMNH 33719. Florida: Newport near Wakulla.
Paratypes: 6. Same locality.
SERRANIDAE
- Centropyge argi*** Woods and Kanazawa, 1951
Fieldiana Zool., 31(53): 636.
Holotype: FMNH 48494. Bermuda: Argus Bank.
POMACANTHIDAE
- Centropyge heraldi*** Woods and Schultz, 1953
Bull. U.S. Natl. Mus., 202(1): 602.
Paratype. Marshall Islands: Bikini Atoll, Boby Island.
POMACANTHIDAE
- Centropyge tutuilae*** Jordan and Jordan, 1922
Mem. Carnegie Mus., 10(1): 62.
Holotype: FMNH 55206 (CM 3902). Samoan Islands: Tetuila, Pago Pago.
POMACANTHIDAE
- Cephalopholis boninius*** Jordan and Thompson, 1914
Mem. Carnegie Mus., 6(4): 248.
Holotype: FMNH 57099 (CM 6038). Japan: Bonin Islands.
SERRANIDAE
- Cephalosilurus fowleri*** Haseman, 1911
Ann. Carnegie Mus., 7(3–4): 317.
Holotype: FMNH 54254 (CM 2880). Brazil: Bahia, Cidade de Barra.
PIMELODIDAE
- Ceratobranchia binghami*** Eigenmann, 1927
Mem. Mus. Comp. Zool. Harv. Coll., 43(4): 357.
Paratypes: 5. Peru: 2 localities.
CHARACIDAE
- Cetopsorhamdia boquillae*** Eigenmann, 1922
Mem. Carnegie Mus., 9(1): 37.
Holotype: FMNH 55212 (CM 3923). Colombia: Boquilla.
Paratypes: 5. Same locality.
PIMELODIDAE
- Cetopsorhamdia nasus*** Eigenmann and Fisher, 1916
In: Eigenmann, Ann. Carnegie Mus., 10(1–2): 83.
Holotype: FMNH 58126 (CM 7124). Colombia: Honda.
PIMELODIDAE
- Cetopsorhamdia phantasia*** Stewart, 1985
Copeia 1985(2): 340.
Holotype: FMNH 94601. Ecuador: Río Jivino, west of Limoncocha.
Paratype. Same locality.
PIMELODIDAE
- Chaenopsis alepidota californiensis*** Böhlke, 1957
Proc. Acad. Nat. Sci. Phila., 109: 99.
Paratypes: 4. California: Santa Catalina Island, Avalon.
CHAENOPSIDAE
- Chaenopsis roseola*** Hastings and Shipp, 1980
Proc. Biol. Soc. Wash., 93(4): 876.
Paratype. Florida: southwest of Panama City Beach.
CHAENOPSIDAE

Chaenothorax eigenmanni Ellis, 1913
Ann. Carnegie Mus., 8(3-4): 393.
Holotype: FMNH 54880 (CM 3542). Brazil: São
Luiz de Cáceres.
Paratype. Same locality.
CALLICHTHYIDAE

Chaetostomus leucomelas Eigenmann, 1917
Proc. Am. Philos. Soc., 56(7): 681.
Paratype. Colombia: Río Patía.
LORICARIIDAE

Characidium blennioides Eigenmann, 1909
Ann. Carnegie Mus., 6(1): 37.
Holotype: FMNH 71043 (CM 1144). Guyana:
Erukin.
Paratypes: 19. Guyana: 6 localities.
CHARACIDAE

Characidium catenatum Eigenmann, 1909
Ann. Carnegie Mus., 6(1): 40.
Holotype: FMNH 52761 (CM 1153). Guyana:
Warraputa.
Paratypes: 6. Guyana: 2 localities.
CHARACIDAE

Characidium caucanum Eigenmann, 1912
Indiana Univ. Stud., 16: 25.
Holotype: FMNH 56057 (CM 4847). Colombia:
Cali.
Paratypes: 20. Colombia: 3 localities.
CHARACIDAE

Characidium laterale Eigenmann, 1909
Ann. Carnegie Mus., 6(1): 36.
Holotype: FMNH 72627 (CM 1141). Guyana:
Amatuk.
CHARACIDAE

Characidium pellucidum Eigenmann, 1909
Ann. Carnegie Mus., 6(1): 39.
Holotype: FMNH 52764 (CM 1156). Guyana:
Gluck Island.
CHARACIDAE

Characidium phoxocephalum Eigenmann, 1912
Indiana Univ. Stud., 16: 26.
Holotype: FMNH 56061 (CM 4851). Colombia:
Paila.
CHARACIDAE

Characidium pteroides Eigenmann, 1909
Ann. Carnegie Mus., 6(1): 40.
Holotype: FMNH 53548 (CM 1157). Guyana:
Konawaruk.
Paratype. Guyana: Rockstone.
CHARACIDAE

Characidium vintoni Eigenmann, 1909
Ann. Carnegie Mus., 6(1): 36.
Holotype: FMNH 52756 (CM 1142). Guyana:
Shrimp Creek.
Paratypes: 22. Same locality.
CHARACIDAE

Characidium zebra Eigenmann, 1909
Ann. Carnegie Mus., 6(1): 38.
Holotype: FMNH 53547 (CM 1151). Guyana:
Ireng River, Maripicru Creek.
Paratypes: 7. Same locality.
CHARACIDAE

Characodon multiradiatus Meek, 1904
Field Mus. Nat. Hist. Zool. Ser., 5: 119.
Holotype: FMNH 4523. Mexico: Río Lerma at
Lerma.
Paratypes: 23. Same locality.
GOODEIDAE

Charax metae Eigenmann, 1922
Mem. Carnegie Mus., 9(1): 238.
Paratypes: 10. Colombia: Quebrada Cramalote
at Villavicencio.
CHARACIDAE

Charax rupununi Eigenmann, 1912
Mem. Carnegie Mus., 5: 402.
Holotype: FMNH 53663 (CM 2135). Guyana:
Rupununi.
Paratype. Same locality.
CHARACIDAE

Chasmias misakius Jordan and Snyder, 1901
Proc. U.S. Natl. Mus., 23(1235): 761.
Paratypes: 35. Japan: Sagami Sea, Misaki.
GOBIIDAE

Chasmocranus brevior Eigenmann, 1912
Mem. Carnegie Mus., 5: 162.
Holotype: FMNH 53213 (CM 1662). Guyana:
Warratuk.
PIMELODIDAE

Chasmocranus chimantanus Inger, 1956
Fieldiana Zool., 34(37): 425.
Holotype: FMNH 45704. Venezuela: Río Abá-
capa, west side of Chimantá-tepui.
Paratypes: 3. Same locality.
PIMELODIDAE

- Chasmocranus longior** Eigenmann, 1912
 Mem. Carnegie Mus., 5: 160.
 Holotype: FMNH 53208 (CM 1655). Guyana: Amatuk.
 Paratypes: 13. Guyana: 5 localities.
 PIMELODIDAE
- Chasmocranus rosae** Eigenmann, 1922
 Mem. Carnegie Mus., 9(1): 220.
 Holotype: FMNH 55140 (CM 3841). Colombia: Río Negro at Villavicencio.
 Paratypes: 2. Same locality.
 PIMELODIDAE
- Chasmodes bosquianus longimaxilla** Williams, 1983
 Bull. Fla. State Mus. Biol. Sci., 29(2): 85.
 Paratypes: 14. Gulf of Mexico: 4 localities.
 BLENNIIDAE
- Cheirocerus eques** Eigenmann, 1917
 Ann. Carnegie Mus., 11(3–4): 398.
 Holotype: FMNH 58255 (CM 7254). Bolivia: Villa Bella.
 Paratypes: 5. Bolivia: 3 localities.
 PIMELODIDAE
- Cheirodon annae** McAtee, 1903
 Proc. Acad. Nat. Sci. Phila., 59: 515.
 Paratypes: 2. South America.
 CHARACIDAE
- Cheirodon dialepturus** Fink and Weitzman, 1974
 Smithson. Contrib. Zool., 172: 5.
 Paratypes: 2. Panama: Río San Pedro basin, west of Santiago.
 CHARACIDAE
- Cheirodon eigenmanni** Meek, 1912
 Field Mus. Nat. Hist. Zool. Ser., 10(7): 70.
 Holotype: FMNH 7683. Costa Rica: La Junta.
 Paratype. Same locality.
 CHARACIDAE
- Cheirodon galusdae** Eigenmann, 1927
 Mem. Natl. Acad. Sci. Wash., 22(2): 44.
 Paratypes (?): 3. Chile: Río Locomilla at San Javier.
 These came to Field Museum via University of Illinois, Urbana, where they had been sent by Eigenmann for a study of fish parasites.
 CHARACIDAE
- Cheirodon interruptus ibicuihensis** Eigenmann, 1915
 Mem. Carnegie Mus., 7(1): 74.
 Syntypes: 8, FMNH 57833 (CM 6816). Brazil: Cacequy.
 CHARACIDAE
- Cheirodon madeirae** Eigenmann, 1915
 Mem. Carnegie Mus., 7(1): 76.
 Holotype: FMNH 57864 (CM 6847). Bolivia: San Joaquín.
 CHARACIDAE
- Cheirodon microdon** Eigenmann, 1915
 Mem. Carnegie Mus., 7(1): 80.
 Holotype: FMNH 57867 (CM 6850). Brazil: Cáceres.
 Paratype. Same locality.
 CHARACIDAE
- Cheirodon notomelas** Eigenmann, 1915
 Mem. Carnegie Mus., 7(1): 74.
 Holotype: FMNH 57829 (CM 6812). Brazil: Rio Tietê basin at Miguel Calmone.
 Paratypes: 10. Same locality.
 CHARACIDAE
- Cheirodon parahybae** Eigenmann, 1915
 Mem. Carnegie Mus., 7(1): 70.
 Holotype: FMNH 57857 (CM 6841). Brazil: Campos.
 Paratypes: 5. Same locality.
 CHARACIDAE
- Cheirodon stenodon** Eigenmann, 1915
 Mem. Carnegie Mus., 7(1): 82.
 Holotype: FMNH 57865 (CM 6848). Brazil: Bebedouro near Rio Grande and Rio Paraná.
 Paratypes: 28. Same locality.
 CHARACIDAE
- Chelidonichthys ischyryus** Jordan and Thompson, 1914
 Mem. Carnegie Mus., 6(4): 282.
 Holotype: FMNH 57517 (CM 6459). Japan: Sagami Bay.
 TRIGLIDAE
- Chirostoma attenuatum** Meek, 1902
 Field Mus. Nat. Hist. Zool. Ser., 3(16): 112.
 Holotype: FMNH 3631. Mexico: Lago de Patzcuaro at Patzcuaro.
 Paratypes: 12. Same locality.
 ATHERINIDAE

- Chirostoma consocium** Jordan and Hubbs, 1919
Stanford Univ. Publ., Univ. Ser., p. 76.
Holotype: FMNH 3672. Mexico: Lago Chapala at La Palma.
Paratypes: 88. Mexico: 2 localities.
ATHERINIDAE
- Chirostoma jordani** Woolman, 1894
Bull. U.S. Fish Comm. 1894, 14: 62.
Lectoparatypes: 10. Mexico: Río de Lerma at Salamanca.
Lectotype designated by Barbour (1973, Tulane Univ. Stud. Zool. Bot., 18(3): 103).
ATHERINIDAE
- Chirostoma labarcae** Meek, 1902
Field Mus. Nat. Hist. Zool. Ser., 3(6): 113.
Holotype: FMNH 3640. Mexico: Río Lerma at La Barca.
Paratypes: 8. Mexico: 2 localities.
ATHERINIDAE
- Chirostoma mezquital** Meek, 1904
Field Mus. Nat. Hist. Zool. Ser., 5: 170.
Holotype: FMNH 4389. Mexico: Río Mezquital at Durango.
Paratypes: 7. Same locality.
ATHERINIDAE
- Chirostoma patzcuaro** Meek, 1902
Field Mus. Nat. Hist. Zool. Ser., 3(6): 113.
Holotype: FMNH 3628. Mexico: Lago de Patzcuaro at Patzcuaro.
Paratypes: 8. Same locality.
ATHERINIDAE
- Chirostoma regani** Jordan and Hubbs, 1919
Stanford Univ. Publ., Univ. Ser., p. 74.
Holotype: FMNH 3687. Mexico: Xochimilco.
Paratypes: 13. Same locality.
ATHERINIDAE
- Chirostoma zirahuén** Meek, 1902
Field Mus. Nat. Hist. Zool. Ser., 3(6): 114.
Holotype: FMNH 3609. Mexico: Lago de Zirahuén at Zirahuén.
Paratypes: 29. Same locality.
ATHERINIDAE
- Chondrostoma cyri orientalis** Bianco and Banarescu, 1982
Cybium, Ser. 3, 6(2): 80.
Paratypes: 2. Iran: Pulwar River near Persepolis.
CYPRINIDAE
- Chonerhinus amabilis** Roberts, 1982
Proc. Calif. Acad. Sci., 43(1): 5.
Paratypes: 2. Indonesia: Kapuas River, southwest of Putussibau.
TETRAODONTIDAE
- Chonerhinus nefastus** Roberts, 1982
Proc. Calif. Acad. Sci., 43(1): 10.
Paratypes: 8. Malaysia: Sarawak, 3 localities.
TETRAODONTIDAE
- Chonerhinus remotus** Roberts, 1982
Proc. Calif. Acad. Sci., 43(1): 11.
Holotype: FMNH 68476. Malaysia: Sabah, mouth of Sungei Deramakot.
Paratypes: 74. Malaysia: Sabah, 4 localities.
TETRAODONTIDAE
- Chonerhinus silus** Roberts, 1982
Proc. Calif. Acad. Sci., 43(1): 12.
Holotype: FMNH 68477. Malaysia: Sarawak, Rejang basin, Sungei Baleh.
Paratypes: 56. Malaysia: Sarawak, 5 localities.
TETRAODONTIDAE
- Chromis alta** Greenfield and Woods, 1980
Copeia, 1980(4): 630.
Paratypes: 14. Mexico and Ecuador, Galápagos: 2 localities.
POMACENTRIDAE
- Chromis atripectoralis** Welander and Schultz, 1951
J. Wash. Acad. Sci., 41(3): 107.
Paratypes: 26. Philippine and Marshall Islands: 4 localities.
POMACENTRIDAE
- Chromis limbaughi** Greenfield and Woods, 1980
Copeia, 1980(4): 627.
Paratypes: 12. Mexico: Baja California, 3 localities.
POMACENTRIDAE
- Chromis meridiana** Greenfield and Woods, 1980
Copeia, 1980(4): 634.
Holotype: FMNH 75774. Chile: Isla San Felix.
Paratypes: 7. Same locality.
POMACENTRIDAE
- Chromis mossambicus** Peters, 1852
Monatsber. Akad. Wiss. Berl., 1852: 681.
Syntypes: 2, FMNH 54267 (CM 2898). Mozambique.
CICHLIDAE

- Chromis randalli** Greenfield and Hensley, 1970
Copeia, 1970(4): 689.
Paratype. Easter Island: Motu Nui.
POMACENTRIDAE
- Chromis sparsidens** Hilgendorf, 1905
Zool. Jahrb. Abt. Syst. Geogr. Biol. Tiere, 22(4): 408.
Syntypes: 2, FMNH 54272 (CM 2902). Tanzania: Manyara Lake.
CICHLIDAE
- Chromis villadolidi** Jordan and Tanaka, 1927
Ann. Carnegie Mus., 17(12): 287.
Holotype: FMNH 59185 (CM 8328). Sea of Japan: Kiusiu.
POMACENTRIDAE
- Chromis weberi** Fowler and Bean, 1928
Bull. U.S. Natl. Mus., 100(7): 41.
Paratype. Philippine Islands: Tataan Island.
POMACENTRIDAE
- Chromis woodsi** Bruner and Arnam, 1979
Fieldiana Zool., 73(3): 50.
Holotype: FMNH 85011. Western Indiana Ocean: Somalia coast.
Paratypes: 13. Western Indian Ocean: 5 localities.
POMACENTRIDAE
- Cichlasoma acutum** Miller, 1907
Bull. Am. Mus. Nat. Hist., 23(2): 117.
Paratype. Guatemala: Río Kilagua at los Amates.
CICHLIDAE
- Cichlasoma alfari** Meek, 1907
Field Mus. Nat. Hist. Zool. Ser., 7(5): 148.
Holotype: FMNH 6032. Costa Rica: Turrialba.
Paratypes: 4. Same locality.
One specimen in the paratype series, FMNH 6033, is outside the size range given in the original description.
CICHLIDAE
- Cichlasoma boliviense** Kullander, 1983
A revision of the South American cichlid genus *Cichlasoma* . . . , Stockholm, p. 165.
Paratypes: 2. Bolivia: Santa Cruz de la Sierra.
CICHLIDAE
- Cichlasoma calobrense** Meek and Hildebrand, 1913
Field Mus. Nat. Hist. Zool. Ser., 10(8): 90.
Holotype: FMNH 7600. Panama: Río Calobre.
Paratypes: 38. Panama: 7 localities.
CICHLIDAE
- Cichlasoma centrale** Meek, 1906
Field Mus. Nat. Hist. Zool. Ser., 7(3): 94.
Holotype: FMNH 5510. Guatemala: Caballo Blanco.
CICHLIDAE
- Cichlasoma dorsatum** Meek, 1907
Field Mus. Nat. Hist. Zool. Ser., 7(4): 123.
Holotype: FMNH 5971. Nicaragua: Lago Managua at Managua.
Paratypes: 4. Nicaragua: 3 localities.
CICHLIDAE
- Cichlasoma eigenmanni** Meek, 1902
Field Mus. Nat. Hist. Zool. Ser., 3(6): 119.
Holotype: FMNH 3712. Mexico: Río Tehuacán at Venta Salada.
Paratype. Same locality.
CICHLIDAE
- Cichlasoma evermanni** Meek, 1904
Field Mus. Nat. Hist. Zool. Ser., 5: 214.
Holotype: FMNH 4727. Mexico: Oaxaca, Tehuantepec.
Paratypes: 5. Same locality.
CICHLIDAE
- Cichlasoma frontale** Meek, 1909
Field Mus. Nat. Hist. Zool. Ser., 7(7): 210.
Holotype: FMNH 6358. Costa Rica: Turru-bares.
Paratype. Same locality.
CICHLIDAE
- Cichlasoma grammodes** Taylor and Miller, 1980
Occas. Pap. Mus. Zool. Univ. Mich., 693: 2.
Paratypes: 9. Mexico: Río Chico at Chiapa.
CICHLIDAE
- Cichlasoma granadense** Meek, 1907
Field Mus. Nat. Hist. Zool. Ser., 7(4): 121.
Holotype: FMNH 5951. Nicaragua: Lago Nicaragua at Granada.
Paratypes: 9. Nicaragua: 4 localities.
CICHLIDAE
- Cichlasoma hartwegi** Taylor and Miller, 1980
Occas. Pap. Mus. Zool. Univ. Mich., 693: 8.
Paratype. Mexico: Río Salado near Chiapilla.
CICHLIDAE
- Cichlasoma hedricki** Meek, 1904
Field Mus. Nat. Hist. Zool. Ser., 5: 208.
Holotype: FMNH 4673. Mexico: Oaxaca, Obispo.
Paratypes: 42. Mexico: 3 localities.
CICHLIDAE

- Cichlasoma meeki** Hildebrand, 1925
Bull. Bur. Fish., 41: 275.
Paratypes: 11. El Salvador: 4 localities.
CICHLIDAE
- Cichlasoma milleri** Meek, 1907
Field Mus. Nat. Hist. Zool. Ser., 7(5): 142.
Holotype: FMNH 5529. Guatemala: El Rancho.
Paratypes: 11. Same locality.
CICHLIDAE
- Cichlasoma mojarra** Meek, 1904
Field Mus. Nat. Hist. Zool. Ser., 5: 217.
Holotype (?): FMNH 4719. Mexico: Oaxaca, San Gerónimo.
See comment in Grey (1947, p. 133); FMNH 4719 agrees closely with published size and illustration. Published number FMNH 4718 is a lot identified by Meek as *C. melanurum*, and recently by R. Miller as *C. zonatum*.
CICHLIDAE
- Cichlasoma nigrum** Meek, 1907
Field Mus. Nat. Hist. Zool. Ser., 7(4): 128.
Holotype: FMNH 5979. Nicaragua: Lago Nicaragua, Granada.
CICHLIDAE
- Cichlasoma ornatum gephyrum** Eigenmann, 1922
Mem. Carnegie Mus., 9(1): 205.
Holotype: FMNH 58614 (CM 7639). Colombia: Río Dagua at Córdoba. Paratypes: 4. Colombia: 3 localities.
CICHLIDAE
- Cichlasoma paranaense** Kullander, 1983
A revision of the South American cichlid genus *Cichlasoma* . . . , Stockholm, p. 241.
Paratypes: 3. Brazil: Rio Tietê at Salto das Cruces.
CICHLIDAE
- Cichlasoma punctatum** Meek, 1909
Field Mus. Nat. Hist. Zool. Ser., 7(7): 210.
Holotype: FMNH 6478. Costa Rica: Buenos Aires de Térraba.
Paratypes: 10. Same locality.
CICHLIDAE
- Cichlasoma spilotum** Meek, 1912
Field Mus. Nat. Hist. Zool. Ser., 10(7): 73.
Holotype: FMNH 7686. Costa Rica: Victoria.
Paratypes (?): 10. Costa Rica: Zent.
See comment in Grey (1947, p. 134); status of paratypes uncertain.
CICHLIDAE
- Cichlasoma tenue** Meek, 1906
Field Mus. Nat. Hist. Zool. Ser., 7(3): 94.
Holotype: FMNH 3781. Mexico: Veracruz, Achotal.
Paratypes: 4. Same locality.
CICHLIDAE
- Cichlasoma tuba** Meek, 1912
Field Mus. Nat. Hist. Zool. Ser., 10(7): 73.
Holotype: FMNH 7681 MISSING. Costa Rica: Victoria.
Paratype (?). Same locality.
See comment in Grey (1947, pp. 134–135); status of the paratype uncertain.
CICHLIDAE
- Cichlasoma tuyrense** Meek and Hildebrand, 1913
Field Mus. Nat. Hist. Zool. Ser., 10(8): 89.
Holotype: FMNH 7599. Panama: Río Tuyra, Boca de Cupe.
Paratypes: 65. Panama: 6 localities.
CICHLIDAE
- Cichlasoma umbriferum** Meek and Hildebrand, 1913
Field Mus. Nat. Hist. Zool. Ser., 10(8): 88.
Holotype: FMNH 7598. Panama: Río Cupe at Cituro.
Paratypes: 20. Panama: 3 localities.
CICHLIDAE
- Cichlasoma urophthalmus mayorum** Hubbs, 1936
Carnegie Inst. Wash., Publ. 457: 277.
Paratypes: 3. Mexico: Yucatán, Chichen Itza, Xtolok Cenote.
CICHLIDAE
- Cichlasoma zonatum** Meek, 1905
Proc. Biol. Soc. Wash., 18: 245.
Holotype: FMNH 3776. Mexico: Oaxaca, Nilotpec.
Paratype. Same locality.
CICHLIDAE
- Cirricaecula johnsoni** Schultz, 1953
Bull. U.S. Natl. Mus., 202(1): 50.
Paratype. Marshall Islands: Rongerik Atoll, Bock Island.
OPHICHTHIDAE
- Clarias malaris** Nichols and Griscom, 1917
Bull. Am. Mus. Nat. Hist., 37(25): 705.
Paratype. Zaïre: Stanleyville.
CLARIIDAE

- Clarias mossambicus** Peters, 1852
Monatsber. Akad. Wiss. Berl., 1852: 682.
Syntype (?): FMNH 54274 (CM 2904). Mozambique: Quelimane.
This precise locality was not mentioned in original description.
CLARIIDAE
- Clinocottus analis australis** Hubbs, 1926
Occas. Pap. Mus. Zool. Univ. Mich., 171: 11.
Paratypes: 146. California: 4 localities.
COTTIDAE
- Cnesterodon carnegiei** Haseman, 1911
Ann. Carnegie Mus., 7(3–4): 385.
Holotype: (CM 2868) MISSING. Brazil: Rio Iguaçu at Serrinha Paraná.
Paratypes: 120. Brazil and Uruguay: 4 localities.
POECILIIDAE
- Cobitis aurata vallahica** Nalbant, 1957
Senck. Biol., 38(3/4): 209.
Paratypes: 5. Romania: Ialomitza River at Crivina.
COBITIDAE
- Cobitis taenia melanoleuca** Nichols, 1925
Am. Mus. Nov., 170: 3.
Paratype. China: Shansi.
COBITIDAE
- Coelorrhynchus gilberti** Jordan and Hubbs, 1925
Mem. Carnegie Mus., 10(2): 327.
Holotype: FMNH 59401 (CM 7960). Japan: market at Shizuoka.
MACROURIDAE
- Compsura heterura** Eigenmann, 1915
Mem. Carnegie Mus., 7(1): 61.
Holotype: FMNH 57825 (CM 6808). Brazil: Rio Itapicurú at Queimadas.
Paratypes: 11. Same locality.
CHARACIDAE
- Conger jordani** Kanazawa, 1958
Proc. U.S. Natl. Mus., 108(3400): 250.
Paratype. Japan: Tokyo Market.
CONGRIDAE
- Copeina osgoodi** Eigenmann, 1922
Mem. Carnegie Mus., 9(1): 232.
Holotype: FMNH 15352. Peru: mouth of Río Yavary at Nazareth.
Paratypes: 6. Same locality.
LEBIASINIDAE
- Copidoglanis novaeguineae** Weber, 1908
Nova Guinea, 5(2): 226.
Syntype: FMNH 52388 (CM 644). New Guinea: Lake Sentani.
PLOTOSIDAE
- Coregonus stanleyi** Kendall, 1903
Bull. U.S. Fish Comm. 1902, 22: 366.
Paratypes: 2. Maine: Aroostook County, Eagle Lake.
SALMONIDAE
- Coripareius septentrionalis** Nichols, 1925
Am. Mus. Nov., 181: 2.
Paratype: MISSING. China: Mongolia, Yellow River at Paotow.
CYPRINIDAE
- Corydoras australe** Eigenmann and Ward, 1907
Ann. Carnegie Mus., 4(2): 123.
Paratype. Paraguay: Corumbá.
CALLICHTHYIDAE
- Corydoras blochi** Nijssen, 1971
Beaufortia, 19(250): 92.
Holotype: FMNH 75951. Guyana: Moco Creek near Lehtem.
Paratypes: 2. Brazil: Rio Amazonas at Santarém.
CALLICHTHYIDAE
- Corydoras bondi coppenamensis** Nijssen, 1970
Beaufortia, 18(230): 19.
Paratypes: 2. Surinam: Tributary of Coppename River.
CALLICHTHYIDAE
- Corydoras ellisae** Gosline, 1940
Stanford Ichthyol. Bull., 2(1): 17.
Holotype: FMNH 57361 (CM 3522). Paraguay: Arroyo Poná at Sapucay.
Paratype. Paraguay: near Sapucay.
CALLICHTHYIDAE
- Corydoras heteromorphus** Nijssen, 1970
Beaufortia, 18(230): 22.
Paratypes: 2. Surinam: tributary of Coppename River.
CALLICHTHYIDAE
- Corydoras schwartzi surinamensis** Nijssen, 1970
Beaufortia, 18(230): 39.
Paratype. Surinam: tributary of Coppename River.
CALLICHTHYIDAE

- Corydoras weitzmani** Nijssen, 1971
Beaufortia, 19(250): 91.
Holotype: FMNH 69934. Peru: Río Vilcanota system at Cuzco.
Paratypes: 2. Same locality.
CALLICHTHYIDAE
- Corymbophanes andersoni** Eigenmann, 1909
Ann. Carnegie Mus., 6(1): 5.
Holotype: FMNH 52675 (CM 1001). Guyana: Potaro River at Aruataima Falls.
LORICARIIDAE
- Coryphaenoides bonanox** Jordan and Thompson, 1914
Mem. Carnegie Mus., 6: 305.
Holotype: FMNH 57520 (CM 6462). Japan: Sagami Bay.
MACROURIDAE
- Coryphopterus punctipectophorus** Springer, 1960
Bull. Mar. Sci. Gulf Caribb., 10(2): 237.
Paratypes: 3. Gulf of Mexico: off Florida.
GOBIIDAE
- Corythoichthys elerae** Evermann and Seale, 1907
Bull. Bur. Fish. 1906, 26: 57.
Paratype. Philippine Islands: Sorsogon, Bacon. Misspelled *Corythoichthys* in Evermann and Seale (1907).
SYNGNATHIDAE
- Corythoichthys polynotatus** Dawson, 1977
Copeia, 1977(2): 331.
Paratypes: 4. Philippine Islands: Negros Island.
SYNGNATHIDAE
- Cotylis nigripinnis woodsi** Schultz, 1944
Proc. U.S. Natl. Mus., 96(3187): 68.
Holotype: FMNH 41794. Cocos Island: Wafer Bay.
GOBIESOCIDAE
- Creagrutus beni** Eigenmann, 1911
Ann. Carnegie Mus., 8(1): 172.
Holotype: FMNH 54585 (CM 3216). Bolivia: Río Beni at Villa Bella.
CHARACIDAE
- Creagrutus brevipinnis** Eigenmann, 1913
Indiana Univ. Stud., 18: 10.
Holotype: FMNH 56095 (CM 4887). Colombia: Piedra de Moler.
Paratypes: 59. Colombia: 3 localities.
CHARACIDAE
- Creagrutus caucanus** Eigenmann, 1913
Indiana Univ. Stud., 18: 9.
Holotype: FMNH 56104 (CM 4895). Colombia: Paila.
Paratypes: 19. Colombia: 4 localities.
CHARACIDAE
- Creagrutus magdalenae** Eigenmann, 1913
Indiana Univ. Stud., 18: 8.
Holotype: FMNH 56088 (CM 4880). Colombia: Girardot.
Paratypes: 108. Colombia: 3 localities.
CHARACIDAE
- Creagrutus melanzonus** Eigenmann, 1909
Ann. Carnegie Mus., 6(1): 30.
Holotype: FMNH 52705 (CM 1067). Guyana: Crab Falls.
Paratypes: 2. Guyana: 2 localities.
CHARACIDAE
- Creagrutus notropoides** Meek and Hildebrand, 1912
Field Mus. Nat. Hist. Zool. Ser., 10(6): 68.
Holotype: FMNH 7574. Panama: Río Indio.
Paratypes: 67. Panama: 5 localities.
CHARACIDAE
- Creagrutus simus** Meek and Hildebrand, 1913
Field Mus. Nat. Hist. Zool. Ser., 10(8): 85.
Holotype: FMNH 7591. Panama: Río Cupe, Cituro.
Paratypes: 232. Panama: 5 localities.
CHARACIDAE
- Crenicara altispinosa** Haseman, 1911
Ann. Carnegie Mus., 7(3-4): 344.
Holotype: FMNH 54090 (CM 2639). Bolivia: Río Mamoré, below mouth of Río Guaporé.
Paratypes: 7. Bolivia: 2 localities.
CICHLIDAE
- Crenicichla alta** Eigenmann, 1912
Mem. Carnegie Mus., 5: 516.
Holotype: FMNH 53777 (CM 2274). Guyana: Gluck Island.
Paratypes: 35. Guyana: 13 localities.
CICHLIDAE
- Crenicichla dorsocellata** Haseman, 1911
Ann. Carnegie Mus., 7(3-4): 355.
Holotype: FMNH 54155 (CM 2721). Brazil: Rio Parahyba at Campos.
Paratype. Brazil: Santarem.
CICHLIDAE

Crenicichla iguassuensis Haseman, 1911
Ann. Carnegie Mus., 7(3–4): 352.
Holotype: FMNH 54159 (CM 2725). Brazil: Rio Iguaçu at Porto União da Victoria.
Paratypes: 4. Brazil: 2 localities.
CICHLIDAE

Crenicichla jaguarensis Haseman, 1911
Ann. Carnegie Mus., 7(3–4): 351.
Holotype: FMNH 54157 (CM 2723). Brazil: Rio Grande of the Paraná at Jaguará.
Paratype. Same locality.
CICHLIDAE

Crenicichla santaremensis Haseman, 1911
Ann. Carnegie Mus., 7(3–4): 354.
Holotype: FMNH 54153 (CM 2719). Brazil: near Santarém.
Paratype. Same locality.
CICHLIDAE

Crenicichla simoni Haseman, 1911
Ann. Carnegie Mus., 7(3–4): 345.
Holotype: FMNH 54097 (CM 2646). Brazil: São Luiz de Cáceres.
Paratypes: 7. Brazil, Bolivia, and Paraguay: 4 localities.
CICHLIDAE

Crossochir koelzi Hubbs, 1933
Proc. U.S. Natl. Mus., 82(2962): 4.
Paratypes: 6. California: San Diego.
EMBIOTOCIDAE

Crossostoma fascicauda Nichols, 1926
Am. Mus. Nov., 224: 2.
Paratype. China: Fukien, Fuching Hsien.
COBITIDAE

Cryptocentrus geniornatus Herre, 1935
Field Mus. Nat. Hist. Zool. Ser., 18(12): 428.
Holotype: FMNH 17364. Indonesia: Waigiu Island.
GOBIIDAE

Cryptotomus crassiceps Bean, 1906
Proc. Biol. Soc. Wash., 19: 32.
Lectotype: FMNH 4964. Bermuda: Cooper's Island.
Lectotype designated by Grey (1947, p. 138).
SCARIDAE

Ctenogobius aterrimus Herre, 1935
Field Mus. Nat. Hist. Zool. Ser., 18(12): 423.
Holotype: FMNH 17384. Solomon Islands: Kulambangra Island.
GOBIIDAE

Ctenogobius leavelli Herre, 1935
Lingnan Sci. J. Canton, 14(3): 396.
Paratypes: 6. China: Kwangsi, Wuchow.
GOBIIDAE

Ctenogobius malekulae Herre, 1935
Field Mus. Nat. Hist. Zool. Ser., 18(12): 423.
Holotype: FMNH 17385. New Hebrides Islands: Malekula Island, Bushman's Bay.
GOBIIDAE

Ctenogobius myxodermus Herre, 1935
Lingnan Sci. J. Canton, 14(3): 395.
Paratypes: 10. China: 2 localities.
GOBIIDAE

Ctenogobius opalescens Herre, 1936
Bull. Raffles Mus., 12: 15.
Paratypes: 3. Singapore.
GOBIIDAE

Ctenogobius villadolidi Herre, 1936
Philipp. J. Sci., 59(3): 361.
Paratypes: 2. Philippine Islands: Negros Island, Dumaguete.
GOBIIDAE

Ctenogobius virgatulus Jordan and Snyder, 1901
Proc. U.S. Natl. Mus., 24(1244): 63.
Paratype: MISSING. Japan: Sagami, Misaki.
GOBIIDAE

Ctenogobius waigiensis Herre, 1935
Field Mus. Nat. Hist. Zool. Ser., 18(12): 424.
Holotype: FMNH 17413. Indonesia: Waigiu Island.
Paratypes: 77. Same locality.
GOBIIDAE

Ctenopoma multispinis Peters, 1844
Monatsber. Akad. Wiss. Berl., 1844: 34 (not seen).
Syntype: FMNH 54262 (CM 2891). Mozambique: Quelimane.
ANABANTIDAE

Culter aokii Oshima, 1919
Ann. Carnegie Mus., 12(2–4): 250.
Holotype: FMNH 59110 (CM 8248). Taiwan: Jitsugetsutan (Lake Candidius).
CYPRINIDAE

- Curimata cerasina** Vari, 1983
Acta Biol. Venez., 11(4): 30.
Paratype. Venezuela: Guárico, pit along side of road from Calabozo to Camaguán.
CURIMATIDAE
- Curimata incompta** Vari, 1983
Acta Biol. Venez., 11(4): 34.
Paratypes: 2. Venezuela: 2 localities.
CURIMATIDAE
- Curimatus atratoensis** Eigenmann, 1912
Indiana Univ. Stud., 16: 19.
Holotype: FMNH 56024 (CM 4814). Colombia: Quibdó.
Paratypes: 48. Same locality.
CURIMATIDAE
- Curimatus gillii** Eigenmann and Kennedy, 1903
Proc. Acad. Nat. Sci. Phila., 59: 510.
Paratype. Paraguay: Chagalalina.
CURIMATIDAE
- Curimatus issororoensis** Eigenmann, 1912
Mem. Carnegie Mus., 5: 266.
Holotype: FMNH 53647 (CM 2119). Guyana: Issorora Rubber Plantation.
Paratype. Same locality.
CURIMATIDAE
- Curimatus metae** Eigenmann, 1922
Mem. Carnegie Mus., 9(1): 230.
Holotype: FMNH 55143 (CM 3845). Colombia: Quebrada Cramalote at Villavicencio.
Paratypes: 2. Same locality.
CURIMATIDAE
- Curimatus morawhannae** Eigenmann, 1912
Mem. Carnegie Mus., 5: 266.
Holotype: FMNH 53650 (CM 2122). Guyana: Morawhanna.
Paratypes: 14. Guyana: 2 localities.
CURIMATIDAE
- Curimatus patiae** Eigenmann, 1914
Indiana Univ. Stud., 19: 12.
Holotype: FMNH 56554 (CM 5368). Colombia: Barbacoas.
Paratypes: 3. Same locality.
CURIMATIDAE
- Curimatus peruanus** Eigenmann, 1922
Mem. Carnegie Mus., 9(1): 104.
Paratype. Peru: Río Chira at Sullana.
CURIMATIDAE
- Cyclopium chapmani** Eigenmann, 1912
Indiana Univ. Stud., 16: 13.
Holotype: FMNH 56071 (CM 4863). Colombia: Boquía.
Paratypes: 2. Same locality.
ASTROBLEPIDAE
- Cyclopium pirrense** Meek and Hildebrand, 1913
Field Mus. Nat. Hist. Zool. Ser., 10(8): 83.
Holotype: FMNH 7586. Panama: Río Cana at Cana.
Paratypes: 33. Same locality.
ASTROBLEPIDAE
- Cyclopium trifasciatum** Eigenmann, 1912
Indiana Univ. Stud., 16: 14.
Holotype: FMNH 56076 (CM 4868). Colombia: Caldas.
Paratypes: 40. Colombia: 2 localities.
ASTROBLEPIDAE
- Cyclopium unifasciatum** Eigenmann, 1912
Indiana Univ. Stud., 16: 15.
Holotype: FMNH 56079 (CM 4871). Colombia: Caldas.
Paratypes: 54. Same locality.
ASTROBLEPIDAE
- Cyclopium vanceae** Eigenmann, 1913
Ann. Carnegie Mus., 8(3-4): 421.
Holotype: FMNH 56066 (CM 4856). Peru: Queta, brook southeast of Tarma.
ASTROBLEPIDAE
- Cyclopium ventrale** Eigenmann, 1912
Indiana Univ. Stud., 16: 15.
Holotype: FMNH 56074 (CM 4866). Colombia: Caldas.
Paratypes: 15. Colombia: 2 localities.
ASTROBLEPIDAE
- Cynodonichthys tenuis** Meek, 1904
Field Mus. Nat. Hist. Zool. Ser., 5: 101.
Holotype: FMNH 4643. Mexico: Oaxaca, El Hule.
APLOCHEILIDAE
- Cynoglossus heterolepis** Weber, 1910
Notes Leiden Mus., 32: 237.
Syntype: FMNH 52501 (CM 759). New Guinea: Lorentz River.
CYNOGLOSSIDAE

- Cynopotamus essequibensis** Eigenmann, 1912
 Mem. Carnegie Mus., 5: 403.
 Holotype: FMNH 53673 (CM 2146). Guyana: Potaro Landing.
 Paratypes: 4. Guyana: 2 localities.
 CHARACIDAE
- Cyprinodon bifasciatus** Miller, 1968
 Occas. Pap. Mus. Zool. Univ. Mich., 659: 1.
 Paratypes: 20. Mexico: Río Puente Colorado near Cuatro Ciénegas.
 CYPRINODONTIDAE
- Cyprinodon dearborni** Meek, 1909
 Field Mus. Nat. Hist. Zool. Ser., 7(7): 208.
 Holotype: FMNH 6484. Lesser Antilles: Curaçao, Willemstad.
 Paratypes: 13. Same locality.
 CYPRINODONTIDAE
- Cyprinodon fontinalis** Smith and Miller, 1980
 Proc. Biol. Soc. Wash., 93(2): 405.
 Paratypes: 15. Mexico: Chihuahua, Ojo de Caribonera.
 CYPRINODONTIDAE
- Cyprinodon meeki** Miller, 1976
 Bull. South. Calif. Acad. Sci., 75(2): 73.
 Paratypes: 77. Mexico: 2 localities.
 CYPRINODONTIDAE
- Cyprinodon orthonotus** Peters, 1844
 Monatsber. Akad. Wiss. Berl., 1844: 35 (not seen).
 Syntype: FMNH 54289 (CM 2920). Mozambique: Quellimane.
 CYPRINODONTIDAE
- Cypselurus alienus** Herre, 1935
 Field Mus. Nat. Hist. Zool. Ser., 18(12): 392.
 Holotype: FMNH 17224. Cuba: south coast, near Santiago.
 EXOCOETIDAE
- Cypselurus antoncichi** Woods and Schultz, 1953
 Bull. U.S. Natl. Mus., 202(1): 181.
 Paratypes: 2. Marshall Islands: 2 localities.
 EXOCOETIDAE
- Dangila sabana** Inger and Chin, 1962
 Fieldiana Zool., 45: 94.
 Holotype: FMNH 44778. Malaysia: Sabah, Danau, Bukit Garam.
 Paratypes: 65. Malaysia: Sabah, 3 localities.
 CYPRINIDAE
- Dasyatis ushieii** Jordan and Hubbs, 1925
 Mem. Carnegie Mus., 10(2): 114.
 Holotype: FMNH 59380 (CM 7778). Japan: Mikawa Bay.
 DASYATIDAE
- Dasyscopelus orientalis** Gilbert, 1913
 Mem. Carnegie Mus., 6(2): 70.
 Holotype: FMNH 55834 (CM 4613). Japan: Sagami Bay, Misaki.
 Paratypes: 52. Same locality.
 MYCTOPHIDAE
- Decapterus dayi** Wakiya, 1924
 Ann. Carnegie Mus., 15(2-3): 158.
 Holotype: FMNH 59468 (CM 7711). Taiwan.
 CARANGIDAE
- Decapterus tabl** Berry, 1968
 Contrib. Mar. Sci., 13: 152.
 Paratype. Western Atlantic: off Colombia.
 CARANGIDAE
- Dermatocheir catablepta** Durbin, 1909
 Ann. Carnegie Mus., 6(1): 55.
 Holotype: FMNH 53553 (CM 1198). Guyana: Tumatumari, above the falls.
 CHARACIDAE
- Deuterodon acanthogaster** Eigenmann, 1911
 Ann. Carnegie Mus., 8(1): 179.
 Holotype: FMNH 54748 (CM 3395). Brazil: Corumbá.
 Paratypes: 14. Brazil: 2 localities.
 CHARACIDAE
- Deuterodon atracaudata** Meek and Hildebrand, 1912
 Field Mus. Nat. Hist. Zool. Ser., 10(6): 68.
 Holotype: FMNH 7573. Panama: Canal Zone, Río Frijoles.
 Paratypes: 497. Panama: 22 localities.
 CHARACIDAE
- Deuterodon pinnatus** Eigenmann, 1909
 Ann. Carnegie Mus., 6(1): 25.
 Holotype: FMNH 53525 (CM 1046). Guyana: Potaro River at Amatuk.
 Paratypes: 46. Guyana: 5 localities.
 CHARACIDAE
- Deuterodon potaroensis** Eigenmann, 1909
 Ann. Carnegie Mus., 6(1): 27.
 Holotype: FMNH 52967 (CM 1053). Guyana: Potaro River at Amatuk.
 Paratypes: 2. Guyana: 2 localities.
 CHARACIDAE

- Diaphus gigas** Gilbert, 1913
 Mem. Carnegie Mus., 6(2): 93.
 Holotype: FMNH 55825 (CM 4601). Japan: Sagami Bay, Misaki.
 Paratype: Same locality.
 MYCTOPHIDAE
- Diaphus glandulifer** Gilbert, 1913
 Mem. Carnegie Mus., 6(2): 90.
 Paratypes: 3. Japan: Sagami Bay.
 MYCTOPHIDAE
- Diaphus latus** Gilbert, 1913
 Mem. Carnegie Mus., 6(2): 95.
 Holotype: FMNH 56208 (CM 4604). Japan: Sagami Bay, Misaki.
 MYCTOPHIDAE
- Diaphus sagamiensis** Gilbert, 1913
 Mem. Carnegie Mus., 6(2): 96.
 Holotype: FMNH 55831 (CM 4608). Japan: Sagami Bay, Misaki.
 Paratypes: 3. Same locality.
 MYCTOPHIDAE
- Diapterus evermanni** Meek and Hildebrand, 1925
 Field Mus. Nat. Hist. Zool. Ser., 15(2): 594.
 Paratypes: 2. Panama: Fox Bay, Colón.
 GERREIDAE
- Diceratias pileatus** Uwate, 1979
 Copeia, 1979(1): 140.
 Paratypes: 2. Western Atlantic Ocean: off Surinam.
 DICERATIIDAE
- Dicrolene kanazawai** Grey, 1958
 Fieldiana Zool., 39(16): 170.
 Paratypes: 6. Gulf of Mexico.
 OPHIDIIDAE
- Dionda diaboli** Hubbs and Brown, 1956
 Southwest Nat., 1(2): 69.
 Paratypes: 2. Texas: Devil's Lake.
 CYPRINIDAE
- Dionda erimyzonops** Hubbs and Miller, 1974
 Occas. Pap. Mus. Zool. Univ. Mich., 671: 4.
 Paratypes: 3. Mexico: Tamaulipas, Forlón.
 CYPRINIDAE
- Diphyacantha chocoensis** Henn, 1916
 Ann. Carnegie Mus., 10(1-2): 114.
 Paratypes: 19. Colombia: creek near mouth of Río Calima.
 POECILIIDAE
- Diplectrum labarum** Rosenblatt and Johnson, 1974
 Calif. Fish Game, 60(4): 181.
 Paratypes: 57. Mexico: Baja California, Bahía Santa Inez.
 SERRANIDAE
- Diplectrum pacificum** Meek and Hildebrand, 1925
 Field Mus. Nat. Hist. Zool. Ser., 15(2): 477.
 Paratypes: 3. Panama: Naos Island.
 SERRANIDAE
- Diplodus bermudensis** Caldwell, 1965
 Fieldiana Zool., 44(23): 217.
 Holotype: FMNH 72520. Bermuda: Saint George's Island, Whalebone Bay.
 Paratypes: 179. Bermuda: 11 localities.
 SPARIDAE
- Diplogonurus maderensis** Noronha, 1926
 Ann. Carnegie Mus., 16(3-4): 381.
 Holotype: MISSING. Cama de Lobos.
 The type, apparently the head (dried?) illustrated in Noronha, was never received by the Carnegie Museum (Henn, 1928, p. 79).
 GEMPYLIDAE
- Diretmus pauciradiatus** Woods, 1973
 Mem. Sears Fnd. Mar. Res., 1(6): 296.
 Holotype: FMNH 66797. Southern Caribbean Sea.
 Paratypes: 15. Caribbean Sea and Gulf of Mexico: 12 localities.
 DIRETMIDAE
- Disparichthys fluviatilis** Herre, 1935
 Field Mus. Nat. Hist. Zool. Ser., 18(12): 384.
 Holotype: FMNH 17192. New Guinea: Sepik River basin, brook at Marienberg.
 CARAPIDAE
- Distichodus stigmaturus** Fowler, 1935
 Ann. Transvaal Mus., 16(2): 260.
 Paratypes: 3. Botswana: Chobe River near Kasana.
 CITHARINIDAE
- Dollfusichthys sinuarabici** Chabanaud, 1931
 Bull. Soc. Zool. Fr., 56: 304.
 Paratype: Egypt: Bay of Suez.
 CYNOGLOSSIDAE
- Doraops zuloagai** Schultz, 1944
 Proc. U.S. Natl. Mus., 94(3172): 271.
 Paratype: Venezuela: Río Apón near mouth of Río Cogollo.
 DORADIDAE

- Doras lentiginosus** Eigenmann, 1917
Ann. Carnegie Mus., 11(3-4): 401.
Holotype: FMNH 58056 (CM 7048). Brazil: Santarém.
Paratype. Brazil: Manaus.
DORADIDAE
- Dormitator gymnocephalus** Eigenmann, 1912
Mem. Carnegie Mus., 5: 523.
Holotype: FMNH 53922 (CM 2438). Guyana: Konawaruk.
ELEOTRIDAE
- Dorosoma anale** Meek, 1904
Field Mus. Nat. Hist. Zool. Ser., 5: 93.
Holotype: FMNH 4637. Mexico: Oaxaca, El Hule.
Paratypes: 15. Mexico: 4 localities.
CLUPEIDAE
- Dorosoma chavesi** Meek, 1907
Field Mus. Nat. Hist. Zool. Ser., 7(4): 112.
Holotype: FMNH 5928. Nicaragua: Laguna Janicero, Granada.
Paratypes: 73. Nicaragua: 4 localities.
Grey (1947, p. 140) noted that the type, FMNH 5928, was MISSING. We now have 1 specimen labeled type and a separated lot of 6 labeled paratypes with both lots bearing FMNH 5928, but no indication of who separated the type.
CLUPEIDAE
- Doryrhamphus melanopleura cooki** Weed and Woods, 1953
Bull. U.S. Natl. Mus., 202(1): 249.
Holotype: FMNH 16220. Cook Archipelago: Aitutaki Island.
Paratype. Same locality.
SYNGNATHIDAE
- Draconetta oregona** Briggs and Berry, 1959
Copeia, 1959(2): 129.
Paratypes: 18. Northeast coast of South America: 4 localities.
DRACONETTIDAE
- Dysomma abyssale** Kamohara, 1938
On the offshore bottom-fishes of Prov. Tosa, Shikoku, Japan, p. 12 (not seen).
Neotype: FMNH 76869. Sagami Sea.
Neotype designated by Robins and Robins (1976, Proc. Acad. Nat. Sci. Phila., 127(18): 256).
SYNAPHOBRANCHIDAE
- Echidna unicolor** Schultz, 1953
Bull. U.S. Natl. Mus., 202(1): 106.
Paratype. Marshall Islands: Bikini Atoll.
MURAENIDAE
- Echidna vineta** Jenkins, 1903
Bull. U.S. Fish Comm. 1902, 22: 429.
Paratype: MISSING. Hawaiian Islands.
Published as FMNH 2753 (field no. 263), but the specimen apparently was never received.
MURAENIDAE
- Echidna zonophaea** Jordan and Evermann, 1903
Bull. U.S. Fish Comm. 1902, 22: 167.
Paratype. Hawaiian Islands: Honolulu.
MURAENIDAE
- Echiodon dawsoni** Williams and Shipp, 1982
Copeia, 1982(4): 486.
Paratypes: 2. Gulf of Mexico: off Florida.
CARAPIDAE
- Ectenias brunneus** Jordan and Thompson, 1914
Mem. Carnegie Mus., 6(4): 241.
Holotype: FMNH 57511 (CM 6453). Japan: Misaki.
Paratype. Same locality.
CORYPHAENIDAE
- Ekemblemaria myersi** Stephens, 1963
Univ. Calif. Publ. Zool., 68: 23.
Paratypes: 5. Mexico: Golfo de California, Bahía Santa Inez.
CHAENOPSIDAE
- Eleotris herwerdenii** Weber, 1910
Notes Leiden Mus., 32: 338.
Syntype: FMNH 52462 (CM 720). New Guinea: Lorentz River.
ELEOTRIDAE
- Eleotris heterodon** Weber, 1908
Nova Guinea, 5(2): 255.
Syntype: FMNH 52460 (CM 718). New Guinea: Lake Sentani.
ELEOTRIDAE
- Eleotris isthmensis** Meek and Hildebrand, 1916
Field Mus. Nat. Hist. Zool. Ser., 10(15): 359.
Holotype: FMNH 8951. Panama: Canal Zone, Mindi Cut.
Paratypes: 29. Panama: 5 localities.
ELEOTRIDAE

Eleotris latifasciatus Meek and Hildebrand, 1912
Field Mus. Nat. Hist. Zool. Ser., 10(6): 68.
Holotype: FMNH 7575. Panama: Río Cárdenas
at Corozal.
Paratypes: 3. Panama: 2 localities.
ELEOTRIDAE

Eleotris macrolepis Meek, 1914
Field Mus. Nat. Hist. Zool. Ser., 10(10): 130.
Holotype: FMNH 7775. Costa Rica: Jesús Ma-
ría.
ELEOTRIDAE

Emblemaria angelica Böhlke and Mead, 1957
Not. Nat. (Phila.), 301: 2.
Paratypes: 24 MISSING. Mexico: Acapulco Bay.
CHAENOPSIDAE

Emblemaria hyltoni Johnson and Greenfield, 1976
Fieldiana Zool., 70(2): 17.
Holotype: FMNH 80412. Honduras: Isla Roa-
tán.
Paratypes: 2. Same locality.
CHAENOPSIDAE

Emblemaria nigra Meek and Hildebrand, 1928
Field Mus. Nat. Hist. Zool. Ser., 15(3): 953.
Paratype. Panama: Porto Bello.
CHAENOPSIDAE

Emblemariopsis pricei Greenfield, 1975
Copeia, 1975(4): 713.
Holotype: FMNH 77481. Belize: Glover's Reef.
Paratype. Same locality.
CHAENOPSIDAE

Encaeura evides Jordan and Hubbs, 1925
Mem. Carnegie Mus., 10(2): 303.
Holotype: FMNH 58821 (CM 7931). Japan:
Wakanoura.
ELEOTRIDAE

Enchelycore carychroa Böhlke and Böhlke, 1976
Proc. Acad. Nat. Sci. Phila., 127(13): 138.
Paratypes: 2. Haiti: Gulf of Gonâve to Port-au-
Prince Bay.
MURAENIDAE

Engyprosopon xenandrus Gilbert, 1905
Bull. U.S. Fish Comm. 1903, 23(2): 687.
Paratypes: 2. Hawaiian Islands.
BOTHIDAE

Enneapterygius punctulatus Herre, 1935
Field Mus. Nat. Hist. Zool. Ser., 18(12): 432.
Holotype: FMNH 17388. New Hebrides Is-
lands: Wala Island.
Paratypes: 2. New Hebrides Islands: 2 localities.
TRIPTERYGIIDAE

Enneapterygius waigiensis Herre, 1935
Field Mus. Nat. Hist. Zool. Ser., 18(12): 434.
Holotype: FMNH 17494. Indonesia: Waigu Is-
land.
TRIPTERYGIIDAE

Entomacrodus randalli Springer, 1967
Proc. U.S. Natl. Mus., 122(3582): 71.
Paratypes: 2. Marquesas Islands: Nuku Hiva Is-
land.
BLENNIIDAE

Entomocorus benjamini Eigenmann, 1917
Ann. Carnegie Mus., 11(3-4): 403.
Holotype: FMNH 58019 (CM 7006). Bolivia:
San Joaquín.
Paratypes: 9. Bolivia: 2 localities.
AUCHENIPTERIDAE

Entonanthias pascalus Jordan and Tanaka, 1927
Ann. Carnegie Mus., 17(12): 385.
Holotype: FMNH 59184 (CM 8327). Japan:
Okinawa, Naha.
SERRANIDAE

Epapterus blohmi Vari, Jewett, Taphorn, and Gil-
bert, 1984
Proc. Biol. Soc. Wash., 97(2): 463.
Paratypes: 2. Venezuela: Guarico, Laguna Los
Guácimos.
AUCHENIPTERIDAE

Epinephelus elongatus Schultz, 1953
Bull. U.S. Natl. Mus., 202(1): 345.
Paratypes: 3. Marshall Islands: Bikini Atoll, 2
localities.
SERRANIDAE

Epinephelus emoryi Schultz, 1953
Bull. U.S. Natl. Mus., 202(1): 340.
Paratype: Marianas Islands: Rota Island.
SERRANIDAE

Epinephelus ionthas Jordan and Metz, 1913
Mem. Carnegie Mus., 6(1): 32.
Holotype: FMNH 55797 (CM 4559). Korea:
Fusan.
SERRANIDAE

- Epinephelus melanostigma** Schultz, 1953
Bull. U.S. Natl. Mus., 202(1): 348.
Paratypes: 2. Samoan Islands: Swains Island.
SERRANIDAE
- Epinephelus spilotoceps** Schultz, 1953
Bull. U.S. Natl. Mus., 202(1): 357.
Paratypes: 2. Marshall Islands: Bikini Atoll, 2 localities.
SERRANIDAE
- Epinnula orientalis americana** Grey, 1953
Copeia, 1953(3): 137.
Holotype: FMNH 46391. Gulf of Mexico.
Paratypes: 8. Gulf of Mexico: 2 localities.
GEMPYLIDAE
- Epinnula orientalis pacifica** Grey, 1953
Copeia, 1953(3): 138.
Holotype: FMNH 59528. Japan: Kumano-Nada, off Owase.
Paratype. Same locality.
GEMPYLIDAE
- Ernstichthys intonsus** Stewart, 1985
Fieldiana Zool., New Ser., 25: 16.
Holotype: FMNH 94603. Ecuador: Río Napo at Añangu.
ASPREDINIDAE
- Etelis evurus** Jordan and Evermann, 1903
Bull. U.S. Fish Comm. 1902, 22: 184.
Paratype. Hawaiian Islands: Hilo.
LUTJANIDAE
- Etheostoma obeyense** Kirsch, 1892
Bull. U.S. Fish Comm. 1890, 10: 292.
Paratypes: 36. Kentucky.
PERCIDAE
- Eucinostomus californiensis elongatus** Meek and Hildebrand, 1925
Field Mus. Nat. Hist. Zool. Ser., 15(2): 586.
Paratypes: 52. Panama: Taboga Island.
GERREIDAE
- Euctenogobius liolepis** Meek and Hildebrand, 1928
Field Mus. Nat. Hist. Zool. Ser., 15(3): 875.
Paratypes: 7. Panama: Canal Zone, on beach near Balboa.
GOBIIDAE
- Euctenogobius panamensis** Meek and Hildebrand, 1928
Field Mus. Nat. Hist. Zool. Ser., 15(3): 874.
Paratypes: 3. Panama: Río Culebra at Culebra.
GOBIIDAE
- Euleptoeleotris clarki** Hildebrand, 1938
Field Mus. Nat. Hist. Zool. Ser., 22(4): 352.
Paratypes: 2. Panama: Canal Zone, Miraflores locks.
ELEOTRIDAE
- Euleptoeleotris shropshirei** Hildebrand, 1938
Field Mus. Nat. Hist. Zool. Ser., 22(4): 353.
Paratype. Panama: Puerto Pilón.
ELEOTRIDAE
- Eumegistus illustris** Jordan and Jordan, 1922
Mem. Carnegie Mus., 10(1): 36.
Holotype: FMNH 59325 (CM 3899). Hawaiian Islands: Honolulu Market.
BRAMIDAE
- Eupomacentrus chrysus** Bean, 1906
Proc. Biol. Soc. Wash., 19: 32.
Holotype: FMNH 5025. Bermuda: White's Flat Channel.
POMACENTRIDAE
- Eupomacentrus rocasensis** Emery, 1972
Copeia, 1972(2): 330.
Paratypes: 12. Atlantic Ocean: off Brazil, Atol das Rocas.
POMACENTRIDAE
- Evarra tlahuacensis** Meek, 1902
Field Mus. Nat. Hist. Zool. Ser., 3(6): 85.
Holotype: FMNH 3681. Mexico: Lago Chalco, Tlahuac.
CYPRINIDAE
- Evermannella ahlstromi** Johnson and Glodek, 1975
Copeia, 1975(4): 716.
Paratypes: 19. Pacific Ocean: 7 localities.
EVERMANNELLIDAE
- Evermannella sicaria** Rofen, 1963
Aquatica, Aquatic Res. Inst. Stockton, 1: 1.
Paratypes: 2. Western North Atlantic: off Bermuda.
EVERMANNELLIDAE

Eviota japonica Jewett and Lachner, 1983
Proc. Biol. Soc. Wash., 96(4): 788.
Paratypes: 19. Japan: Ryukyu Islands, Aikawa
Rikuzen.
GOBIIDAE

Evorthodus minutus Meek and Hildebrand, 1928
Field Mus. Nat. Hist. Zool. Ser., 15(3): 870.
Paratype. Panama: Canal Zone, Corozal.
GOBIIDAE

Exerpes asper earli Hubbs, 1952
Stanford Ichthyol. Bull., 4(2): 94.
Paratypes: 5. Mexico: Baja California, Laguna
San Ignacio.
LABRISOMIDAE

Farlowella azygia Eigenmann and Vance, 1917
Ann. Carnegie Mus., 11(1-2): 299.
Holotype: FMNH 55091 (CM 3782). Brazil:
Santarém.
LORICARIIDAE

Farlowella hasemani Eigenmann and Vance, 1917
Ann. Carnegie Mus., 11(1-2): 301.
Holotype: FMNH 55089 (CM 3781). Brazil:
Pará.
Paratype. Same locality.
LORICARIIDAE

Farlowella jauruensis Eigenmann and Vance, 1917
Ann. Carnegie Mus., 11(1-2): 300.
Holotype: FMNH 55088 (CM 3780). Brazil:
Jaurú.
LORICARIIDAE

Felichthys amblops Meek and Hildebrand, 1913
Field Mus. Nat. Hist. Zool. Ser., 10(8): 77.
Holotype: FMNH 7576. Panama: Río Tuyra,
Marriganti.
Paratypes: 3. Panama: 2 localities.
ARIIDAE

Fitzroyia eigenmanni Haseman, 1911
Ann. Carnegie Mus., 7(3-4): 385.
Holotype: FMNH 69527 (CM 2866). Brazil:
creek near Serrinha Paraná.
Paratypes: 2 (mixed with 8 topotypes). Same
locality.
JENYNSIIDAE

Flammeo scythrops Jordan and Evermann, 1903
Bull. U.S. Fish Comm. 1902, 22: 174.
Paratype. Hawaiian Islands: Honolulu.
HOLOCENTRIDAE

Formosania gilberti Oshima, 1919
Ann. Carnegie Mus., 12(2-4): 194.
Holotype: FMNH 59085 (CM 8222). Taiwan:
Tamusui River near Shinten.
CYPRINIDAE

Fowlerina franciscensis Eigenmann, 1914
Indiana Univ. Stud., 20: 45.
Holotype: FMNH 56426 (CM 5240). Brazil: La-
goas of Rio Grande, Barreiras.
Paratypes: 143. Brazil: 10 localities.
Redescribed as "*Ephippicharax franciscoen-
sis*, sp. nov." by Eigenmann and Myers
(1928, Mem. Mus. Comp. Zool. Harv.
Coll., 43(4): 506).
CHARACIDAE

Franzia ardens Jordan and Thompson, 1914
Mem. Carnegie Mus., 6(4): 251.
Holotype: FMNH 57509 (CM 6451). Japan:
Misaki.
SERRANIDAE

Fundulus cubensis Eigenmann, 1903
Bull. U.S. Fish Comm. 1902, 22: 222.
Paratype. Cuba: Pinar del Río.
CYPRINODONTIDAE

Fundulus neumanni Hilgendorf, 1905
Zool. Jahrb. Abt. Syst. Geogr. Biol. Tiere,
22(4): 417.
Syntype: FMNH 54290 (CM 2921). Tanzania:
North Ugogo.
CYPRINODONTIDAE

Fundulus oaxacae Meek, 1902
Field Mus. Nat. Hist. Zool. Ser., 3(6): 90.
Holotype: FMNH 3721. Mexico: Río Verde at
Oaxaca.
Paratypes: 40. Same locality.
CYPRINODONTIDAE

Galeichthys aguadulce Meek, 1904
Field Mus. Nat. Hist. Zool. Ser., 5: 9.
Holotype: FMNH 4678. Mexico: Río Tesecho-
cán at Pérez.
ARIIDAE

Gambusia alvarezi Hubbs and Springer, 1957
Tex. J. Sci., 9(3): 310.
Paratypes: 34. Mexico: Chihuahua, El Ojo de
San Gregorio.
POECILIIDAE

Gambusia bonita Meek, 1904

Field Mus. Nat. Hist. Zool. Ser., 5: 132.

Holotype (?): FMNH 4630. Mexico: Veracruz, Río Tonto at Refugio.

Allotype and paratypes: 57. Mexico: 2 localities. FMNH 4630 has 2 specimens. Both smaller than the published size of the holotype; thus both may be paratypes and the holotype lost or mixed with a paratype lot.

POECILIIDAE

Gambusia caliensis Eigenmann and Henn, 1916

In: Henn, Ann. Carnegie Mus., 10(1–2): 113.

Holotype: FMNH 57721 (CM 6700). Colombia: Cali.

POECILIIDAE

Gambusia cana Meek and Hildebrand, 1913

Field Mus. Nat. Hist. Zool. Ser., 10(8): 87.

Holotype: FMNH 7596. Panama: Río Satiganti, Cana.

Allotype and paratypes: 35. Same locality.

POECILIIDAE

Gambusia cascajalensis Meek and Hildebrand, 1913

Field Mus. Nat. Hist. Zool. Ser., 10(8): 86.

Holotype: FMNH 7594. Panama: Río Cascajal at Porto Bello.

Allotype and paratypes: 17. Same locality.

POECILIIDAE

Gambusia darienensis Meek and Hildebrand, 1913

Field Mus. Nat. Hist. Zool. Ser., 10(8): 88.

Holotype: FMNH 7597. Panama: Darién, Río Capetí.

Allotype and paratypes: 61. Panama: 8 localities.

POECILIIDAE

Gambusia fasciata Meek, 1904

Field Mus. Nat. Hist. Zool. Ser., 5: 129.

Holotype: FMNH 4715. Mexico: Oaxaca, San Gerónimo.

Paratypes: 19. Mexico: Oaxaca, 2 localities.

POECILIIDAE

Gambusia heterochir Hubbs, 1957

Tulane Stud. Zool., 5: 3.

Paratypes: 7. Texas: Menard County, Clear Creek.

POECILIIDAE

Gambusia hurtadoi Hubbs and Springer, 1957

Tex. J. Sci., 9(3): 307.

Paratypes: 30. Mexico: Chihuahua, El Ojo de la Hacienda Dolores.

POECILIIDAE

Gambusia infans Woolman, 1894

Bull. U.S. Fish Comm. 1894, 14: 62.

Lectoparatypes: 6. Mexico: Río de Lerma at Salamanca.

Lectotype designated by Jordan and Evermann (1896, Bull. U.S. Natl. Mus., 47(1): 680).

POECILIIDAE

Gambusia latipunctata Meek and Hildebrand, 1913

Field Mus. Nat. Hist. Zool. Ser., 10(8): 87.

Holotype: FMNH 7595. Panama: Arriján.

Paratypes: 16. Same locality.

POECILIIDAE

Gambusia nigroventralis Eigenmann and Henn, 1912

In: Eigenmann, Indiana Univ. Stud., 16: 26.

Holotype: FMNH 56045 (CM 4835). Colombia: Río San Juan at Itsmina.

Paratypes: 5. Same locality.

POECILIIDAE

Gambusia panuco Hubbs, 1926

Misc. Publ. Mus. Zool. Univ. Mich., 16: 30.

Holotype: FMNH 14060. Mexico: San Luis Potosí, Valles.

Allotype and paratypes: 8. Same locality.

POECILIIDAE

Gambusia parismina Meek, 1912

Field Mus. Nat. Hist. Zool. Ser., 10(7): 71.

Holotype: FMNH 7678. Costa Rica: Parismina.

Paratypes: 39. Same locality.

POECILIIDAE

Gambusia regani Hubbs, 1926

Misc. Publ. Mus. Zool. Univ. Mich., 16: 28.

Holotype: FMNH 14033. Mexico: Tamaulipas, Forlón.

Allotype and paratypes: 12. Same locality.

POECILIIDAE

Gambusia turrubarensis Meek, 1912

Field Mus. Nat. Hist. Zool. Ser., 10(7): 71.

Holotype: FMNH 7676. Costa Rica: Turrubares.

Allotype and paratypes: 95. Same locality.

POECILIIDAE

Gambusia umbratilis Meek, 1912

Field Mus. Nat. Hist. Zool. Ser., 10(7): 70.
Holotype: FMNH 7684. Costa Rica: Guapilis.
Paratypes: 47. Same locality.
POECILIIDAE

Gambusia vittata Hubbs, 1926

Misc. Publ. Mus. Zool. Univ. Mich., 16: 26.
Holotype: FMNH 14046. Mexico: Tamaulipas,
Forlón.
Paratypes: 12. Same locality.
POECILIIDAE

Gambusia xanthosoma Greenfield, 1983

Copeia, 1983(2): 459.
Holotype: FMNH 94188. Grand Cayman Is-
land: West Bay.
Paratypes: 10. Same locality.
POECILIIDAE

Gambusia yucatana australis Greenfield, 1985

Copeia, 1985(2): 375.
Holotype: FMNH 94901. Belize: Saint George's
Cay, Barrier Reef.
Paratypes: 58. Belize: 4 localities.
POECILIIDAE

Gastromyzon fasciatus Inger and Chin, 1961

Copeia, 1961(2): 173.
Holotype: FMNH 68119. Malaysia: Sarawak,
Sungei Dapu River near mouth of Sungei
Putai.
Paratypes: 29. Malaysia: Sarawak, 3 localities.
HOMALOPTERIDAE

Gastromyzon lepidogaster Roberts, 1982

Proc. Calif. Acad. Sci., 42(20): 509.
Paratypes: 8. Malaysia: Sabah, 2 localities.
HOMALOPTERIDAE

Gastromyzon megalepis Roberts, 1982

Proc. Calif. Acad. Sci., 42(20): 510.
Holotype: FMNH 68126. Malaysia: Sarawak,
tributary of Baleh River.
Paratypes: 8. Malaysia: Sarawak, 2 localities.
HOMALOPTERIDAE

Gastromyzon pauciradiatus Inger and Chin, 1961

Copeia, 1961(2): 174.
Holotype: FMNH 68121. Malaysia: Sarawak,
tributary of Baleh River.
Paratypes: 21. Malaysia: Sarawak, 2 localities.
HOMALOPTERIDAE

Gastromyzon punctulatus Inger and Chin, 1961

Copeia, 1961(2): 173.
Holotype: FMNH 68116. Malaysia: Sarawak,
Sungei Dapu River near mouth of Sungei
Putai.
Paratypes: 16. Malaysia: Sarawak, 3 localities.
HOMALOPTERIDAE

Gastromyzon ridens Roberts, 1982

Proc. Calif. Acad. Sci., 42(20): 515.
Paratype. Indonesia: Kapuas River basin, Sun-
gei Pinoh.
HOMALOPTERIDAE

Gazza achlamys Jordan and Starks, 1917

Ann. Carnegie Mus., 11: 446.
Holotype: FMNH 58939 (CM 8074). Sri Lanka:
Colombo.
Paratypes: 2. Same locality.
LEIOGNATHIDAE

Gelanoglanis stroudi Böhlke, 1980

Proc. Acad. Nat. Sci. Phila., 132: 152.
Paratypes: 6. Colombia: Río Manacacias near
Puerto Gaitán.
AUCHENIPTERIDAE

Genycharax tarpon Eigenmann, 1912

Indiana Univ. Stud., 16: 22.
Holotype: FMNH 56018 (CM 4808). Colombia:
Cartago.
Paratypes: 15. Colombia: 3 localities.
CHARACIDAE

Geophagus brasiliensis iporangensis Haseman,
1911

Ann. Carnegie Mus., 7(3-4): 364.
Holotype: FMNH 54202 (CM 2792). Brazil:
Iporanga.
Paratypes: 3. Same locality.
CICHLIDAE

Geophagus brasiliensis itapicuruensis Haseman,
1911

Ann. Carnegie Mus., 7(3-4): 365.
Holotype: FMNH 54365 (CM 2793). Brazil: Rio
Itapicuru at Queimadas.
Paratypes: 7. Brazil: 3 localities.
CICHLIDAE

Gephyrocharax caucanus Eigenmann, 1912

Indiana Univ. Stud., 16: 24.
Holotype: FMNH 56012 (CM 4802). Colombia:
Cartago.
Paratypes: 37. Colombia: 3 localities.
CHARACIDAE

- Gephyrocharax chocoensis** Eigenmann, 1912
Indiana Univ. Stud., 16: 23.
Holotype: FMNH 56016 (CM 4806). Colombia: Istmina.
Paratypes: 25. Same locality.
CHARACIDAE
- Gephyrocharax intermedius** Meek and Hildebrand, 1916
Field Mus. Nat. Hist. Zool. Ser., 10(15): 278.
Holotype: FMNH 8945. Panama: Río Chame.
Paratypes: 28. Same locality.
CHARACIDAE
- Gephyrocharax melanocheir** Eigenmann, 1912
Indiana Univ. Stud., 16: 24.
Holotype: FMNH 56049 (CM 4839). Colombia: Bernal Creek near Honda.
Paratypes: 19. Colombia: 2 localities.
CHARACIDAE
- Gephyrocharax whaleri** Hildebrand, 1938
Field Mus. Nat. Hist. Zool. Ser., 22(4): 254.
Paratypes: 8. Panama: between Campana and La Venta.
CHARACIDAE
- Gerhardinus nudus** Meek and Hildebrand, 1928
Field Mus. Nat. Hist. Zool. Ser., 15(3): 889.
Paratypes: 7. Panama: tide pools at Panama City.
GOBIIDAE
- Gibbonsia elegans erroli** Hubbs, 1952
Stanford Ichthyol. Bull., 4(2): 151.
Paratypes: 6. Mexico: Baja California, Isla Guadalupe.
CLINIDAE
- Gibbonsia elegans montereyensis** Hubbs, 1927
Pap. Mich. Acad. Sci. Arts and Lett., 7: 354.
Paratypes: 8. California: 2 localities.
CLINIDAE
- Gibbonsia elegans velifera** Hubbs, 1952
Stanford Ichthyol. Bull., 4(2): 149.
Paratypes: 17. California: 2 localities.
CLINIDAE
- Gibbonsia metzi** Hubbs, 1927
Pap. Mich. Acad. Sci. Arts and Lett., 7: 360.
Paratypes: 8. California: 4 localities.
CLINIDAE
- Gibbonsia norae** Hubbs, 1952
Stanford Ichthyol. Bull., 4(2): 126.
Paratypes: 9. Mexico: Baja California, Isla Guadalupe.
CLINIDAE
- Gila minacae** Meek, 1902
Field Mus. Nat. Hist. Zool. Ser., 3(6): 80.
Holotype: FMNH 3573. Mexico: Río Papigochic at Miñaca.
CYPRINIDAE
- Gillellus greyae** Kanazawa, 1952
Fieldiana Zool., 34(7): 88.
Paratypes: 2. Bermuda and Florida: 2 localities.
DACTYLOSCOPIDAE
- Girardinus garmani** Eigenmann, 1903
Bull. U.S. Fish Comm. 1902, 22: 226.
Paratype: Cuba: Palacios.
POECILIIDAE
- Gladiogobius ensifer** Herre, 1933
Copeia, 1933(1): 23.
Holotype: FMNH 17412. Indonesia: Waigiu Island.
GOBIIDAE
- Glandulocauda inequalis** Eigenmann, 1911
Ann. Carnegie Mus., 8(1): 169.
Holotype: FMNH 54893 (CM 3555). Brazil: Porto Alegre.
Paratypes: 4. Same locality.
CHARACIDAE
- Glandulocauda melanogenys** Eigenmann, 1911
Ann. Carnegie Mus., 8(1): 168.
Holotype: FMNH 54891 (CM 3553). Brazil: São Paulo, Alto da Serra.
Paratypes: 20. Same locality.
CHARACIDAE
- Glandulocauda melanopleura** Eigenmann, 1911
Ann. Carnegie Mus., 8(1): 170.
Holotype: FMNH 54895 (CM 3557). Brazil: Rio Iguaçu at Serrinha Paraná.
Paratypes: 3. Same locality.
CHARACIDAE
- Glanidium ribeiroi** Haseman, 1911
Ann. Carnegie Mus., 7(3–4): 381.
Holotype: FMNH 54253 (CM 2877). Brazil: Rio Iguaçu, Porto União da Victoria.
Paratypes: 5. Same locality.
AUCHENIPTERIDAE

- Glaridichthys falcatus** Eigenmann, 1903
Bull. U.S. Fish Comm. 1902, 22: 224.
Paratypes: 4. Cuba: 3 localities.
POECILIIDAE
- Glossanodon pygmaeus** Cohen, 1958
Bull. Fla. State Mus. Biol. Sci., 3(3): 156.
Holotype: FMNH 64345. Gulf of Mexico: off Tortugas.
Paratypes: 7. Gulf of Mexico: 3 localities.
ARGENTINIDAE
- Glossogobius abacopus** Jordan and Richardson, 1909
Mem. Carnegie Mus., 4(4): 200.
Holotype: FMNH 52210 (CM 357). Taiwan: Takao.
Paratype: Same locality.
GOBIIDAE
- Glossogobius koragensis** Herre, 1935
Field Mus. Nat. Hist. Zool. Ser., 18(12): 419.
Holotype: FMNH 17365. New Guinea: Sepik River at Koragu.
Paratype: New Guinea: Kerame River at Ambot.
GOBIIDAE
- Glossogobius parvus** Oshima, 1919
Ann. Carnegie Mus., 12(2-4): 305.
Holotype: FMNH 59138 (CM 8276). Taiwan: Kizanto Island near Giran.
GOBIIDAE
- Glossogobius sandakanensis** Inger, 1957
Fieldiana Zool., 36(3): 393.
Holotype: FMNH 59526. Malaysia: Sabah, Mankala River at Sandakan.
GOBIIDAE
- Glyphisodon troscheli** Gill, 1863
Proc. Acad. Nat. Sci. Phila., 1862: 150.
Paratype: Mexico: Baja California, Cabo San Lucas.
Misspelled *Glyphidodon* in Gill (1863).
POMACENTRIDAE
- Glyptothorax housei** Herre, 1942
Stanford Ichthyol. Bull., 2(4): 117.
Paratypes: 3. India: stream in Anaimallai Hills, east of Valparai.
SISORIDAE
- Gnathodentex oculomaculatus** Herre, 1935
Field Mus. Nat. Hist. Zool. Ser., 18(12): 407.
Holotype: FMNH 17335. Solomon Islands: Isabel Island, Tenibuli.
Paratype: Same locality.
LETHRINIDAE
- Gnatholepis corlettei** Herre, 1935
Field Mus. Nat. Hist. Zool. Ser., 18(12): 418.
Holotype: FMNH 17367. New Hebrides Islands: Malekula Island, Bushman's Bay.
Paratypes: 3. Same locality.
GOBIIDAE
- Gnatholepis knighti** Jordan and Evermann, 1903
Bull. U.S. Fish Comm. 1902, 22: 204.
Paratypes: 5. Hawaiian Islands: Hilo.
GOBIIDAE
- Gnathopogon iijimae** Oshima, 1919
Ann. Carnegie Mus., 12(2-4): 219.
Holotype: FMNH 59096 (CM 8234). Taiwan: Tozen River.
CYPRINIDAE
- Gnathopogon ishikawae** Jordan and Thompson, 1914
Mem. Carnegie Mus., 6(4): 215.
Paratypes: 3. Japan: Chikugo River at Kurume.
FMNH 57077 (CM 6014), the published number for the lot which included the holotype has 3 specimens, all too small to be the holotype; perhaps the holotype went to Stanford University.
CYPRINIDAE
- Gnathopogon longifilis** Jordan and Hubbs, 1925
Mem. Carnegie Mus., 10(2): 169.
Holotype: FMNH 58711 (CM 7817). Korea: Ping-yang River.
Published number CM 7819 is an error.
CYPRINIDAE
- Ganthopogon majimae** Jordan and Hubbs, 1925
Mem. Carnegie Mus., 10(2): 167.
Holotype: FMNH 58710 (CM 7816). Korea: Ping-yang River.
CYPRINIDAE
- Gnathopogon suwae** Jordan and Hubbs, 1925
Mem. Carnegie Mus., 10(2): 166.
Holotype: FMNH 58708 (CM 7814). Japan: Lake Suwa at Kamisuwa.
CYPRINIDAE

- Gnathopogon tsuchigae** Jordan and Hubbs, 1925
Mem. Carnegie Mus., 10(2): 170.
Holotype: FMNH 58712 (CM 7818). Korea:
Ping-yang River.
CYPRINIDAE
- Gobiella birtwistlei** Herre, 1934
Bull. Raffles Mus., 9: 85.
Syntypes: 18, FMNH 47158 and 47230. Singapore.
GOBIIDAE
- Gobiesox costaricensis** Meek, 1912
Field Mus. Nat. Hist. Zool. Ser., 10(7): 74.
Holotype: FMNH 7677. Costa Rica: Zent.
Paratypes: 4. Costa Rica: 2 localities.
GOBIESOCIDAE
- Gobiesox fluviatilis** Briggs and Miller, 1960
Occas. Pap. Mus. Zool. Univ. Mich., 616: 2.
Paratype. Mexico: Jalisco, Barranca de Barranquitas.
GOBIESOCIDAE
- Gobiesox fulvus** Meek, 1907
Field Mus. Nat. Hist. Zool. Ser., 7(5): 149.
Holotype: FMNH 6035. Costa Rica: Cocos Island.
GOBIESOCIDAE
- Gobiesox marijeanae** Briggs, 1960
Copeia, 1960(3): 215.
Paratype. Mexico: Islas Tres Marías, Isla San Juanito.
GOBIESOCIDAE
- Gobiesox mexicanus** Briggs and Miller, 1960
Occas. Pap. Mus. Zool. Univ. Mich., 616: 7.
Paratype. Mexico: Oaxaca, Río Tonameca.
GOBIESOCIDAE
- Gobiesox paradiseus** Herre, 1935
Field Mus. Nat. Hist. Zool. Ser., 18(12): 432.
Holotype: FMNH 17404. Ecuador: Islas Galápagos, Isla Eden.
GOBIESOCIDAE
- Gobiesox potamius** Briggs, 1955
Stanford Ichthyol. Bull., 6: 93.
Holotype: FMNH 7815. Costa Rica: Turru-bares.
Paratype. Same locality.
Both are from the paratype series of *Gobiesox costaricensis*.
GOBIESOCIDAE
- Gobiocichla ethelwynnae** Roberts, 1982
Proc. K. Ned. Akad. Wet. Ser. C Biol. Med. Sci., 85(4): 577.
Paratypes: 15. Cameroon: Cross River, downstream from Mamfé.
CICHLIDAE
- Gobiocichla wonderi** Kanazawa, 1951
Ann. Mag. Nat. Hist., Ser. 12, 4: 378.
Holotype: FMNH 50423. Mali: Niger River at Sotuba.
CICHLIDAE
- Gobionellus saepepallens** Gilbert and Randall, 1968
Not. Nat. (Phila.), 411: 2.
Paratypes: 2. The Bahamas: Green Turtle Cay, east of Little Abaco.
GOBIIDAE
- Gobiopsis aporia** Lachner and McKinney, 1978
Smithson. Contrib. Zool., 262: 12.
Paratype. Sri Lanka: Galle.
GOBIIDAE
- Gobiopsis woodsi** Lachner and McKinney, 1978
Smithson. Contrib. Zool., 262: 21.
Paratypes: 39. India: Gulf of Mannar, Musal Tivu.
GOBIIDAE
- Gobio coriparoides** Nichols, 1925
Am. Mus. Nov., 181: 4.
Paratypes: 2. China: Shansi, Chin-ssu.
CYPRINIDAE
- Gobio kesleri banaticus** Banarescu, 1953
Stud. Cercet. Stiint., 4(1-2): 300.
Paratypes: 22. Romania: Banat, Timis River.
CYPRINIDAE
- Gobio rivuloides** Nichols, 1925
Am. Mus. Nov., 181: 5.
Paratypes: 3. China: Shansi, Niang-tze-kwan.
CYPRINIDAE
- Gobio soldatovi minulus** Nichols, 1925
Am. Mus. Nov., 181: 3.
Paratypes: 4. China: Shansi, Kweihwa.
CYPRINIDAE
- Gobius claytonii** Meek, 1902
Field Mus. Nat. Hist. Zool. Ser., 3(6): 121.
Holotype: FMNH 3740. Mexico: Río San Francisco at La Antigua.
Paratypes: 8. Same locality.
GOBIIDAE

- Gobius cliffordpopei** Nichols, 1925
Am. Mus. Nov., 185: 5.
Paratypes: 4. China: Hunan, Tungting Lake.
GOBIIDAE
- Gobius daguae** Eigenmann, 1917
Proc. Am. Philos. Soc., 56(7): 685.
Holotype: FMNH 58479 (CM 7481). Colombia:
mouth of Río Dagua.
Paratype. Same locality.
GOBIIDAE
- Gobius parvus** Meek, 1902
Field Mus. Nat. Hist. Zool. Ser., 3(6): 121.
Holotype: FMNH 3738. Mexico: Río San Francisco at La Antigua.
Paratypes: 29. Same locality.
GOBIIDAE
- Gonostoma ebelingi** Grey, 1960
Bull. Mus. Comp. Zool. Harv. Coll., 122(2): 109.
Paratype. Western Pacific Ocean: Marshall Islands area.
CLUPEIDAE
- Goodea captiva** Hubbs, 1924
Occas. Pap. Mus. Zool. Univ. Mich., 148: 4.
Holotype: FMNH 5557. Mexico: San Luis Potosí, Jesús María.
Paratypes: 172. Same locality.
GOODEIDAE
- Goodea toweri** Meek, 1904
Field Mus. Nat. Hist. Zool. Ser., 5: 138.
Holotype: FMNH 4519. Mexico: San Luis Potosí, Río Verde.
Paratypes: 26. Same locality.
GOODEIDAE
- Goodea whitei** Meek, 1904
Field Mus. Nat. Hist. Zool. Ser., 5: 137.
Holotype: FMNH 4547. Mexico: Morelos, Yau-tepec.
Paratypes: 420. Mexico: 2 localities.
GOODEIDAE
- Grammonus mowbrayi** Grey, 1951
Copeia, 1951(2): 154.
Holotype: FMNH 49185. Western North Atlantic: off Bermuda.
Paratypes: 4. Same locality.
OPHIDIIDAE
- Grasseichthys gabonensis** Géry, 1964
C. R. Seances Acad. Sci. Paris, 259: 4806.
Paratypes: 6. Gabon: tributary of Ntsimy River near Nzingmeyong.
KNERIIDAE
- Grimatrocetes bullisi** Grey, 1958
Fieldiana Zool., 39(16): 150.
Paratypes: 3. Gulf of Mexico.
ALEPOCEPHALIDAE
- Gunnelllichthys viridesens** Dawson, 1968
Proc. Biol. Soc. Wash., 81: 61.
Paratype. Maldives: Addu Atoll, Bushy Island.
MICRODESMIDAE
- Gymnapogon gracilicauda** Lachner, 1953
Bull. U.S. Natl. Mus., 202(1): 497.
Paratype. Marshall Islands: Bikini Atoll, Airi Island.
APOGONIDAE
- Gymnapogon urospilotes** Lachner, 1953
Bull. U.S. Natl. Mus., 202(1): 494.
Paratypes: 2. Marshall Islands: 2 localities.
APOGONIDAE
- Gymnorhamphichthys hypostomus** Ellis, 1912
Mem. Carnegie Mus., 5: 436.
Holotype: FMNH 54554 (CM 3182). Bolivia: San Joaquín.
Paratypes: 4. Brazil: 3 localities.
RHAMPHICHTHYIDAE
- Gymnothorax bayeri** Schultz, 1953
Bull. U.S. Natl. Mus., 202(1): 124.
Paratype. Marshall Islands: Bikini Atoll, Erik Island.
MURAENIDAE
- Gymnothorax bikiniensis** Schultz, 1953
Bull. U.S. Natl. Mus., 202(1): 116.
Paratype. Marshall Islands: Bikini Atoll, Boro Island.
MURAENIDAE
- Gymnothorax efatensis** Herre, 1935
Field Mus. Nat. Hist. Zool. Ser., 18(12): 384.
Holotype: FMNH 17193. New Hebrides Islands: Efate Island, Vila.
MURAENIDAE

Gymnothorax leucostigma Jordan and Richardson, 1909

Mem. Carnegie Mus., 4(4): 174.

Holotype: FMNH 52124 (CM 253). Taiwan: Takao.

MURAENIDAE

Hadropterus scierus apristis Hubbs and Hubbs, 1954

In: Hubbs, Am. Midl. Nat., 52(1): 211.

Paratypes: 13. Texas: San Marcos River.

PERCIDAE

Halichoeres fijiensis Herre, 1935

Field Mus. Nat. Hist. Zool. Ser., 18(12): 410.

Holotype: FMNH 17341. Fiji Islands: Ovalau Island.

Paratypes: 3. Same locality.

LABRIDAE

Hampala macrolepidota sabana Inger and Chin, 1962

Fieldiana Zool., 45: 81.

Holotype: FMNH 68218. Malaysia: Sabah, tributary of Kinabatangan River near Deramakot.

Paratypes: 134. Malaysia: Sabah, 5 localities.

CYPRINIDAE

Hasemania bilineata Ellis, 1911

Ann. Carnegie Mus., 8(2): 150.

Holotype: FMNH 54383 (CM 3001). Brazil: São Paulo, Alto da Serra.

Paratypes: 6. Brazil: Rio Tietê at Mogi das Cruces.

CHARACIDAE

Hasemania maxillaris Ellis, 1911

Ann. Carnegie Mus., 8(2): 148.

Holotype: FMNH 54303 (CM 2937). Brazil: Rio Iguaçu at Porto União.

CHARACIDAE

Hasemania melanura Ellis, 1911

Ann. Carnegie Mus., 8(2): 149.

Holotype: FMNH 54384 (CM 3002). Brazil: Rio Iguaçu at Porto União.

Paratypes: 39. Same locality.

CHARACIDAE

Helogenes unidorsalis Glodek and Carter, 1978

Fieldiana Zool., 72(6): 77.

Holotype: FMNH 80463. Ecuador: Río Bobonaza between Sarayacu and Montalvo.

Paratypes: 5. Same locality.

HELOGENIDAE

Helotes lorentzi Weber, 1910

Notes Leiden Mus., 32: 236.

Syntype: FMNH 52495 (CM 753). New Guinea: Lorentz River.

TERAPONIDAE

Hemiancistrus daguae Eigenmann, 1912

Indiana Univ. Stud., 16: 11.

Holotype: FMNH 56052 (CM 4842). Colombia: Caldas.

Paratypes: 11. Colombia: 2 localities.

LORICARIIDAE

Hemiancistrus mayoloi Eigenmann, 1912

Indiana Univ. Stud., 16: 10.

Holotype: FMNH 56036 (CM 4826). Colombia: Istmina.

Paratypes: 2. Same locality.

LORICARIIDAE

Hemiancistrus wilsoni Eigenmann, 1917

Proc. Am. Philos. Soc., 56(7): 678.

Holotype: FMNH 58550 (CM 7570). Colombia: Truandó.

Paratypes: 3. Same locality.

LORICARIIDAE

Hemibrycon boquiae Eigenmann, 1913

Indiana Univ. Stud., 18: 20.

Holotype: FMNH 56259 (CM 5059). Colombia: Boquia.

Paratypes: 10. Same locality.

Published originally as "*Bryconamericus* or *Hemibrycon*".

CHARACIDAE

Hemibrycon colombianus Eigenmann, 1914

In: Eigenmann, Henn, and Wilson, Indiana Univ. Stud., 19: 8.

Holotype: FMNH 56653 (CM 5470). Colombia: Río San Gil, Santander.

Paratypes: 7. Colombia: 3 localities.

CHARACIDAE

Hemibrycon dariensis Meek and Hildebrand, 1916

Field Mus. Nat. Hist. Zool. Ser., 10(15): 285.

Holotype: FMNH 8947. Panama: Río Tuyra basin, Río Yape.

Paratypes: 28. Panama: 7 localities.

CHARACIDAE

Hemibrycon helleri Eigenmann, 1927

Mem. Mus. Comp. Zool. Harv. Coll., 43(4): 406.

Paratypes: 3. Peru: Río Comerciato.

CHARACIDAE

Hemicarax rhomboides Meek and Hildebrand, 1925

Field Mus. Nat. Hist. Zool. Ser., 15(2): 345.
Paratype. Panama: Fox Bay, Colón.
CARANGIDAE

Hemicetopsis amphioxus Eigenmann, 1914

In: Eigenmann, Henn, and Wilson, Indiana Univ. Stud., 19: 14.
Holotype: FMNH 56519 (CM 5332). Colombia: Río Patía basin, creek near San Lorenzo.
CETOPSIDAE

Hemicetopsis macilentus Eigenmann, 1912

Mem. Carnegie Mus., 5: 211.
Holotype: FMNH 53260 (CM 1726). Guyana: creek below Potaro Landing.
CETOPSIDAE

Hemicetopsis minutus Eigenmann, 1912

Mem. Carnegie Mus., 5: 211.
Holotype: FMNH 53262 (CM 1728). Guyana: Amatuk.
CETOPSIDAE

Hemicetopsis othonops Eigenmann, 1912

Indiana Univ. Stud., 16: 17.
Holotype: FMNH 56040 (CM 4830). Colombia: Girardot.
Paratypes: 16. Colombia: 2 localities.
CETOPSIDAE

Hemiculterella engraulis Nichols, 1925

Am. Mus. Nov., 182: 7.
Paratypes: 2. China: Hunan, Tungting Lake.
CYPRINIDAE

Hemidoras leporhinus Eigenmann, 1912

Mem. Carnegie Mus., 5: 195.
Holotype: FMNH 53185 (CM 1624). Guyana: Tumatumari.
Paratype. Guyana: Crab Falls.
DORADIDAE

Hemidoras micropoeus Eigenmann, 1912

Mem. Carnegie Mus., 5: 195.
Holotype: (CM 1636) MISSING. Guyana: Wismar.
Paratype. Guyana: Lama Stop-off.
DORADIDAE

Hemidoras microstomus Eigenmann, 1912

Mem. Carnegie Mus., 5: 193.
Holotype: FMNH 53206 (CM 1650). Guyana: Rockstone.
Paratypes: 3. Guyana: Essequibo River, Crab Falls.
DORADIDAE

Hemidoras notospilus Eigenmann, 1912

Mem. Carnegie Mus., 5: 196.
Holotype: FMNH 53184 (CM 1623). Guyana: Crab Falls.
DORADIDAE

Hemieleotris levis Eigenmann, 1917

Proc. Am. Philos. Soc., 56(7): 684.
Paratypes: 5. Colombia: pools in Buenaventura.
ELEOTRIDAE

Hemigrammus analis Durbin, 1909

Ann. Carnegie Mus., 6(1): 64.
Holotype: FMNH 53041 (CM 1466). Guyana: Rockstone.
Paratypes: 45. Guyana: 3 localities.
CHARACIDAE

Hemigrammus brevis Ellis, 1911

Ann. Carnegie Mus., 8(2): 161.
Holotype: FMNH 54444 (CM 3068). Brazil: Lagoas of Rio Grande at Barreiras.
Paratypes: 37. Brazil: 3 localities.
CHARACIDAE

Hemigrammus compressus Meek, 1904

Field Mus. Nat. Hist. Zool. Ser., 5: 87.
Holotype: FMNH 4641. Mexico: Río Papaloapan basin at El Hule.
Paratypes: 28. Mexico: 2 localities.
CHARACIDAE

Hemigrammus cylindricus Durbin, 1909

Ann. Carnegie Mus., 6(1): 62.
Holotype: FMNH 53426 (CM 1461). Guyana: Tumatumari.
Paratypes: 20. Guyana: 4 localities.
CHARACIDAE

Hemigrammus erythrozonus Durbin, 1909

Ann. Carnegie Mus., 6(1): 56.
Holotype: FMNH 53546 (CM 1448). Guyana: Erukin.
Paratypes: 7. Same locality.
CHARACIDAE

Hemigrammus iota Durbin, 1909

Ann. Carnegie Mus., 6(1): 60.
Holotype: FMNH 53034 (CM 1458). Guyana: Gluck Island.
Paratypes: 4. Guyana: 2 localities.
CHARACIDAE

- Hemigrammus kennedyi** Eigenmann, 1903
Proc. Acad. Nat. Sci. Phila., 59: 520.
Paratypes: 4. Paraguay: Campo Grande.
CHARACIDAE
- Hemigrammus marginatus** Ellis, 1911
Ann. Carnegie Mus., 8(2): 159.
Holotype: FMNH 54430 (CM 3053). Brazil: Rio Itapicurú at Queimadas.
Paratypes: 130. Brazil and Paraguay: 18 localities.
CHARACIDAE
- Hemigrammus melanopterus** Eigenmann and Kennedy, 1903
Proc. Acad. Nat. Sci. Phila., 59: 518.
Paratypes: 2. Paraguay: Arroyo Pypucú.
Spelled *melasopterus* in Eigenmann and Kennedy (1903), apparently a typographic error.
CHARACIDAE
- Hemigrammus minutus** Meek and Hildebrand, 1912
Field Mus. Nat. Hist. Zool. Ser., 10(6): 67.
Holotype: FMNH 7572 MISSING or mixed.
Panama: Canal Zone, Río Agua Clara.
Paratypes: 71. Panama: 7 localities.
CHARACIDAE
- Hemigrammus orthus** Durbin, 1909
Ann. Carnegie Mus., 6(1): 61.
Holotype: FMNH 53052 (CM 1477). Guyana: Tukeit.
Paratypes: 19. Guyana: 3 localities.
CHARACIDAE
- Hemigrammus riddlei** Meek, 1907
In: Eigenmann and Ogle, Proc. U.S. Natl. Mus., 33(1556): 13.
Holotype: MISSING. Venezuela: Los Castillos.
See comment in Grey (1947, p. 156); this was published as at the Field Museum, but neither the catalogue entry nor the fish have been found.
CHARACIDAE
- Hemigrammus rodwayi** Durbin, 1909
Ann. Carnegie Mus., 6(1): 58.
Holotype: FMNH 53026 (CM 1450). Guyana: Georgetown trenches.
Paratypes: 248. Guyana: 4 localities.
CHARACIDAE
- Hemiodus orthonops** Eigenmann and Kennedy, 1903
Proc. Acad. Nat. Sci. Phila., 59: 511.
Paratype. Paraguay: Asunción.
HEMIODONTIDAE
- Hemiodus parnaguae** Eigenmann and Henn, 1916
Ann. Carnegie Mus., 10(1–2): 87.
Holotype: FMNH 56886 (CM 5701). Brazil: Lagoa de Parnagua.
Paratypes: 2. Same locality.
HEMIODONTIDAE
- Hemiodus quadrimaculatus** Pellegrin, 1908
Bull. Mus. d'Hist. Nat., 14(7): 343.
Paratype: MISSING. French Guiana: Camopi River.
HEMIODONTIDAE
- Hemipimelodus papillifer** Herre, 1935
Field Mus. Nat. Hist. Zool. Ser., 18(12): 390.
Holotype: FMNH 17211. New Guinea: Sepik River at Timbunke.
Paratype. Same locality.
ARIIDAE
- Hemipimelodus velutinus** Weber, 1908
Nova Guinea, 5(2): 225.
Syntype: FMNH 52392 (CM 648). New Guinea: Tawarin River.
ARIIDAE
- Hemipteronotus baldwini** Jordan and Evermann, 1903
Bull. U.S. Fish Comm. 1902, 22: 192.
Paratypes: 2. Hawaiian Islands: Honolulu.
LABRIDAE
- Hemipteronotus evides** Jordan and Richardson, 1909
Mem. Carnegie Mus., 4(4): 196.
Holotype: FMNH 52197 (CM 343). Taiwan: Takao.
LABRIDAE
- Hemiramphus archipelagicus** Collette, 1978
Proc. Biol. Soc. Wash., 91(3): 732.
Paratypes: 2. Indonesia: Java, Batavia.
HEMIRAMPHIDAE
- Hemiramphus bermudensis** Collette, 1962
Bull. Mar. Sci. Gulf Caribb., 12(3): 432.
Holotype: FMNH 48115. Bermuda: Saint George's Island, Whalebone Bay.
Paratypes: 6. Bermuda: 6 localities.
HEMIRAMPHIDAE

Heptapterus stewarti Haseman, 1911

Ann. Carnegie Mus., 7(3-4): 376.

Holotype: FMNH 54234 (CM 2850). Brazil:
creek at Serrinha Paraná.

PIMELODIDAE

Heros beani Jordan, 1889

Proc. U.S. Natl. Mus. 1888, 11: 332.

Paratype. Mexico: Río Presidio, Mazatlán.

CICHLIDAE

Heterandria colombianus Eigenmann and Henn,
1912

In: Eigenmann, Indiana Univ. Stud., 16: 27.

Holotype: FMNH 56047 (CM 4837). Colombia:
mouth of Río Dagua.

Paratypes: 2. Same locality.

POECILIIDAE

Heterandria cubensis Eigenmann, 1903

Bull. U.S. Natl. Mus. 1902, 22: 227.

Paratype. Cuba: Pinar del Río.

POECILIIDAE

Heterandria hasemani Henn, 1916

Ann. Carnegie Mus., 10(1-2): 116.

Holotype: FMNH 55874 (CM 4663). Bolivia:
Puerto Suárez.

Paratypes: 3. Same locality.

POECILIIDAE

Heterandria lutzi Meek, 1902

Field Mus. Nat. Hist. Zool. Ser., 3(6): 106.

Holotype: FMNH 3718. Mexico: Río Quioatepec
at Cuicatlán.

Paratypes: 381. Mexico: 3 localities.

POECILIIDAE

Heterenchelys biaggii Böhlke, 1956

Not. Nat. (Phila.), 289: 8.

Holotype: FMNH 61774. Puerto Rico: Maya-
güez pier.

Paratype. Same locality.

HETERENCHELYIDAE

Heterocharax macrolepis Eigenmann, 1912

Mem. Carnegie Mus., 5: 406.

Holotype: FMNH 53669 (CM 2142). Guyana:
Rockstone.

Paratypes: 2. Guyana: 2 localities.

CHARACIDAE

Heterogramma corumbae Eigenmann and Ward,
1907

Ann. Carnegie Mus., 4(2): 146.

Paratype. Paraguay: Corumbá.

CICHLIDAE

Heterogramma ortmanni Eigenmann, 1912

Mem. Carnegie Mus., 5: 506.

Holotype: FMNH 53801 (CM 2306). Guyana:
Erukin.

Paratypes: 25. Guyana: 6 localities.

CICHLIDAE

Heterogramma ritense Haseman, 1911

Ann. Carnegie Mus., 7(3-4): 362.

Holotype: FMNH 54187 (CM 2765). Brazil: Rio
Santa Rita at Santa Rita.

Paratypes: 3. Same locality.

CICHLIDAE

Heterogramma taeniatum pertense Haseman, 1911

Ann. Carnegie Mus., 7(3-4): 359.

Holotype: FMNH 54171 (CM 2741). Brazil:
Manaus.

Paratype: MISSING. Brazil: Rio Tapajós at
Santarém.

CICHLIDAE

Heterogramma trifasciatum maciliense Haseman,
1911

Ann. Carnegie Mus., 7(3-4): 360.

Syntypes: 2, FMNH 54177 (CM 2751) MISS-
ING. Brazil: São Antonio de Guaporé.

CICHLIDAE

Hildadoras boliviensis Fernández-Yépez, 1968

Bol. Inst. Oceanogr. Univ. Oriente, 7(1): 43.

Holotype: FMNH 84067. Venezuela: Orinoco
system, Quebrada El Pilón.

Paratype. Same locality.

DORADIDAE

Hippocampus brunneus Bean, 1906

Proc. Biol. Soc. Wash., 19: 32.

Holotype: FMNH 5494. Bermuda: Long Bird
Island.

SYNGNATHIDAE

Hippocampus fisheri Jordan and Evermann, 1903

Bull. U.S. Fish Comm. 1902, 22: 169.

Paratype. Hawaiian Islands: Hilo.

SYNGNATHIDAE

Holacanthus lunatus Blosser, 1909

Ann. Carnegie Mus., 6(1): 299.

Holotype: FMNH 52831 (CM 1235). Lesser An-
tilles: Saint Croix Island.

Misspelled *Holocanthus* in Blosser (1909).

POMACANTHIDAE

Holesthes heterodon Eigenmann, 1915

Mem. Carnegie Mus., 7(1): 87.

Holotype: FMNH 57890 (CM 6875). Brazil: Jaguará.

Paratypes: 86. Brazil: 13 localities.

CHARACIDAE

Holocentrus bullisi Woods, 1955

Fieldiana Zool., 37(4): 113.

Paratypes: 16. Gulf of Mexico and North Atlantic: 6 localities.

HOLOCENTRIDAE

Holocentrus ensifer Jordan and Evermann, 1903

Bull. U.S. Fish Comm. 1902, 22: 176.

Paratype. Hawaiian Islands: Honolulu.

HOLOCENTRIDAE

Holocentrus meeki Bean, 1906

Proc. Biol. Soc. Wash., 19: 31.

Lectotype: FMNH 5079. Bermuda: Saint David's Island, Hayward's Bay.

Lectoparatype. Same locality.

Lectotype designated by Grey (1947, p. 157).

HOLOCENTRIDAE

Holocentrus xantherythrus Jordan and Evermann, 1903

Bull. U.S. Fish Comm. 1902, 22: 175.

Paratypes: 2. Hawaiian Islands: Honolulu.

HOLOCENTRIDAE

Hololepis saludae Hubbs and Cannon, 1935

Misc. Publ. Mus. Zool. Univ. Mich., 30: 50.

Paratype. South Carolina: Saluda County, Richland Creek.

PERCIDAE

Holopristis riddlei Meek, 1907

In: Eigenmann and Ogle, Proc. U.S. Natl. Mus., 33(1556): 11.

Paratypes: 3. Venezuela: Los Castillos.

See comment in Grey (1947, p. 158); misspelled *Holopristes* in Meek (1907).

CHARACIDAE

Homaloptera hingi Herre, 1934

Lingnan Sci. J. Canton, 13(2): 287.

Paratypes: 4. Hong Kong: stream at Pok Fu Lam.

HOMALOPTERIDAE

Hoplomyzon papillatus Stewart, 1985

Fieldiana Zool., New Ser., 25: 8.

Holotype: FMNH 94908. Ecuador: Río Aguarico, 1 km upstream from confluence with Río Shushufindi.

ASPREDINIDAE

Hoplosternum magdalenae Eigenmann, 1913

In: Ellis, Ann. Carnegie Mus., 8(3-4): 412.

Holotype: FMNH 56281 (CM 5081). Colombia: Soplaviento.

Paratypes: 12. Colombia: 2 localities.

CALLICHTHYIDAE

Hoplosternum punctatum Meek and Hildebrand, 1916

Field Mus. Nat. Hist. Zool. Ser., 10(15): 264.

Holotype: FMNH 8943. Panama: Río Marte. Armade east of Panama City.

Paratype. Same locality.

CALLICHTHYIDAE

Hoplostethus occidentalis Woods, 1973

Mem. Sears Fnd. Mar. Res., 1(6): 319.

Holotype: FMNH 66790. Gulf of Mexico.

Paratypes: 110. Tropical Western Atlantic: 37 localities.

TRACHICHTHYIDAE

Hybognathus hankinsoni Hubbs, 1929

In: Jordan, Manual of Vertebrate Animals of Northeast United States, p. 88.

Paratypes: 13. Iowa: 4 localities.

CYPRINIDAE

Hybopsis x-punctata trautmani Hubbs and Crowe, 1956

Occas. Pap. Mus. Zool. Univ. Mich., 578: 7.

Paratype. Indiana: Park County, Sugar Creek.

CYPRINIDAE

Hybopsis x-punctata x-punctata Hubbs and Crowe, 1956

Occas. Pap. Mus. Zool. Univ. Mich., 578: 7.

Paratypes: 2. Missouri: Arlington.

Two additional specimens with localities "Illinois" and "no data" were also labeled paratypes of *H. x-punctata* by Hubbs, but the subspecies designations were not given.

CYPRINIDAE

Hymenocephalus aterrimus Gilbert, 1905

Bull. U.S. Fish Comm. 1903, 23(2): 666.

Paratype. Hawaiian Islands.

MACROURIDAE

Hymenocephalus grimaldii Weber, 1913

Die Fische der Siboga-Expedition, Leiden, p. 169.

Syntype: FMNH 52450 (CM 708). Indonesia: Bali Sea.

MACROURIDAE

***Hypessobrycon bifasciatus* Ellis, 1911**

Ann. Carnegie Mus., 8(2): 156.

Holotype: FMNH 54404 (CM 3026). Brazil: Rio Parahyba at Campos.

Paratypes: 201. Brazil: 8 localities.

CHARACIDAE

***Hypessobrycon duragenys* Ellis, 1911**

Ann. Carnegie Mus., 8(2): 155.

Holotype: (CM 3023) MISSING. Brazil: Rio Tietê at Mogy das Cruces.

Paratypes: 14. Brazil: 2 localities.

CHARACIDAE

***Hypessobrycon ecuadoriensis* Eigenmann and Henn, 1914**

In: Eigenmann, Henn, and Wilson, Indiana Univ. Stud., 19: 9.

Paratypes: 107. Ecuador: forest pool near Vines.

CHARACIDAE

***Hypessobrycon eos* Durbin, 1909**

Ann. Carnegie Mus., 6(1): 69.

Holotype: FMNH 52795 (CM 1194). Guyana: creek between Potaro Landing and Kanagaruma.

Paratypes: 23. Guyana: 2 localities.

CHARACIDAE

***Hypessobrycon melanopleurus* Ellis, 1911**

Ann. Carnegie Mus., 8(2): 157.

Holotype: FMNH 54413 (CM 3035). Brazil: São Paulo, Alto da Serra.

Paratypes: 2. Same locality.

CHARACIDAE

***Hypessobrycon minimus* Durbin, 1909**

Ann. Carnegie Mus., 6(1): 68.

Holotype: FMNH 52794 (CM 1193). Guyana: Cane Grove Corner.

CHARACIDAE

***Hypessobrycon minor* Durbin, 1909**

Ann. Carnegie Mus., 6(1): 65.

Holotype: FMNH 52789 (CM 1189) MISSING. Guyana: Konawaruk.

CHARACIDAE

***Hypessobrycon panamensis daguae* Eigenmann, 1922**

Mem. Carnegie Mus., 9(1): 141.

Syntypes: 9, FMNH 56526–56529 and 58316. Colombia: Río Patía basin.

CHARACIDAE

***Hypessobrycon parvulus* Ellis, 1911**

Ann. Carnegie Mus., 8(2): 153.

Holotype: FMNH 54392 (CM 3011). Brazil: Rio Catu at Alagoinhas.

Paratypes: 13. Brazil: 4 localities.

CHARACIDAE

***Hypessobrycon poecilioides* Eigenmann, 1913**

Indiana Univ. Stud., 18: 29.

Holotype: FMNH 56290 (CM 5091). Colombia: Cali.

Paratypes: 36. Same locality.

CHARACIDAE

***Hypessobrycon proteus* Eigenmann, 1913**

Indiana Univ. Stud., 18: 28.

Holotype: FMNH 56293 (CM 5094). Colombia: Quibdó.

Paratypes: 198. Colombia: 8 localities.

CHARACIDAE

***Hypessobrycon reticulatus* Ellis, 1911**

Ann. Carnegie Mus., 8(2): 153.

Holotype: FMNH 54397 (CM 3018). Brazil: Rio Parahyba at Campos.

Paratypes: 17. Brazil: 6 localities.

CHARACIDAE

***Hypessobrycon rosaceus* Durbin, 1909**

Ann. Carnegie Mus., 6(1): 67.

Holotype: FMNH 52791 (CM 1190). Guyana: Gluck Island.

Paratypes: 10. Guyana: 2 localities.

CHARACIDAE

***Hypessobrycon stictus* Durbin, 1909**

Ann. Carnegie Mus., 6(1): 71.

Holotype: FMNH 52798 (CM 1197). Guyana: Lama Stop-off.

Paratypes: 146. Guyana: 5 localities.

CHARACIDAE

***Hypessobrycon taurocephalus* Ellis, 1911**

Ann. Carnegie Mus., 8(2): 151.

Holotype: FMNH 54389 (CM 3007). Brazil: Rio Iguaçu at Serrinha Paraná.

Paratypes: 59. Brazil: 2 localities.

CHARACIDAE

***Hypoatherina barnesi* Schultz, 1953**

Bull. U.S. Natl. Mus., 202(1): 304.

Paratypes: 4. Marshall Islands: 2 localities.

ATHERINIDAE

Hyporhamphus hildebrandi Jordan and Evermann, 1927

Proc. Calif. Acad. Sci., 16(15): 503.

Lectotype: FMNH 8260. Panama: Colón, Fox Bay at Toro Point.

Lectoparatypes: 6. Same locality.

Lectotype designated by Weed (1933, Field Mus. Nat. Hist. Zool. Ser., 20: 60).

HEMIRAMPHIDAE

Hyporhamphus melanopterus Collette and Parin, 1978

Proc. Biol. Soc. Wash., 91(3): 738.

Paratypes: 2. Malaysia: Sabah, Sandakan, fish market.

HEMIRAMPHIDAE

Hyporhamphus snyderi Meek and Hildebrand, 1923

Field Mus. Nat. Hist. Zool. Ser., 15(1): 240.

Paratypes: 2. Panama: Chame Point.

HEMIRAMPHIDAE

Hypsoblennius lignus Meek and Hildebrand, 1928

Field Mus. Nat. Hist. Zool. Ser., 15(3): 946.

Paratype. Panama: Chame Point.

BLENNIIDAE

Hysteronotus megalostomus Eigenmann, 1911

Ann. Carnegie Mus., 8(1): 171.

Holotype: FMNH 54889 (CM 3551). Brazil: Rio das Velhas.

Paratypes: 2. Same locality.

CHARACIDAE

Iburiella kasawae Jordan and Hubbs, 1925

Mem. Carnegie Mus., 10(2): 291.

Holotype: FMNH 58796 (CM 7906). Japan: Tomakomai, near Mororan.

AGONIDAE

Ichthyomyzon unicuspis Hubbs and Trautman, 1937

Misc. Publ. Mus. Zool. Univ. Mich., 35: 53.

Paratypes: 8. Illinois and Ohio: 3 localities.

PETROMYZONTIDAE

Icticus ischanus Jordan and Thompson, 1914

Mem. Carnegie Mus., 6(4): 242.

Holotype: FMNH 57097 (CM 6036). Japan: Ryukyu Islands, Okinawa.

NOMEIDAE

Ilisha argentata Meek and Hildebrand, 1923

Field Mus. Nat. Hist. Zool. Ser., 15(1): 190.

Paratypes: 15. Panama: 5 localities.

CLUPEIDAE

Imparfinis hollandi Haseman, 1911

Ann. Carnegie Mus., 7(3–4): 383.

Holotype: FMNH 54244 (CM 2864). Brazil: Rio Iguaçu at Pôrto União da Victoria.

Haseman stated “Type unique”, but lot CM 2864 included 3 specimens. The 2 topotypes are now FMNH 54245.

PIMELODIDAE

Imparfinis microps Eigenmann and Fisher, 1916

In: Eigenmann, Ann. Carnegie Mus., 10(1–2): 82.

Holotype: FMNH 57793 (CM 6776). Colombia: Río Negro at Villavicencio.

Paratypes (?): 5. Same locality.

Published number CM 8778 is an error. Original description was based on “type only”; but 5 paratypes were listed in Eigenmann (1922, Mem. Carnegie Mus., 9(1): 221).

PIMELODIDAE

Imparfinis mirini Haseman, 1911

Ann. Carnegie Mus., 7(3–4): 318.

Holotype: FMNH 54335 (CM 2981). Brazil: Rio Piracicaba-mirini near Piracicaba.

Paratypes: 6. Same locality.

PIMELODIDAE

Iridio bimaculata Wilson, 1916

Ann. Carnegie Mus., 10(1–2): 68.

Holotype: FMNH 56466 (CM 5280). Colombia: Río Dagua at Buenaventura.

Paratype. Same locality.

LABRIDAE

Iridio decoratus Bean, 1906

Proc. Biol. Soc. Wash., 19: 29.

Lectotype: FMNH 5128. Bermuda: Nonsuch Island.

Lectoparatype. Same locality.

Lectotype designated by Grey (1947, p. 160).

LABRIDAE

Iridio elegans Bean, 1906

Proc. Biol. Soc. Wash., 19: 30.

Lectotype: FMNH 5129. Bermuda: Nonsuch Island, Castle Harbor.

Lectoparatypes: 3. Bermuda: Cooper’s Island, Jones Bay.

Lectotype designated by Grey (1947, p. 160).

LABRIDAE

- Iridio meyeri** Bean, 1906
Proc. Biol. Soc. Wash., 19: 29.
Holotype: FMNH 4938. Bermuda: Nonsuch Island.
Published number FMNH 5496 is an error.
LABRIDAE
- Iridio microstomus** Bean, 1906
Proc. Biol. Soc. Wash., 19: 30.
Holotype: FMNH 5138. Bermuda: Nonsuch Island.
LABRIDAE
- Ischnomembras gabunensis** Fowler, 1903
Proc. Acad. Nat. Sci. Phila., 55: 731.
Paratype. Gabon: Gabon River.
Locality probably wrong since this is synonym of *Menidia beryllina*, a New World taxon; see remarks in Böhlke (1984, Spec. Publ. Acad. Nat. Sci. Phila., 14: 23).
ATHERINIDAE
- Jenkinsia majua majua** Whitehead, 1963
Bull. Brit. Mus. (Nat. Hist.) Zool., 10(6): 356.
Paratypes: 54. Gulf of Mexico: Campeche Bank.
CLUPEIDAE
- Jenkinsia majua woodsi** Whitehead, 1963
Bull. Brit. Mus. (Nat. Hist.) Zool., 10(6): 356.
Paratypes: 10. Caribbean Sea: off Quita Sueño Bank.
CLUPEIDAE
- Joturus daguae** Eigenmann, 1917
Proc. Am. Philos. Soc., 56(7): 681.
Holotype: FMNH 58457 (CM 7458). Colombia: Río Dagua at Caldas.
MUGILIDAE
- Kaupichthys atlanticus** Böhlke, 1956
Proc. Acad. Nat. Sci. Phila., 108: 66.
Paratype. Bermuda: Castle Harbor.
XENOCONGRIDAE
- Kaupichthys brachychirus** Schultz, 1953
Bull. U.S. Natl. Mus., 202(1): 67.
Paratype. Marshall Islands: Bikini Atoll, Roromuk Island.
XENOCONGRIDAE
- Kryptopterus parvanalis** Inger and Chin, 1959
Fieldiana Zool., 39(27): 284.
Holotype: FMNH 68014. Malaysia: Sabah, Kinabatangan River at Deramakot.
Paratypes: 25. Malaysia: Sabah, 14 localities.
SILURIDAE
- Kuhlia bilunulata** Herre, 1935
Field Mus. Nat. Hist. Zool. Ser., 18(12): 404.
Holotype: FMNH 17313. Fiji Islands: Viti Levu Island, Suva Harbor.
Paratypes: 9. Same locality.
KUHLIIDAE
- Labeo fisheri** Jordan and Starks, 1917
Ann. Carnegie Mus., 11: 436.
Holotype: FMNH 58911 (CM 8044). Sri Lanka: Mahaweli River at Kandy.
CYPRINIDAE
- Labeo jordani** Oshima, 1919
Ann. Carnegie Mus., 12(2-4): 204.
Holotype: FMNH 59089 (CM 8226). Taiwan: hatchery at Shori.
CYPRINIDAE
- Labrisomus lentiginosus** Bean, 1906
Proc. Biol. Soc. Wash., 19: 30.
Holotype: FMNH 5142. Bermuda: Paget Parish, tide pool at Doe's Rock.
LABRISOMIDAE
- Labrisomus multiporosus** Hubbs, 1953
Zoologica (N.Y.), 38(9): 131.
Paratypes: 2. California: 2 localities.
LABRISOMIDAE
- Lampanyctus jordani** Gilbert, 1913
Mem. Carnegie Mus., 6(2): 104.
Holotype: FMNH 55837 (CM 4617). Japan: Hokkaido, Nemuro.
MYCTOPHIDAE
- Lampanyctus micropunctatus** Chapman, 1939
Proc. U.S. Natl. Mus., 86(3062): 527.
Paratype. North Pacific: off Washington.
MYCTOPHIDAE
- Lamprologus obliquus** Nichols and Griscom, 1917
Bull. Am. Mus. Nat. Hist., 37(25): 731.
Paratypes: 3. Zaire: Stanleyville.
CICHLIDAE
- Landonia latidens** Eigenmann and Henn, 1914
In: Eigenmann, Henn, and Wilson, Indiana Univ. Stud., 19: 2.
Paratypes: 2. Ecuador: Vinces.
CHARACIDAE

Lasiancistrus brevispinis Heitmans, Nijssen, and Isbrucker, 1983
Bijdr. Dierkd., 53(1): 38.
Paratypes: 2. Surinam: Fallawatra River at Stondansie Fall.
LORICARIIDAE

Lasiancistrus caucanus Eigenmann, 1912
Indiana Univ. Stud., 16: 11.
Holotype: FMNH 56034 (CM 4824). Colombia: Cartago.
Paratypes: 4. Same locality.
LORICARIIDAE

Leiocassis robustus Inger and Chin, 1959
Fieldiana Zool., 39(27): 290.
Holotype: FMNH 68001. Malaysia: Sabah, Kinabatangan River at Deramakot.
Paratypes: 5. Malaysia: Sabah, 2 localities.
BAGRIDAE

Leiogaster melanopus Weber, 1913
Die Fische der Siboga-Expedition, Leiden, p. 180.
Syntype: FMNH 52454 (CM 712). Indonesia: Ceram Sea.
BAGRIDAE

Lepidaplois strophodes Jordan and Evermann, 1903
Bull. U.S. Fish Comm. 1902, 22: 190.
Paratype. Hawaiian Islands: Honolulu.
Misspelled *Lepidoplois* in Jordan and Evermann (1903).
LABRIDAE

Lepidomeda mollispinis mollispinis Miller and Hubbs, 1960
Misc. Publ. Mus. Zool. Univ. Mich., 115: 18.
Paratypes: 12. Nevada: Clark County, Virgin River.
CYPRINIDAE

Lepomis occidentalis Meek, 1902
Field Mus. Nat. Hist. Zool. Ser., 3(6): 118.
Holotype: FMNH 3560. Mexico: Río Conchos at Jiménez.
Paratypes: 28. Mexico: 2 localities.
CENTRARCHIDAE

Lepophidium aporrhox Robins, 1961
Copeia, 1961(2): 218.
Paratypes: 4. Atlantic Ocean: off Guyana and Surinam.
OPHIDIIDAE

Lepophidium pheromystax Robins, 1960
Bull. Mar. Sci. Gulf Caribb., 10(1): 84.
Paratypes: 19. Western Atlantic Ocean: 6 localities.
OPHIDIIDAE

Lepophidium staurophor Robins, 1958
Bull. Mar. Sci. Gulf Caribb., 8(4): 360.
Holotype: FMNH 64383. Western Caribbean Sea.
Paratype. Caribbean Sea.
OPHIDIIDAE

Leporinodus retropinnis Eigenmann, 1922
Mem. Carnegie Mus., 9(1): 116.
Holotype: FMNH 55174 (CM 3875). Brazil: Piracicaba.
ANOSTOMIDAE

Leporinus alternus Eigenmann, 1912
Mem. Carnegie Mus., 5: 307.
Holotype: FMNH 53361 (CM 1827). Guyana: Tukeit.
Paratypes: 5. Guyana: 2 localities.
ANOSTOMIDAE

Leporinus arcus Eigenmann, 1912
Mem. Carnegie Mus., 5: 300.
Holotype: FMNH 53366 (CM 1832). Guyana: Tukeit.
Paratypes: 4. Guyana: 4 localities.
ANOSTOMIDAE

Leporinus brunneus Myers, 1950
Stanford Ichthyol. Bull., 3(4): 195.
Paratype. Venezuela: Casiquiare, Chemoni, near bifurcation of Río Orinoco.
ANOSTOMIDAE

Leporinus ecuadoriensis Eigenmann and Henn, 1916
Ann. Carnegie Mus., 10(1-2): 88.
Paratypes: 2. Ecuador: Río Barranca Alta near Naranjito.
ANOSTOMIDAE

Leporinus granti Eigenmann, 1912
Mem. Carnegie Mus., 5: 307.
Holotype: FMNH 53383 (CM 1851). Guyana: Maripicru Creek.
Paratypes: 2. Same locality.
ANOSTOMIDAE

Leporinus steyermarki Inger, 1956

Fieldiana Zool., 34(37): 431.

Holotype: FMNH 45701. Venezuela: Río Abácapa, west side of Chimantá-tepui.

ANOSTOMIDAE

Leptocephalus ectenurus Jordan and Richardson, 1909

Mem. Carnegie Mus., 4(4): 171.

Holotype: FMNH 52117 (CM 245). Taiwan: Takao.

CONGRIDAE

Leptocephalus harringtonensis Mowbray, 1931

Fauna Bermudensis, 1: 1.

Holotype: FMNH 48442. Bermuda: Harrington Sound.

CONGRIDAE

Leptocottus armatus australis Hubbs, 1921

Occas. Pap. Mus. Zool. Univ. Mich., 94: 5.

Paratypes: 62. California: 10 localities.

COTTIDAE

Leptodoras linnelli Eigenmann, 1912

Mem. Carnegie Mus., 5: 191.

Holotype: FMNH 53561 (CM 1626). Guyana: Tumatumari.

Paratypes: 16. Guyana: 4 localities.

DORADIDAE

Leptoglanis essequibensis Eigenmann, 1912

Mem. Carnegie Mus., 5: 158.

Holotype: FMNH 53331 (CM 1652). Guyana: Crab Falls.

Paratypes: 3. Guyana: 2 localities.

PIMELODIDAE

Leptophilypnus fluviatilis Meek and Hildebrand, 1916

Field Mus. Nat. Hist. Zool. Ser., 10(5): 361.

Holotype: FMNH 8952. Panama: Canal Zone, creek at Mindi.

Paratypes: 20. Same locality.

ELEOTRIDAE

Leucichthys alpenae Koelz, 1924

Occas. Pap. Mus. Zool. Univ. Mich., 146: 1.

Paratypes: 2. Michigan: 2 localities.

SALMONIDAE

Leucichthys kiyi Koelz, 1921

Occas. Pap. Mus. Zool. Univ. Mich., 104: 1.

Paratypes: 2. Michigan: Lake Michigan at Ludington.

SALMONIDAE

Leucichthys reighardi Koelz, 1924

Occas. Pap. Mus. Zool. Univ. Mich., 146: 5.

Paratypes: 2. Wisconsin and Indiana: 2 localities.

SALMONIDAE

Leuciscus carletoni Kendall, 1903

Bull. U.S. Fish Comm. 1902, 22: 357.

Paratypes: 9. Maine: East Branch, Penobscot River.

CYPRINIDAE

Leucogobio polytaenia Nichols, 1925

Am. Mus. Nov., 181: 6.

Paratypes: 3. China: Shansi, Niang-tze-kwan.

CYPRINIDAE

Limia hollandi Henn, 1916

Ann. Carnegie Mus., 10(1-2): 138.

Holotype: FMNH 55861 (CM 4643). Brazil: Rio São Francisco at Penedo.

Paratypes: 56. Brazil: 11 localities.

POECILIIDAE

Limia rivasi Franz and Burgess, 1983

Northeast Gulf Sci., 6(1): 51.

Paratypes: 10. Haiti: Ile de la Gonâve.

POECILIIDAE

Liobagrus anguillicauda Nichols, 1926

Am. Mus. Nov., 224: 1.

Paratypes: 2. China: Fukien, Chungan Hsien.

BAGRIDAE

Liobagrus nantoenis Oshima, 1919

Ann. Carnegie Mus., 12(2-4): 183.

Holotype: FMNH 59080 (CM 8217). Taiwan: Dainansho, Nanto.

BAGRIDAE

Liopempheris sasakii Jordan and Hubbs, 1925

Mem. Carnegie Mus., 10(2): 228.

Holotype: FMNH 58751 (CM 7860). Japan: Toba market.

Published number CM 7861 is an error.

PEMPHERIDAE

Liopropoma mowbrayi Woods and Kanazawa, 1951

Fieldiana Zool., 31(53): 633.

Holotype: FMNH 48544. Bermuda: south shore.

SERRANIDAE

Lithogenes villosus Eigenmann, 1909
Ann. Carnegie Mus., 6(1): 6.
Holotype: FMNH 52960 (CM 1002). Guyana:
Potaro River at Aruataima Falls.
LORICARIIDAE

Lithoxus lithoides Eigenmann, 1912
Mem. Carnegie Mus., 5: 242.
Holotype: FMNH 53557 (CM 1527). Guyana:
Amatuk.
Paratypes: 122. Guyana: 6 localities.
LORICARIIDAE

Liza formosae Oshima, 1922
Ann. Carnegie Mus., 13(3–4): 251.
Holotype: FMNH 59145 (CM 8283). Taiwan:
Anpin.
MUGILIDAE

Liza parva Oshima, 1922
Ann. Carnegie Mus., 13(3–4): 253.
Holotype: FMNH 59146 (CM 8284). Taiwan:
Anpin.
MUGILIDAE

Liza pescadorensis Oshima, 1922
Ann. Carnegie Mus., 13(3–4): 254.
Holotype: FMNH 59147 (CM 8285). Taiwan:
Pescadores Island at Bako.
MUGILIDAE

Lonchopisthus higmani Mead, 1959
Stud. Fauna Suriname Other Guyanas, 2(5):
104.
Paratype. Western Atlantic Ocean: off Surinam.
OPISTOGNATHIDAE

Lophiobagrus aquilus Bailey and Stewart, 1984
Misc. Publ. Mus. Zool. Univ. Mich., 168: 24.
Paratypes: 5. Zambia: Lake Tanganyika.
BAGRIDAE

Lophiobagrus brevispinis Bailey and Stewart, 1984
Misc. Publ. Mus. Zool. Univ. Mich., 168: 35.
Paratypes: 4. Zambia: Lake Tanganyika.
BAGRIDAE

Lophiodes beroe Caruso, 1981
Copeia, 1981(3): 542.
Paratypes: 2. Western North Atlantic: 2 locali-
ties.
LOPHIIDAE

Lophiodes reticulatus Caruso and Suttkus, 1979
Bull. Mar. Sci., 29(4): 491.
Paratypes: 7. Gulf of Mexico.
LOPHIIDAE

Loricaria capetensis Meek and Hildebrand, 1913
Field Mus. Nat. Hist. Zool. Ser., 10(8): 80.
Holotype: FMNH 7582. Panama: Río Capetí.
LORICARIIDAE

Loricaria filamentosa latiura Eigenmann and
Vance, 1912
In: Eigenmann, Indiana Univ. Stud., 16: 13.
Syntypes: 7, FMNH 55115 (CM 3806). Colom-
bia: Boca de Certegui.
LORICARIIDAE

Loricaria filamentosa seminuda Eigenmann and
Vance, 1912
In: Eigenmann, Indiana Univ. Stud., 16: 13.
Holotype: FMNH 55116 (CM 3807). Colombia:
Girardot.
LORICARIIDAE

Loricaria fimbriata Eigenmann and Vance, 1912
In: Eigenmann, Indiana Univ. Stud., 16: 12.
Holotype: FMNH 55117 (CM 3808). Colombia:
Boca de Certegui.
Paratype. Same locality.
LORICARIIDAE

Loricaria griseus Eigenmann, 1909
Ann. Carnegie Mus., 6(1): 8.
Holotype: FMNH 53077 (CM 1504). Guyana:
Konawaruk.
Paratypes: 13. Guyana: 2 localities.
LORICARIIDAE

Loricaria gymnogaster Eigenmann and Vance,
1912
In: Eigenmann, Indiana Univ. Stud., 16: 12.
Paratypes: 29. Colombia: Girardot.
LORICARIIDAE

Loricaria gymnogaster lagoichthys Schultz, 1944
Proc. U.S. Natl. Mus., 94(3172): 331.
Paratypes: 5. Venezuela: Río Motatán north of
Motatán.
LORICARIIDAE

Loricaria microdon Eigenmann, 1909
Ann. Carnegie Mus., 6(1): 7.
Holotype: FMNH 53555 (CM 1507). Guyana:
Rupununi.
LORICARIIDAE

- Loricaria stewarti** Eigenmann, 1909
Ann. Carnegie Mus., 6(1): 9.
Holotype: FMNH 53330 (CM 1508). Guyana:
Ireng River system, Chipoo Creek.
Paratypes: 4. Same locality.
LORICARIIDAE
- Loricaria submarginatus** Eigenmann, 1909
Ann. Carnegie Mus., 6(1): 10.
Holotype: FMNH 53080 (CM 1510). Guyana:
creek below Potaro Landing.
LORICARIIDAE
- Loricaria tuyrensis** Meek and Hildebrand, 1913
Field Mus. Nat. Hist. Zool. Ser., 10(8): 81.
Holotype: FMNH 7583. Panama: Río Tuyra,
Boca de Cupe.
Paratypes: 7. Same locality.
LORICARIIDAE
- Loricaria uracantha rupestre** Schultz, 1944
Proc. U.S. Natl. Mus., 94(3172): 325.
Paratypes: 5. Venezuela: Río San Juan south of
Mene Grande.
LORICARIIDAE
- Loricaria variegata venezuelae** Schultz, 1944
Proc. U.S. Natl. Mus., 94(3172): 329.
Paratypes: 5. Venezuela: Río Sucuy, above its
mouth.
LORICARIIDAE
- Luciobrotula corethromycter** Cohen, 1954
Bull. Mar. Sci. Gulf Caribb., 14(3): 391.
Paratype. Gulf of Mexico.
OPHIDIIDAE
- Lumpenus longirostris** Evermann and Goldsborough, 1907
Bull. Bur. Fish. 1906, 26: 340.
Paratypes: 2. Alaska: Albatross Station 4252.
STICHAEIDAE
- Lycodes tanakae** Jordan and Thompson, 1914
Mem. Carnegie Mus., 6(4): 299.
Holotype: FMNH 57068 (CM 6004). Japan:
Hondo, Noto.
ZOARCIDAE
- Macgregorella badia** Herre, 1935
Field Mus. Nat. Hist. Zool. Ser., 18(12): 415.
Holotype: FMNH 17373. Fiji Islands: Ovalau
Island.
GOBIIDAE
- Macgregorella santa** Herre, 1935
Field Mus. Nat. Hist. Zool. Ser., 18(12): 416.
Holotype: FMNH 17374. New Hebrides Is-
lands: Espiritu Santo Island, Hog Harbor.
Paratypes: 7. New Hebrides Islands: 2 localities.
GOBIIDAE
- Macrodonogobius wilburi** Herre, 1936
Philipp. J. Sci., 59: 279.
Paratype (?). Palau Islands.
GOBIIDAE
- Macropodus filamentosus** Oshima, 1919
Ann. Carnegie Mus., 12(2-4): 278.
Holotype: FMNH 59123 (CM 8261). Taiwan:
Kotosho (Botel Tobago Island).
BELONTIIDAE
- Macropsobrycon uruguayanae** Eigenmann, 1915
Mem. Carnegie Mus., 7(1): 48.
Holotype: FMNH 57910 (CM 6895). Brazil:
Cacequy.
Paratypes: 5. Brazil: 2 localities.
CHARACIDAE
- Macrostoma quercinum japonicum** Tanaka, 1908
J. Coll. Sci. Imp. Univ. Tokyo, 23(13): 5.
Holotype: FMNH 80459. Japan: Sagami Sea.
MYCTOPHIDAE
- Macrurus richardi** Weber, 1913
Die Fische der Siboga-Expedition, Leiden, p.
154.
Syntype: FMNH 52442 (CM 700). Indonesia:
Flores Sea.
MACROURIDAE
- Malacoctenus afueræ multipunctatus** Springer,
1958
Inst. Mar. Sci., 5: 462.
Paratypes: 52. Mexico: Golfo de California, Ba-
hía Santa Inez.
LABRISOMIDAE
- Malacoctenus ebisui** Springer, 1958
Inst. Mar. Sci., 5: 478.
Paratype. Costa Rica: Golfo de Nicoya, Isla Tor-
tuga.
LABRISOMIDAE
- Malacoctenus gigas** Springer, 1958
Inst. Mar. Sci., 5: 457.
Paratypes: 48. Mexico: Golfo de California, Isla
Tiburón.
LABRISOMIDAE

- Malacoctenus hubbsi hubbsi** Springer, 1958
Inst. Mar. Sci., 5: 465.
Paratypes: 125. Mexico: Guaymas, 2 localities.
LABRISOMIDAE
- Malacoctenus hubbsi polyporosus** Springer, 1958
Inst. Mar. Sci., 5: 468.
Paratypes: 32. Mexico: Golfo de California, Isla Pájaro.
LABRISOMIDAE
- Malacoctenus margaritae mexicanus** Springer, 1958
Inst. Mar. Sci., 5: 449.
Paratypes: 47. Mexico: Golfo de California, Bahía Santa Inez.
LABRISOMIDAE
- Malacoctenus puertoricensis** Evermann and Marsh, 1900
Rept. U.S. Fish Comm. 1899, 25: 358.
Paratypes: 2. Puerto Rico: Culebra.
LABRISOMIDAE
- Malacoctenus triangulatus** Springer, 1958
Inst. Mar. Sci., 5: 477.
Paratypes: 8. Gulf of Mexico: 2 localities.
LABRISOMIDAE
- Malacoctenus zacae** Springer, 1958
Inst. Mar. Sci., 5: 470.
Paratypes: 12. Mexico: Golfo de California, Bahía San Lucas.
LABRISOMIDAE
- Malacichthys wakiyae** Jordan and Hubbs, 1925
Mem. Carnegie Mus., 10(2): 233.
Holotype: FMNH 58754 (CM 7863). Japan: Kagoshima Bay.
PERCICHTHYIDAE
- Mastacembelus keithi** Herre, 1940
Bull. Raffles Mus., 16: 24.
Paratype. Malaysia: Sabah, Kabili River.
MASTACEMBELIDAE
- Mastacembelus oatesii** Boulenger, 1893
Ann. Mag. Nat. Hist., Ser. 6, 12: 199.
Syntype: FMNH 5707. Burma: Fort Stedman.
MASTACEMBELIDAE
- Megalamphodus megalopterus** Eigenmann, 1915
Mem. Carnegie Mus., 7(1): 50.
Holotype: FMNH 57823 (CM 6806). Brazil: Cáceres.
Paratypes: 2. Same locality.
CHARACIDAE
- Megalamphodus micropterus** Eigenmann, 1915
Mem. Carnegie Mus., 7(1): 54.
Holotype: FMNH 57916 (CM 6900). Brazil: Rio São Francisco at Lagoa do Porto.
Paratypes: 54. Brazil: 4 localities.
CHARACIDAE
- Megalonema platycephalum** Eigenmann, 1912
Mem. Carnegie Mus., 5: 150.
Holotype: FMNH 53224 (CM 1684). Guyana: Tumatumari.
Paratypes: 2. Guyana: Crab Falls.
PIMELODIDAE
- Megalonema punctatum** Meek and Hildebrand, 1913
Field Mus. Nat. Hist. Zool. Ser., 10(8): 77.
Holotype: FMNH 7577. Panama: Río Tuyra, Boca de Cupe.
PIMELODIDAE
- Megalonema robustum** Meek and Hildebrand, 1913
Field Mus. Nat. Hist. Zool. Ser., 10(8): 78.
Holotype: FMNH 7578. Panama: Río Tuyra, Marriganti.
Paratypes: 3. Panama: 2 localities.
PIMELODIDAE
- Megalonema xanthum** Eigenmann, 1912
Indiana Univ. Stud., 16: 16.
Holotype: FMNH 56032 (CM 4822). Colombia: Girardot.
Paratypes: 90. Same locality.
PIMELODIDAE
- Melamphaes pumilus** Ebeling, 1962
Dana-Rep. Carlsberg Found. 58: 106.
Paratypes: 31. Western North Atlantic: 6 localities.
MELAMPHAIDAE
- Melaniris balsanus** Meek, 1902
Field Mus. Nat. Hist. Zool. Ser., 3(6): 117.
Holotype: FMNH 3706. Mexico: Río Balsas at Balsas.
Paratypes: 99. Same locality.
ATHERINIDAE

- Melaniris sardina** Meek, 1907
Field Mus. Nat. Hist. Zool. Ser., 7(4): 114.
Holotype: FMNH 5937. Nicaragua: Lago Managua at Managua.
Paratypes: 309. Nicaragua: 3 localities.
ATHERINIDAE
- Melanotaenia kabia** Herre, 1935
Field Mus. Nat. Hist. Zool. Ser., 18(12): 397.
Holotype: FMNH 17230. New Guinea: Sepik River at Nyaurangai.
Paratypes: 14. New Guinea: 2 localities.
MELANOTAENIIDAE
- Melanotaenia rosacea** Herre, 1935
Field Mus. Nat. Hist. Zool. Ser., 18(12): 398.
Holotype: FMNH 17245. New Guinea: Sepik River at Koragu.
Paratypes: 19. New Guinea: 2 localities.
MELANOTAENIIDAE
- Menidia chagresi** Meek and Hildebrand, 1914
Field Mus. Nat. Hist. Zool. Ser., 10(10): 119.
Paratypes: 59. Panama and Costa Rica: 3 localities.
ATHERINIDAE
- Menidia lisa** Meek, 1904
Field Mus. Nat. Hist. Zool. Ser., 5: 182.
Holotype: FMNH 4633. Mexico: Veracruz, Refugio.
Paratypes: 48. Mexico: 2 localities.
ATHERINIDAE
- Menidia peninsulae atrimentis** Kendall, 1902
Rept. U.S. Fish Comm. 1901, 27: 258.
Paratypes: 5. Florida: Lake Monroe.
ATHERINIDAE
- Menidia starksi** Meek and Hildebrand, 1923
Field Mus. Nat. Hist. Zool. Ser., 15(1): 267.
Paratypes: 4. Panama: Taboga Island.
ATHERINIDAE
- Menticirrhus focaliger** Ginsburg, 1952
J. Wash. Acad. Sci., 42(3): 97.
Paratypes: 2. Florida: Cape San Blas.
SCIAENIDAE
- Metynnis guaporensis** Eigenmann, 1915
Ann. Carnegie Mus., 9(3-4): 267.
Holotype: FMNH 56913 (CM 5729). Brazil: Rio Guaporé at Maciel.
Paratypes: 2. Brazil and Bolivia: 2 localities.
CHARACIDAE
- Metynnis mola** Eigenmann and Kennedy, 1903
Proc. Acad. Nat. Sci. Phila., 59: 528.
Paratype. Paraguay: Arroyo Trementina.
CHARACIDAE
- Metynnis roosevelti** Eigenmann, 1915
Ann. Carnegie Mus., 9(3-4): 268.
Holotype: FMNH 56925 (CM 5740). Brazil: Manaus.
Paratypes: 8. Brazil: 3 localities.
CHARACIDAE
- Microbrycon minutus** Eigenmann and Wilson, 1914
In: Eigenmann, Henn, and Wilson, Indiana Univ. Stud., 19: 3.
Holotype: FMNH 56606 (CM 5422). Colombia: Río Truando.
CHARACIDAE
- Microdesmus carri** Gilbert, 1966
Copeia, 1966(2): 326.
Paratypes: 2. Costa Rica: Limón Province near mouth of Tortuguero.
MICRODESMIDAE
- Microdesmus woodsi** Kanazawa, 1952
Fieldiana Zool., 34(7): 93.
Holotype: FMNH 48876. Bermuda: Saint David's Island.
MICRODESMIDAE
- Microeleotris mindii** Meek and Hildebrand, 1916
Field Mus. Nat. Hist. Zool. Ser., 10(15): 364.
Holotype: FMNH 8954. Panama: Canal Zone, creek at Mindi.
Paratypes: 25. Panama: 2 localities.
ELEOTRIDAE
- Microeleotris panamensis** Meek and Hildebrand, 1916
Field Mus. Nat. Hist. Zool. Ser., 10(15): 363.
Holotype: FMNH 8953. Panama: Río Juan Díaz.
Paratypes: 18. Panama: 2 localities.
ELEOTRIDAE
- Microgenys minutus** Eigenmann, 1913
Indiana Univ. Stud., 18: 22.
Holotype: FMNH 56215 (CM 5007). Colombia: Piedra de Moler.
Paratype. Same locality.
CHARACIDAE

Microglanis iheringi Gomes, 1946

Occas. Pap. Mus. Zool. Univ. Mich., 494: 9.
Holotype: FMNH 35350. Venezuela: Aragua,
Río Turmero.

Paratypes: 2. Same locality.

PIMELODIDAE

Microglanis poecilus Eigenmann, 1912

Mem. Carnegie Mus., 5: 155.

Holotype: FMNH 47365 (CM 1676). Guyana:
below Packeo Falls.

Paratype. Same locality.

PIMELODIDAE

Microglanis secundus Mees, 1974

Zool. Verh. (Leiden), 132: 235.

Paratypes: 3. Surinam: Coppename River.

PIMELODIDAE

Microglanis variegatus Eigenmann, 1914

In: Eigenmann, Henn, and Wilson, Indiana
Univ. Stud., 19: 14.

Paratype. Ecuador: Vines.

PIMELODIDAE

Micrognathus suvensis Herre, 1935

Field Mus. Nat. Hist. Zool. Ser., 18(12): 396.

Holotype: FMNH 17229. Fiji Islands: Viti Levu
Island, Suva Harbor.

SYNGNATHIDAE

Micropterus pseudaplites Hubbs, 1927

Occas. Pap. Mus. Zool. Univ. Mich., 184: 8.

Paratypes: 13. Arkansas: 6 localities.

CENTRARCHIDAE

Microschemobrycon guaporensis Eigenmann, 1915

Mem. Carnegie Mus., 7(1): 56.

Holotype: FMNH 57926 (CM 6910). Brazil: Rio
Guaporé at Maciel.

CHARACIDAE

Microspathodon frontatus Emery, 1970

Stud. Trop. Oceanogr. Inst. Mar. Sci. Miami,
4(2): 294.

Paratypes: 35. Gulf of Guinea: 3 localities.

POMACENTRIDAE

Mionorus pacificus Herre, 1935

Field Mus. Nat. Hist. Zool. Ser., 18(12): 403.

Holotype: FMNH 17311. Ecuador: Islas Galá-
pagos, Isla Albemarle, Tagus Cove.

Paratype. Same locality.

APOGONIDAE

Misgurnus anguillicaudatus tungting Nichols, 1925

Am. Mus. Nov., 169: 3.

Paratypes: 2. China: Hunan, Tungting Lake.

COBITIDAE

Misgurnus mizolepis fukien Nichols, 1925

Am. Mus. Nov., 169: 4.

Paratype. China: Fukien.

COBITIDAE

Misgurnus mizolepis grangeri Nichols, 1925

Am. Mus. Nov., 169: 5.

Paratypes: 2. China: Szechwan.

COBITIDAE

Misgurnus mohoity yunnan Nichols, 1925

Am. Mus. Nov., 169: 5.

Paratype. China: Yunnan.

COBITIDAE

Mnierpes macrocephalus catherinae Hubbs, 1952

Stanford Ichthyol. Bull., 4(2): 61.

Paratypes: 3. Nicaragua and Costa Rica: 2 lo-
calities.

LABRISOMIDAE

Moenkhausia browni Eigenmann, 1909

Ann. Carnegie Mus., 6(1): 13.

Holotype: FMNH 52733 (CM 1106). Guyana:
Potaro River at Aruataima Falls.

Paratypes: 28. Guyana: 7 localities.

CM 1106 was received from Carnegie Mu-
seum with 2 specimens and 2 labels, one
of each reads "1106 type (a)". FMNH
53732 (64 mm total length) is most likely
the holotype, but we defer a final deter-
mination to a revisor.

CHARACIDAE

Moenkhausia lepidura hasemani Eigenmann, 1917

Mem. Carnegie Mus., 43(1): 102.

Holotype: FMNH 55055 (CM 3746). Brazil:
Santarém.

CHARACIDAE

Moenkhausia metae Eigenmann, 1922

Mem. Carnegie Mus., 9(1): 234.

Paratypes: 4. Colombia: Villavicencio.

CHARACIDAE

Moenkhausia phaeonota Fink, 1979

Breviora, 450: 2.

Paratype. Brazil: Mato Grosso, Rio Arinos.

CHARACIDAE

- Moenkhausia profunda** Eigenmann, 1912
 Mem. Carnegie Mus., 5: 322.
 Holotype: FMNH 53717 (CM 2207). Guyana:
 Issorora Rubber Plantation.
 CHARACIDAE
- Moenkhausia shideleri** Eigenmann, 1909
 Ann. Carnegie Mus., 6(1): 15.
 Holotype: FMNH 52961 (CM 1012). Guyana:
 Bartica.
 CHARACIDAE
- Monacanthus tuckeri** Bean, 1906
 Proc. Biol. Soc. Wash., 19: 33.
 Lectotype: FMNH 5183. Bermuda: The Flatts.
 Lectoparatypes: 2. Bermuda: 2 localities.
 Lectotype designated by Grey (1947, p. 168).
 BALISTIDAE
- Monolene megalepis** Woods, 1961
 Copeia, 1961(2): 192.
 Holotype: FMNH 65746. Puerto Rico: west of
 Arecibo.
 Paratypes: 9. Caribbean Sea: 4 localities.
 BOTHIDAE
- Monomitopus kumae** Jordan and Hubbs, 1925
 Mem. Carnegie Mus., 10(2): 324.
 Holotype: FMNH 58844 (CM 7954). Japan:
 Misaki.
 OPHIDIIDAE
- Moroco yamamotis** Jordan and Hubbs, 1925
 Mem. Carnegie Mus., 10(2): 182.
 Holotype: FMNH 58723 (CM 7829). Korea:
 Lake Yamanaka.
 CYPRINIDAE
- Moxostoma rubreques** Hubbs, 1930
 Misc. Publ. Mus. Zool. Univ. Mich., 20: 24.
 Paratypes: 2. Michigan: Raisin River at Dundee.
 CATOSTOMIDAE
- Mugil anpinensis** Oshima, 1922
 Ann. Carnegie Mus., 13(3-4): 245.
 Holotype: FMNH 59143 (CM 8281). Taiwan:
 Anpin.
 MUGILIDAE
- Mugilops cyanellus** Meek and Hildebrand, 1923
 Field Mus. Nat. Hist. Zool. Ser., 15(1): 271.
 Paratypes: 2. Panama: Canal Zone, 2 localities.
 ATHERINIDAE
- Mulloidies flammeus** Jordan and Evermann, 1903
 Bull. U.S. Fish Comm. 1902, 22: 186.
 Paratype: MISSING. Hawaiian Islands: Hilo.
 MULLIDAE
- Myctophum hollandi** Jordan and Jordan, 1922
 Mem. Carnegie Mus., 10(1): 11.
 Holotype: FMNH 55191 (CM 3897). Hawaiian
 Islands: Honolulu Market.
 MYCTOPHIDAE
- Mylossoma ocellatum** Eigenmann, 1915
 Ann. Carnegie Mus., 9(3-4): 265.
 Holotype: FMNH 56816 (CM 5629). Paraguay:
 Río Confuso at Villa Hays.
 Paratypes: 15. Same locality.
 CHARACIDAE
- Myoglanis potaroensis** Eigenmann, 1912
 Mem. Carnegie Mus., 5: 159.
 Holotype: FMNH 53333 (CM 1664). Guyana:
 Tukeit.
 Paratypes: 14. Guyana: 5 localities.
 PIMELODIDAE
- Myripristis argyromus** Jordan and Evermann, 1903
 Bull. U.S. Fish Comm. 1902, 22: 172.
 Paratype. Hawaiian Islands: Honolulu.
 HOLOCENTRIDAE
- Myripristis berndti** Jordan and Evermann, 1903
 Bull. U.S. Fish Comm. 1902, 22: 170.
 Paratype. Hawaiian Islands: Honolulu.
 HOLOCENTRIDAE
- Myripristis bowditchae** Woods, 1953
 Bull. U.S. Natl. Mus., 202(1): 202.
 Paratypes: 4. Marshall Islands: 3 localities.
 HOLOCENTRIDAE
- Myripristis chryseres** Jordan and Evermann, 1903
 Bull. U.S. Fish Comm. 1902, 22: 171.
 Paratype. Hawaiian Islands: Hilo.
 HOLOCENTRIDAE
- Myripristis mooreanus** Herre, 1935
 Field Mus. Nat. Hist. Zool. Ser., 18(12): 394.
 Holotype: FMNH 17225. Society Islands: Moo-
 rea Island.
 Paratypes: 2. Same locality.
 HOLOCENTRIDAE

- Myripristis randalli** Greenfield, 1974
Nat. Hist. Mus. Los Ang. Cty. Sci. Bull., 19: 34.
Paratype. Austral Islands: Rurutu.
HOLOCENTRIDAE
- Myripristis symmetricus** Jordan and Evermann, 1903
Bull. U.S. Fish Comm. 1902, 22: 173.
Paratype. Hawaiian Islands: Honolulu.
HOLOCENTRIDAE
- Myripristis tiki** Greenfield, 1974
Nat. Hist. Mus. Los Ang. Cty. Sci. Bull., 19: 9.
Paratype. Chile: Easter Island, Hanga Pico.
HOLOCENTRIDAE
- Myripristis undecimalis** Herre, 1935
Field Mus. Nat. Hist. Zool. Ser., 18(12): 393.
Holotype: FMNH 17228. Tuamotu Islands: Takaroa Island.
HOLOCENTRIDAE
- Myripristis woodsi** Greenfield, 1974
Nat. Hist. Mus. Los Ang. Cty. Sci. Bull., 19: 32.
Paratypes: 3. Marshall Islands: Eniwetok Atoll, Girinien Island.
HOLOCENTRIDAE
- Mystus sabanus** Inger and Chin, 1959
Fieldiana Zool., 39(27): 294.
Holotype: FMNH 68088. Malaysia: Sabah, Kinabatangan River at Deramakot.
Paratypes: 13. Malaysia: Sabah, 3 localities.
BAGRIDAE
- Myxine biniplicata** Richardson and Jowett, 1951
Zool. Publ. Victoria Univ. Wellington, 12: 3.
Paratype. New Zealand: Cook Strait.
MYXINIDAE
- Myxus malayanus** Herre, 1936
Philipp. J. Sci., 59: 286.
Paratypes: 2. Sulawesi Island (Celebes): Lembeh Strait.
MUGILIDAE
- Nannacara bimaculata** Eigenmann, 1912
Mem. Carnegie Mus., 5: 487.
Holotype: FMNH 53799 (CM 2304). Guyana: Erukin.
CICHLIDAE
- Nannorhamdia nemacheir** Eigenmann and Fisher, 1916
In: Eigenmann, Ann. Carnegie Mus., 10(1-2): 83.
Holotype: FMNH 58127 (CM 7125). Colombia: Girardot.
Paratype. Same locality.
PIMELODIDAE
- Nannostomus marginatus** Eigenmann, 1909
Ann. Carnegie Mus., 6(1): 41.
Holotype: FMNH 53550 (CM 1171). Guyana: Maduni Creek.
Paratypes: 17. Guyana: 7 localities.
LEBIASINIDAE
- Nannostomus minimus** Eigenmann, 1909
Ann. Carnegie Mus., 6(1): 42.
Holotype: FMNH 52771 (CM 1165). Guyana: Erukin.
Paratype. Guyana: Amatuk.
LEBIASINIDAE
- Nannostomus simplex** Eigenmann, 1909
Ann. Carnegie Mus., 6(1): 42.
Holotype: FMNH 52773 (CM 1167). Guyana: Lama Stop-off.
LEBIASINIDAE
- Nanochromis transvestitus** Stewart and Roberts, 1984
Copeia, 1984(1): 82.
Holotype: FMNH 93584. Zaïre: Lac Mandombe near Ipeque.
Paratypes: 80. Same locality.
CICHLIDAE
- Nansenia ardesiaca** Jordan and Thompson, 1914
Mem. Carnegie Mus., 6(4): 210.
Holotype: FMNH 57086 (CM 6023). Japan: Sagami Sea, Okinose.
BATHYLAGIDAE
- Neetroplus carpintis** Jordan and Snyder, 1900
Bull. U.S. Fish Comm. 1899, 19: 146.
Paratype. Mexico: Laguna del Carpinte near Tampico.
CICHLIDAE
- Neetroplus fluviatilis** Meek, 1912
Field Mus. Nat. Hist. Zool. Ser., 10(7): 74.
Holotype: FMNH 7685. Costa Rica: Costa Rica River at Guapilis.
Paratype. Same locality.
CICHLIDAE

- Neetroplus panamensis** Meek and Hildebrand, 1913
Field Mus. Nat. Hist. Zool. Ser., 10(8): 90.
Holotype: FMNH 7601. Panama: Río Mandin-
go at Bas Obispo.
Paratypes: 20. Panama: 6 localities.
CICHLIDAE
- Nemaclinus atelestos** Böhlke and Springer, 1975
Proc. Acad. Nat. Sci. Phila., 127(7): 58.
Paratype. Caribbean Sea: off Virgin Islands.
LABRISOMIDAE
- Nematabramis alestes borneensis** Inger and Chin, 1962
Fieldiana Zool., 45: 47.
Holotype: FMNH 44791. Malaysia: Sabah, Kota
Belud.
Paratypes: 7. Same locality.
CYPRINIDAE
- Nematobrycon amphioxus** Eigenmann and Wil-
son, 1914
In: Eigenmann, Henn, and Wilson, Indiana
Univ. Stud., 19: 13.
Holotype: (CM 5050) MISSING. Colombia:
Boca de Raspadura.
Paratypes: 15. Colombia: 3 localities.
CHARACIDAE
- Nematobrycon lacortei** Weitzman and Fink, 1971
Beaufortia, 19(248): 59.
Paratype. Aquarium specimen, probably from
Colombia: Río Calima.
CHARACIDAE
- Neoharriotta carri** Bullis and Carpenter, 1966
Copeia, 1966(3): 443.
Paratypes: 2. Southern Caribbean Sea: 2 local-
ities.
RHINOCHIMAERIDAE
- Neoheterandria elegans** Henn, 1916
Ann. Carnegie Mus., 10(1-2): 118.
Holotype: FMNH 57007 (CM 5823). Colombia:
Truandó.
Paratypes: 7. Same locality.
POECILIIDAE
- Neostethus borneensis** Herre, 1939
Proc. Biol. Soc. Wash., 52: 143.
Paratypes: 12. Malaysia: Sabah, Kibili River.
NEOSTETHIDAE
- Nesiotes purpurascens** De Vis, 1884
Proc. Linn. Soc. N. S. Wales, 8(4): 453.
Neotype: FMNH 17340. New Hebrides Islands:
Efate Island, Vila.
Neotype designated by Herre (1936, Field
Mus. Nat. Hist. Zool. Ser., 21: 165).
PSEUDOCHROMIDAE
- Netuma osakae** Jordan and Kasawa, 1925
In: Jordan and Hubbs, Mem. Carnegie Mus.,
10(2): 157.
Holotype: FMNH 59388 (CM 7808). Japan:
market at Osaka.
ARIIDAE
- Netuma planifrons** Higuchi, Reis, and Araújo, 1982
Atlántica, Rio Grande, 5(1): 12.
Paratypes: 5. Brazil: Rio Grande do Sul, Lagoa
dos Patos.
ARIIDAE
- Nototropis forlonensis** Meek, 1904
Field Mus. Nat. Hist. Zool. Ser., 5: 70.
Holotype: FMNH 4478. Mexico: Tamaulipas,
Forlón.
Paratypes: 28. Mexico: 2 localities.
CYPRINIDAE
- Nototropis nazas** Meek, 1904
Field Mus. Nat. Hist. Zool. Ser., 5: 70.
Holotype: FMNH 4375. Mexico: Río Nazas at
Santiago Papasquiaro.
Paratypes: 133. Same locality.
CYPRINIDAE
- Notropis aztecus** Woolman, 1894
Bull. U.S. Fish Comm. 1894, 14: 63.
Syntypes: 11, FMNH 6604. Mexico: Canal in
Mexico City.
CYPRINIDAE
- Notropis chihuahua** Woolman, 1894
Am. Nat., 26(3): 260.
Syntype: FMNH 1889. Mexico: Chihuahua, Río
de los Conchos.
CYPRINIDAE
- Notropis edwardraneyi** Suttkus and Clemmer, 1968
Tulane Stud. Zool. Bot., 15(1): 19.
Paratypes: 100. Alabama: Alabama River, at
Yellow Jacket Bar.
CYPRINIDAE

- Notropis greenei** Hubbs and Ortenburger, 1929
 Publ. Univ. Okla. Biol. Surv., 1: 78.
 Paratypes: 37. Arkansas: Kinderhook.
 CYPRINIDAE
- Notropis muskoka** Meek, 1899
 Field Mus. Nat. Hist. Zool. Ser., 1(17): 308.
 Holotype: FMNH 2964. Canada: Ontario, Gull Lake.
 Paratypes: 20. Same locality.
 CYPRINIDAE
- Notropis panarcys** Hubbs and Miller, 1978
 Copeia, 1978(4): 582.
 Paratypes: 10. Mexico: Chihuahua, Río San Pedro at Meoqui.
 CYPRINIDAE
- Notropis robustus** Meek, 1902
 Field Mus. Nat. Hist. Zool. Ser., 3(6): 82.
 Holotype: FMNH 3548. Mexico: Río Conchos at Jiménez.
 Paratypes: 19. Mexico: 2 localities.
 CYPRINIDAE
- Notropis saladonis** Hubbs and Hubbs, 1958
 Copeia, 1958(4): 297.
 Paratypes: 18. Mexico: 2 localities.
 CYPRINIDAE
- Notropis santarosaliae** Meek, 1902
 Field Mus. Nat. Hist. Zool. Ser., 3(6): 85.
 Holotype: FMNH 3535. Mexico: Río Noavaco at Santa Rosalia.
 Paratypes: 17. Same locality.
 CYPRINIDAE
- Noturus albater** Taylor, 1969
 U.S. Natl. Mus. Bull., 282: 144.
 Paratypes: 4. Arkansas: Main Fork White River at Fayetteville.
 ICTALURIDAE
- Noturus flavater** Taylor, 1969
 U.S. Natl. Mus. Bull., 282: 104.
 Paratype. Missouri: Shannon County, Montier.
 ICTALURIDAE
- Noturus munitus** Suttkus and Taylor, 1965
 Proc. Biol. Soc. Wash., 78: 171.
 Paratypes: 10. Mississippi: Marion County, Pearl River.
 ICTALURIDAE
- Novumbra hubbsi** Schultz, 1929
 Publ. Univ. Wash. Coll. Fish., 2(6): 4.
 Paratype. Washington: one-half mile west of Satsop.
 UMBRIDAE
- Ochmacanthus flabelliferus** Eigenmann, 1912
 Mem. Carnegie Mus., 5: 213.
 Holotype: FMNH 53263 (CM 1729). Guyana: Konawaruk.
 TRICHOMYCTERIDAE
- Odontognathus compressus** Meek and Hildebrand, 1923
 Field Mus. Nat. Hist. Zool. Ser., 15(1): 194.
 Paratypes: 23. Panama: Fox Bay, Colón.
 CLUPEIDAE
- Odontorhamphus cancellori** Weed, 1933
 Field Mus. Nat. Hist. Zool. Ser., 20(7): 52.
 Holotype: FMNH 16208. Cook Archipelago: Aitutaki Island.
 HEMIRAMPHIDAE
- Odontostilbe hastatus** Eigenmann, 1913
 Indiana Univ. Stud., 18: 27.
 Holotype: FMNH 56383 (CM 5103). Colombia: Soplaviento.
 Paratypes: 13. Colombia: Bernal Creek.
 CHARACIDAE
- Odontostilbe melandetus** Eigenmann, 1912
 Mem. Carnegie Mus., 5: 312.
 Holotype: (CM 1878) MISSING. Guyana.
 CHARACIDAE
- Odontostilbe paraguayensis** Eigenmann and Kennedy, 1903
 Proc. Acad. Nat. Sci. Phila., 59: 512.
 Paratype. Paraguay: Arroyo Trementina.
 CHARACIDAE
- Odontostilbe trementinae** Eigenmann and Kennedy, 1903
 Proc. Acad. Nat. Sci. Phila., 59: 513.
 Paratypes: 2. Paraguay: Arroyo Trementina.
 CHARACIDAE
- Ogocephalus corniger** Bradbury, 1980
 Proc. Calif. Acad. Sci., 42(7): 274.
 Paratypes: 23. Caribbean Sea and Atlantic Ocean: 12 localities.
 OGCOEPHALIDAE

- Ogcocephalus declivirostris** Bradbury, 1980
Proc. Calif. Acad. Sci., 42(7): 269.
Holotype: FMNH 64136. Gulf of Mexico: off Texas.
Paratypes: 21. Gulf of Mexico: 14 localities.
OGCOCEPHALIDAE
- Ogcocephalus pantostictus** Bradbury, 1980
Proc. Calif. Acad. Sci., 42(7): 264.
Holotype: FMNH 50765. Florida: Pensacola, Santa Rosa Sound.
Paratypes: 8. Gulf of Mexico: 6 localities.
OGCOCEPHALIDAE
- Ogcocephalus pumilus** Bradbury, 1980
Proc. Calif. Acad. Sci., 42(7): 272.
Paratypes: 75. Caribbean Sea: 14 localities.
OGCOCEPHALIDAE
- Ogcocephalus rostellum** Bradbury, 1980
Proc. Calif. Acad. Sci., 42(7): 267.
Paratypes: 17. Atlantic Ocean: North Carolina to Florida, 4 localities.
OGCOCEPHALIDAE
- Oligobrycon microstomus** Eigenmann, 1915
Mem. Carnegie Mus., 7(1): 57.
Holotype: FMNH 57913 (CM 6898). Brazil: Rio Paratyba at Jacarehy.
Paratype. Same locality.
CHARACIDAE
- Ompok sabanus** Inger and Chin, 1959
Fieldiana Zool., 39(27): 282.
Holotype: FMNH 44828. Malaysia: Sabah, Segama River near Lahad Datu.
Paratypes: 106. Malaysia: Sabah, 15 localities.
SILURIDAE
- Oncorhynchus adonis** Jordan and McGregor, 1925
In: Jordan and Hubbs, Mem. Carnegie Mus., 10(2): 127.
Holotype: FMNH 58680 (CM 7784). Japan: Sagami, Lake Hakone.
SALMONIDAE
- Oncorhynchus ishikawae** Jordan and McGregor, 1925
In: Jordan and Hubbs, Mem. Carnegie Mus., 10(2): 132.
Holotype: FMNH 58682 (CM 7786). Japan: Lake Biwa.
SALMONIDAE
- Oncorhynchus kawamurae** Jordan and McGregor, 1925
In: Jordan and Hubbs, Mem. Carnegie Mus., 10(2): 128.
Holotype: FMNH 58681 (CM 7785). Japan: Ugo, Lake Toyama.
SALMONIDAE
- Oncorhynchus rhodurus** Jordan and McGregor, 1925
In: Jordan and Hubbs, Mem. Carnegie Mus., 10(2): 137.
Holotype: FMNH 59389 (CM 7794). Japan: Sagami, Lake Hakone.
Paratype. Same locality.
SALMONIDAE
- Ophichthus evermanni** Jordan and Richardson, 1909
Mem. Carnegie Mus., 4(4): 172.
Holotype: FMNH 52118 (CM 246). Taiwan: Takao.
OPHICHTHIDAE
- Ophichthus hyposagmatus** McCosker and Böhlke, 1984
Proc. Acad. Nat. Sci. Phila., 136: 24.
Paratype. Caribbean Sea: between Honduras and Jamaica.
OPHICHTHIDAE
- Ophioscion punctatissimus** Meek and Hildebrand, 1925
Field Mus. Nat. Hist. Zool. Ser., 15(2): 644.
Paratypes: 2. Panama: 2 localities.
SCIAENIDAE
- Ophioscion venezuelae** Schultz, 1945
Proc. U.S. Natl. Mus., 96(3192): 131.
Paratype. Venezuela: Caño de Sagua north of Sinamaica.
SCIAENIDAE
- Opisthonema berlangai** Berry and Barrett, 1963
Inter-Am. Trop. Tuna Comm. Bull., 7(2): 119.
Paratypes: 6. Ecuador: Islas Galápagos, Isla Barrington (Santa Fe).
CLUPEIDAE
- Opisthonema medirastre** Berry and Barrett, 1963
Inter-Am. Trop. Tuna Comm. Bull., 7(2): 118.
Paratypes: 5. Panama and Mexico: 2 localities.
CLUPEIDAE

- Optonurus atherodon** Gilbert and Cramer, 1896
Proc. U.S. Natl. Mus., 19(1114): 431.
Paratype. Hawaiian Islands.
MACROURIDAE
- Oreias furcatus** Banareescu and Nalbant, 1976
Nimphaca, 4: 188.
Paratypes: 4. China: Szechwan, Baurong.
COBITIDAE
- Orestias mundus** Parenti, 1984
Bull. Am. Mus. Nat. Hist., 178(2): 199.
Holotype: FMNH 41135. Peru: Cuzco.
Paratypes: 110. Same locality.
CYPRINODONTIDAE
- Ostichthys acanthorhinus** Randall, Shimizu, and Yamakawa, 1982
Jpn. J. Ichthyol., 29(1): 10.
Holotype: FMNH 70251. Arabian Sea: Gulf of Oman.
Paratypes: 3. Arabian Sea: Gulf of Oman, 3 localities.
HOLOCENTRIDAE
- Otolithes lini** Herre, 1935
Lingnan Sci. J. Canton, 14(4): 603.
Paratype. China: Canton Market.
SCIAENIDAE
- Otophidium schmidtii** Woods and Kanazawa, 1951
Fieldiana Zool., 31(53): 640.
Holotype: FMNH 48175. Bermuda: Castle Harbor, Cock Rock.
OPHIDIIDAE
- Oxyloricaria citurensis** Meek and Hildebrand, 1913
Field Mus. Nat. Hist. Zool. Ser., 10(8): 82.
Holotype: FMNH 7585. Panama: Río Cupe at Cituro.
Paratypes: 150. Panama: 5 localities.
LORICARIIDAE
- Oxyloricaria dariensis** Meek and Hildebrand, 1913
Field Mus. Nat. Hist. Zool. Ser., 10(8): 81.
Holotype: FMNH 7584 MISSING or mixed.
Panama: Río Tuyra, Boca de Cupe.
Paratypes: 39. Panama: 7 localities.
LORICARIIDAE
- Pangasius tubbi** Inger and Chin, 1959
Fieldiana Zool., 39(27): 287.
Holotype: FMNH 68047. Malaysia: Sabah, confluence of Deramakot River with Kinabatangan River.
Paratypes: 18. Malaysia: Sabah, 4 localities.
PANGASIIDAE
- Pantosteus jordani** Evermann, 1893
Bull. U.S. Fish Comm. 1892, 12: 51.
Paratypes: 2 (disintegrated). South Dakota: Hat Creek, Ardmore.
CATOSTOMIDAE
- Pantosteus santaanae** Snyder, 1908
Proc. U.S. Natl. Mus., 34(1595): 33.
Paratypes: 6. California: Santa Ana River near Riverside.
CATOSTOMIDAE
- Paraclinus integripinnis zacae** Hubbs, 1952
Stanford Ichthyol. Bull., 4(2): 87.
Paratypes: 18. Mexico: Baja California, Punta Aberrojos.
LABRISOMIDAE
- Paracrossochilus acerus** Inger and Chin, 1962
Fieldiana Zool., 45: 100.
Holotype: FMNH 68240. Malaysia: Sarawak, Sungei Dapu River.
Paratypes: 600. Malaysia: Sarawak, 10 localities.
CYPRINIDAE
- Parahollandia schmidtii** Woods, 1959
Copeia, 1959(3): 223.
Paratypes: 7. Western Caribbean Sea: 3 localities.
TRIACANTHODIDAE
- Paralepis philippinus** Fowler, 1934
Proc. Acad. Nat. Sci. Phila., 85: 281.
Paratypes: 2. Philippine Islands: Leyte.
PARALEPIDAE
- Parapelecus eigenmanni** Jordan and Metz, 1913
Mem. Carnegie Mus., 6(1): 21.
Holotype: FMNH 55802 (CM 4565). Korea: Suigen.
CYPRINIDAE
- Parascolopsis townsendi** Boulenger, 1901
Ann. Mag. Nat. Hist., Ser. 7, 7: 262.
Syntype: FMNH 5736. Sea of Oman.
NEMIPTERIDAE
- Parastremma sadina** Eigenmann, 1912
Indiana Univ. Stud., 16: 20.
Holotype: FMNH 56022 (CM 4812). Colombia: Istmina.
Paratypes: 11. Same locality.
CHARACIDAE

Paratrachichthys argyrophanus Woods, 1961

Fieldiana Zool., 39(45): 525.

Paratypes: 13. Atlantic Ocean: off Brazil, 3 localities.

TRACHICHTHYIDAE

Parecbasis cyclolepis Eigenmann, 1914

Indiana Univ. Stud., 20: 45.

Holotype: FMNH 56677 (CM 5495). Brazil: São Antonio de Rio Madeira.

Paratype. Same locality.

CHARACIDAE

Parodon bifasciatus Eigenmann, 1912

Mem. Carnegie Mus., 5: 274.

Holotype: FMNH 53582 (CM 1925). Guyana: Maripicru Creek.

PARODONTIDAE

Parodon dariensis Meek and Hildebrand, 1913

Field Mus. Nat. Hist. Zool. Ser., 10(8): 83.

Holotype: FMNH 7587. Panama: Río Cupe at Cituro.

Paratype. Same locality.

PARODONTIDAE

Parodon ecuadoriensis Eigenmann and Henn, 1914

In: Eigenmann, Henn, and Wilson, Indiana Univ. Stud., 19: 12.

Holotype: FMNH 56597 (CM 5413). Ecuador: Vinces.

Paratypes: 8. Ecuador: 2 localities.

PARODONTIDAE

Parodon terminalis Eigenmann and Henn, 1914

In: Eigenmann, Henn, and Wilson, Indiana Univ. Stud., 19: 12.

Holotype: FMNH 56599 (CM 5415). Ecuador: Vinces.

Paratype. Same locality.

PARODONTIDAE

Parotocinclus jimi Garavello, 1976

Arq. Zool. (São Paulo), 28(4): 9.

Paratypes: 3. Brazil: Bahia, Rio do Peixe at Fazenda Pedra Branca, Itagiba.

LORICARIIDAE

Pellonula tenuis Nichols and Griscom, 1917

Bull. Am. Mus. Nat. Hist., 37(25): 675.

Paratypes: 5. Zaïre: Stanleyville.

CLUPEIDAE

Pempheris dispar Herre, 1935

Field Mus. Nat. Hist. Zool. Ser., 18(12): 407.

Holotype: FMNH 17326. Solomon Islands: Isabel Island, Tenibuli.

Paratypes: 6. Same locality.

PEMPHERIDAE

Peristedion gilberti Jordan, 1922

Proc. U.S. Natl. Mus., 59(2392): 655.

Paratype. Hawaiian Islands: deep water of Oahu.

TRIGLIDAE

Peristedion greyae Miller, 1967

Bull. Mar. Sci., 17(1): 27.

Paratypes: 2. Western North Atlantic and Caribbean Sea: 2 localities.

TRIGLIDAE

Petersius barnardi Herre, 1936

Proc. Biol. Soc. Wash., 49: 100.

Paratypes: 8. Mozambique: near Beira.

CHARACIDAE

Petersius conserialis Hilgendorf, 1894

Sitzungsber. Ges. naturf. Freunde Berl., 1894: 173.

Paratype. Tanzania: Kingani River at Dunda.

CHARACIDAE

Petroscirtes kulambangrae Herre, 1935

Field Mus. Nat. Hist. Zool. Ser., 18(12): 436.

Holotype: FMNH 17392. Solomon Islands: Kulambangra Island.

BLENNIIDAE

Phalloptychus eigenmanni Henn, 1916

Ann. Carnegie Mus., 10(1-2): 121.

Holotype: FMNH 55876 (CM 4665). Brazil: Rio Catu at Alagoinhas.

Paratypes: 10. Same locality.

POECILIIDAE

Phallotorynus fasciolatus Henn, 1916

Ann. Carnegie Mus., 10(1-2): 129.

Holotype: FMNH 55061 (CM 3752). Brazil: São Paulo, Jacarehy.

Paratypes: 10. Same locality.

POECILIIDAE

Phenacogaster beni Eigenmann, 1911

Ann. Carnegie Mus., 8(1): 174.

Holotype: FMNH 54597 (CM 3229). Bolivia: Río Beni at Villa Bella.

Paratype. Same locality.

CHARACIDAE

Phenacogaster franciscoensis Eigenmann, 1911
Ann. Carnegie Mus., 8(1): 173.
Holotype: FMNH 54599 (CM 3231). Brazil: Boqueirão, near mouth of Rio Preto.
Paratypes: 16. Brazil: 4 localities.
CHARACIDAE

Phenacogaster megalostictus Eigenmann, 1909
Ann. Carnegie Mus., 6(1): 29.
Holotype: FMNH 53530 (CM 1056). Guyana: Tumatumari.
Paratypes: 47. Guyana: 6 localities.
CHARACIDAE

Phenacogaster microstictus Eigenmann, 1909
Ann. Carnegie Mus., 6(1): 30.
Holotype: FMNH 52970 (CM 1063). Guyana: Potaro River at Tumatumari.
Paratypes: 5. Guyana: 3 localities.
CHARACIDAE

Phenagoniates wilsoni Eigenmann, 1914
In: Eigenmann, Henn, and Wilson, Indiana Univ. Stud., 19: 2.
Holotype: FMNH 56540 (CM 5354). Colombia: Manigrú.
Paratypes: 2. Colombia: 2 localities.
CHARACIDAE

Phoxinus cumberlandensis Starnes and Starnes, 1978
Copeia, 1978(3): 509.
Paratypes: 16. Kentucky: South Fork, Dog Slaughter Creek.
CYPRINIDAE

Phoxinus orcuttii Eigenmann and Eigenmann, 1890
Proc. Calif. Acad. Sci., Ser. 2, 3: 2.
Syntypes: 10, FMNH 54985 (CM 3675). California: Temecula River at Riverside.
CYPRINIDAE

Phoxiscus kikuchii Oshima, 1919
Ann. Carnegie Mus., 12(2–4): 226.
Holotype: FMNH 59099 (CM 8237). Taiwan: Bokusekikaku.
CYPRINIDAE

Phrynelox atra Schultz, 1957
Proc. U.S. Natl. Mus., 107(3383): 76.
Holotype: FMNH 21705. Australia: Sydney.
Paratype. Same locality.
ANTENNARIIDAE

Phrynelox zebrinus Schultz, 1957
Proc. U.S. Natl. Mus., 107(3383): 75.
Paratypes: 2. Australia: 2 localities.
ANTENNARIIDAE

Physailia villiersi Boulenger, 1912
Ann. Mus. Congo Belge Ser. Quarto Zool., Ser. 1, 2(3): 17.
Syntypes: 2, FMNH 56151 (CM 4942). Zaïre: Luculla River.
SCHILBIDAE

Physiculus grinnelli Jordan and Jordan, 1922
Mem. Carnegie Mus., 10(1): 22.
Holotype: FMNH 55192 (CM 3898). Hawaiian Islands: Honolulu Market.
MORIDAE

Piabina analis Eigenmann, 1914
In: Eigenmann, Henn, and Wilson, Indiana Univ. Stud., 19: 8.
Holotype: FMNH 56661 (CM 5478). Brazil: Cáceres.
CHARACIDAE

Pikea aurora Jordan and Evermann, 1903
Bull. U.S. Fish Comm. 1902, 22: 178.
Paratype. Hawaiian Islands: Honolulu.
SERRANIDAE

Pikea mexicanus atlanticus Schultz, 1958
Proc. U.S. Natl. Mus., 108(3405): 326.
Paratype. Atlantic Ocean: off Surinam.
SERRANIDAE

Pikea mexicanus mexicanus Schultz, 1958
Proc. U.S. Natl. Mus., 108(3405): 323.
Paratype. Gulf of Mexico: off Florida, south of Panama City.
SERRANIDAE

Pimelodella avanhandavae Eigenmann, 1917
Mem. Carnegie Mus., 7(4): 240.
Holotype: FMNH 57981 (CM 6969). Brazil: Rio Tietê above Salto Avanhandava.
Paratypes: 30. Brazil: 2 localities.
PIMELODIDAE

Pimelodella boliviana Eigenmann, 1917
Mem. Carnegie Mus., 7(4): 245.
Holotype: FMNH 57976 (CM 6964). Bolivia: Santa Cruz de la Sierra.
Paratype. Same locality.
PIMELODIDAE

- Pimelodella eigenmanni* Meek, 1905**
 Proc. Biol. Soc. Wash., 18: 241.
 Holotype: FMNH 3400. Brazil: São Paulo.
 Paratypes: 5. Same locality.
 PIMELODIDAE
- Pimelodella griffini* Eigenmann, 1917**
 Mem. Carnegie Mus., 7(4): 250.
 Holotype: FMNH 57974 (CM 6962). Paraguay:
 near Sapucay.
 Paratypes: 7. Same locality.
 PIMELODIDAE
- Pimelodella hasemani* Eigenmann, 1917**
 Mem. Carnegie Mus., 7(4): 241.
 Holotype: FMNH 57980 (CM 6968). Brazil: São
 Antonio de Rio Madeira.
 PIMELODIDAE
- Pimelodella itapicuruensis* Eigenmann, 1917**
 Mem. Carnegie Mus., 7(4): 247.
 Holotype: FMNH 57986 (CM 6974). Brazil: Rio
 Itapicuru at Queimadas.
 Paratypes: 21. Brazil: 4 localities.
 PIMELODIDAE
- Pimelodella laticeps* Eigenmann, 1917**
 Mem. Carnegie Mus., 7(4): 243.
 Holotype: FMNH 57969 (CM 6957). Brazil:
 Uruguayana.
 Paratypes: 3. Same locality.
 PIMELODIDAE
- Pimelodella laticeps australis* Eigenmann, 1917**
 Mem. Carnegie Mus., 7(4): 243.
 Holotype: FMNH 57962 (CM 6950). Brazil:
 Uruguayana.
 Paratypes: 43. Same locality.
 PIMELODIDAE
- Pimelodella macturki* Eigenmann, 1912**
 Mem. Carnegie Mus., 5: 170.
 Holotype: FMNH 53234 (CM 1695). Guyana:
 creek in Mora Passage.
 Paratypes: 11. Guyana: 3 localities.
 PIMELODIDAE
- Pimelodella megalops* Eigenmann, 1912**
 Mem. Carnegie Mus., 5: 169.
 Holotype: FMNH 53231 (CM 1692). Guyana:
 Tumatumari.
 Paratypes: 35. Guyana: 2 localities.
 PIMELODIDAE
- Pimelodella metae* Eigenmann, 1917**
 Mem. Carnegie Mus., 7(4): 244.
 Holotype: FMNH 58441 (CM 7441). Colombia:
 Río Negro at Villavicencio.
 Paratypes: 2. Colombia: Río Meta at Barrigona.
 PIMELODIDAE
- Pimelodella notomelas* Eigenmann, 1917**
 Mem. Carnegie Mus., 7(4): 244.
 Holotype: FMNH 57967 (CM 6955). Brazil:
 Cáceres.
 Paratypes: 9. Brazil: 2 localities.
 PIMELODIDAE
- Pimelodella serrata* Eigenmann, 1917**
 Mem. Carnegie Mus., 7(4): 235.
 Holotype: FMNH 57979 (CM 6967). Bolivia:
 San Joaquín.
 Paratype. Same locality.
 PIMELODIDAE
- Pimelodus clarias coprophagus* Schultz, 1944**
 Proc. U.S. Natl. Mus., 94(3172): 203.
 Paratypes: 4. Venezuela: Río Palmar near To-
 tuma.
 PIMELODIDAE
- Pimelodus heteropleurus* Eigenmann, 1912**
 Mem. Carnegie Mus., 5: 176.
 Holotype: FMNH 53336 (CM 1734). Guyana:
 Rupununi Pan.
 PIMELODIDAE
- Pimelodus ortmanni* Eigenmann, 1911**
 Ann. Carnegie Mus., 7(3-4): 379.
 Holotype: FMNH 54240 (CM 2856). Brazil: Rio
 Iguaçu at Pôrto União da Victoria.
 Paratypes: 5. Same locality.
 PIMELODIDAE
- Pimephales anuli* Kendall, 1903**
 Bull. U.S. Fish Comm. 1902, 22: 360.
 Paratypes: 15. Maine: Lunkasoos Lake.
 CYPRINIDAE
- Plagiotremus phenax* Smith-Vaniz, 1976**
 Monogr. Acad. Nat. Sci. Phila., 19: 131.
 Paratypes: 11. Maldive Islands: 3 localities.
 BLENNIIDAE
- Platophrys inermis* Gilbert, 1905**
 Bull. U.S. Fish Comm. 1903, 23(2): 685.
 Paratype. Hawaiian Islands.
 BOTHIDAE

Platypoecilus tropicus Meek, 1907

Field Mus. Nat. Hist. Zool. Ser., 7(5): 146.
Holotype: FMNH 6027. Costa Rica: Turrialba.
Paratypes: 2. Same locality.
POECILIIDAE

Platypoecilus variatus Meek, 1904

Field Mus. Nat. Hist. Zool. Ser., 5: 146.
Holotype: FMNH 4501. Mexico: San Luis Potosí, Valles.
Paratypes: 109. Mexico: 5 localities.
POECILIIDAE

Platysilurus barbatus Haseman, 1911

Ann. Carnegie Mus., 7(3–4): 320.
Holotype: FMNH 60306 (CM 2987). Brazil: São Antonio de Rio Madeira.
Paratypes: 2. Bolivia: Río Machupo near San Joaquín.
PIMELODIDAE

Plecostomus derbyi Haseman, 1911

Ann. Carnegie Mus., 7(3–4): 384.
Holotype: FMNH 54246 (CM 2865). Brazil: Rio Iguaçu at Pôrto União da Victoria.
LORICARIIDAE

Plecostomus hemiurus Eigenmann, 1912

Mem. Carnegie Mus., 5: 224.
Holotype: FMNH 53110 (CM 1544). Guyana: Amaturuk.
Paratypes: 10. Guyana: 6 localities.
LORICARIIDAE

Plectranthias garrupellus Robins and Starck, 1961

Proc. Acad. Nat. Sci. Phila., 113(11): 295.
Paratype. Western North Atlantic: no station data.
SERRANIDAE

Plesiops cephalotaenia Inger, 1955

Pac. Sci., 9(3): 272.
Holotype: FMNH 44708. Philippine Islands: Sitangkai.
Paratypes: 3. Same locality.
PLESIOPIDAE

Pleurogrammus azonus Jordan and Metz, 1913

Mem. Carnegie Mus., 6(1): 47.
Holotype: FMNH 55796 (CM 4558). Korea: Chinnampo.
HEXAGRAMMIDAE

Plotosus papuensis Weber, 1910

Notes Leiden Mus., 32: 228.
Syntype: FMNH 52494 (CM 752). New Guinea: Lorentz River.
PLOTOSIDAE

Pneumatophorus peruanus Jordan and Hubbs, 1925

Mem. Carnegie Mus., 10(2): 211.
Holotype: FMNH 58739 (CM 7847). Peru: Callao.
SCOMBRIDAE

Poblana ferdebueni Solórzano and López, 1965

Ciencia (Mex. City), 23(5): 209.
Paratypes: 3. Mexico: Puebla, Laguna de Almeyala o Chignahuapan.
ATHERINIDAE

Poecilia amates Miller, 1907

Bull. Am. Mus. Nat. Hist., 23(2): 108.
Paratypes: 6. Guatemala: Los Amates.
POECILIIDAE

Poecilia caudata Meek, 1909

Field Mus. Nat. Hist. Zool. Ser., 7(7): 209.
Holotype: FMNH 6360. Costa Rica: Turruabares.
Allotype and paratypes: 10. Costa Rica: 3 localities.
POECILIIDAE

Poecilia fria Eigenmann and Henn, 1914

In: Eigenmann, Henn, and Wilson, Indiana Univ. Stud., 19: 13.
Holotype: FMNH 56603 (CM 5420). Ecuador: Vinces.
Paratypes: 112. Same locality.
POECILIIDAE

Poecilia latipunctata Meek, 1904

Field Mus. Nat. Hist. Zool. Ser., 5: 150.
Holotype: FMNH 4484. Mexico: Tamaulipas, Forlón.
Paratypes: 12. Same locality.
POECILIIDAE

Poecilia pittieri Meek, 1912

Field Mus. Nat. Hist. Zool. Ser., 10(7): 71.
Holotype: FMNH 7680. Costa Rica: Río Reventazón at La Junta.
Allotype and paratypes: 15. Same locality.
POECILIIDAE

Poecilia tenuis Meek, 1907

Field Mus. Nat. Hist. Zool. Ser., 7(5): 147.
Holotype (?): FMNH 6028. Costa Rica: Tiribi.
Paratypes: 4. Costa Rica: 2 localities.

See comment in Grey (1947, p. 175). One of
2 specimens in FMNH 6029 is also la-
beled "type", and is closer to the size
given for the type in Meek (1907).

POECILIIDAE

Poeciliopsis balsas Hubbs, 1926

Misc. Publ. Mus. Zool. Univ. Mich., 16: 66.
Holotype: FMNH 3702. Mexico: Balsas, Gue-
rrero.

Paratypes: 91. Mexico: 6 localities.

POECILIIDAE

Poecilobrycon auratus Eigenmann, 1909

Ann. Carnegie Mus., 6(1): 43.
Holotype: FMNH 52768 (CM 1161). Guyana:
Konawaruk.

Paratypes: 8. Guyana: 3 localities.

LEBIASINIDAE

Poecilobrycon erythrurus Eigenmann, 1909

Ann. Carnegie Mus., 6(1): 44.
Holotype: FMNH 52774 (CM 1168). Guyana:
Rockstone sand-bank.

Paratypes: 6. Guyana: 4 localities.

LEBIASINIDAE

Poecilobrycon harrisoni Eigenmann, 1909

Ann. Carnegie Mus., 6(1): 43.
Holotype: FMNH 52767 (CM 1160). Guyana:
canal at Christianburg.

LEBIASINIDAE

Poecilobrycon ocellatus Eigenmann, 1909

Ann. Carnegie Mus., 6(1): 45.
Holotype: FMNH 52975 (CM 1179). Guyana:
Wismar.

Paratypes: 25. Guyana: 6 localities.

LEBIASINIDAE

Poecilocharax bovalii Eigenmann, 1909

Ann. Carnegie Mus., 6(1): 34.
Holotype: FMNH 53544 (CM 1136). Guyana:
creek at Savanna Landing.

Paratypes: 171. Guyana: 4 localities.

CHARACIDAE

Poecilurichthys moenkhausii Eigenmann and
Kennedy, 1903

Proc. Acad. Nat. Sci. Phila., 59: 522.
Paratypes: 2. Paraguay: brook near Arroyo Tre-
mentina.

CHARACIDAE

Pogoneleotris microps Weber, 1908

Nova Guinea, 5(2): 258.

Syntypes: 2, FMNH 53459 (CM 717). New
Guinea: Merauke River.

ELEOTRIDAE

Polyacanthonotus merretti Sulak, Crabtree, and
Hureau, 1984

Cybium, 8(4): 59.

Paratype. Atlantic Ocean: off Virginia, Norfolk
Canyon.

NOTACANTHIDAE

Polyipnus nuttingii Gilbert, 1905

Bull. U.S. Fish Comm. 1903, 23(2): 609.

Paratypes: 6. Hawaiian Islands.

STERNOPTYCHIDAE

Pomacentrus cranei Herre, 1935

Field Mus. Nat. Hist. Zool. Ser., 18(12): 409.

Holotype: FMNH 17338. Solomon Islands: Is-
abel Isand, Tenibuli.

Paratype. Same locality.

POMACENTRIDAE

Pomacentrus jenkinsi Jordan and Evermann, 1903

Bull. U.S. Fish Comm. 1902, 22: 189.

Paratypes: 3. Hawaiian Islands.

POMACENTRIDAE

Pomacentrus reidi Fowler and Bean, 1928

Bull. U.S. Natl. Mus., 100(7): 99.

Paratypes: 2. Philippine Islands: Dalaganem Is-
land.

POMACENTRIDAE

Pomacentrus stigma Fowler and Bean, 1928

Bull. U.S. Natl. Mus., 100(7): 100.

Paratype. Philippine Islands: Mindoro, Paluan
Bay.

POMACENTRIDAE

Pomadasy grandis Meek, 1907

Field Mus. Nat. Hist. Zool. Ser., 7(4): 116.

Holotype: FMNH 5939. Nicaragua: Lago Ni-
caragua at Granada.

HAEMULIDAE

Pomadasy starri Meek, 1904

Field Mus. Nat. Hist. Zool. Ser., 5: 200.

Holotype: FMNH 4693. Mexico: Veracruz, Pé-
rez.

HAEMULIDAE

- Pomadasy templei** Meek, 1904
Field Mus. Nat. Hist. Zool. Ser., 5: 201.
Holotype: FMNH 4504. Mexico: San Luis Potosí, Valles.
Paratypes: 2. Mexico: Veracruz, Pérez.
HAEMULIDAE
- Porichthys bathoiketes** Gilbert, 1968
Bull. Mar. Sci., 18(3): 710.
Paratype. Caribbean Sea: off Honduras.
BATRACHOIDIDAE
- Porichthys kymosemeum** Gilbert, 1968
Bull. Mar. Sci., 18(3): 724.
Holotype: FMNH 66688. Atlantic Ocean: off Brazil.
BATRACHOIDIDAE
- Porichthys myriaster** Hubbs and Schultz, 1939
Proc. U.S. Natl. Mus., 86(3060): 478.
Paratypes: 4. California: off Mira Mar pier.
BATRACHOIDIDAE
- Porogadus silus** Carter and Sulak, 1984
Bull. Mar. Sci., 34(3): 361.
Paratypes: 2. Western North Atlantic: The Bahamas region.
OPHIDIIDAE
- Porotergus gimbeli** Ellis, 1912
Mem. Carnegie Mus., 5: 441.
Holotype: FMNH 54566 (CM 3197). Brazil: Pará.
Paratypes: 2. Same locality.
APTERONOTIDAE
- Porotergus gymnotus** Ellis, 1912
Mem. Carnegie Mus., 5: 441.
Holotype: FMNH 53575 (CM 1759). Guyana: Amatuk.
Paratype. Guyana: Konawaruk.
APTERONOTIDAE
- Priapichthys annectens hesperis** Hubbs, 1924
Misc. Publ. Mus. Zool. Univ. Mich., 13: 22.
Holotype: FMNH 14106. Costa Rica: San José.
Paratypes: 630. Costa Rica: 6 localities.
POECILIIDAE
- Priapichthys fosteri** Hildebrand, 1925
Bull. Bur. Fish., 41: 260.
Paratypes: 18. El Salvador: 2 localities.
POECILIIDAE
- Priapichthys letonai** Hildebrand, 1925
Bull. Bur. Fish., 41: 258.
Paratypes: 45. El Salvador: 5 localities.
POECILIIDAE
- Priapichthys olomina** Meek, 1914
Field Mus. Nat. Hist. Zool. Ser., 10(10): 114.
Holotype: FMNH 7827. Costa Rica: Orotina.
Allotype and paratypes: 132. Costa Rica: 4 localities.
The 16 "Pacific Side" paratypes which Grey (1947, p. 178) noted as MISSING are probably among the 7 lots (29 specimens) collected by Alfaro that are now in our main collection as *Brachyrhaphis olomina*.
POECILIIDAE
- Priapichthys panamensis** Meek and Hildebrand, 1916
Field Mus. Nat. Hist. Zool. Ser., 10(15): 322.
Holotype: FMNH 8950. Panama: Chame Point.
Allotype and paratypes: 17. Panama: 2 localities.
POECILIIDAE
- Prionodes baldwini** Evermann and Marsh, 1899
Rept. U.S. Fish Comm. 1899, 25: 353.
Paratypes: 5. Puerto Rico: 2 localities.
SERRANIDAE
- Prionotus aspersus** Meek and Hildebrand, 1928
Field Mus. Nat. Hist. Zool. Ser., 15(3): 856.
Paratypes: 3. Panama: 3 localities.
TRIGLIDAE
- Pristella aubynei** Eigenmann, 1909
Ann. Carnegie Mus., 6(1): 24.
Holotype: FMNH 52698 (CM 1042). Guyana: Lama Stop-off.
Paratypes: 47. Guyana: 3 localities.
CHARACIDAE
- Pristipomoides freemani** Anderson, 1966
Bull. Mar. Sci., 16(4): 816.
Paratype. Caribbean Sea: off Colombia.
LUTJANIDAE
- Pristiurus sauteri** Jordan and Richardson, 1909
Mem. Carnegie Mus., 4(4): 160.
Holotype: FMNH 55193 (CM 219). Taiwan: Takao.
Paratype. Same locality.
SCYLIORHINIDAE

- Probolodus heterostomus** Eigenmann, 1911
Ann. Carnegie Mus., 8(1): 164.
Holotype: FMNH 54328 (CM 2973). Brazil: Campos.
Paratypes: 5. Brazil: 2 localities.
CHARACIDAE
- Prochilodus maripicru** Eigenmann, 1912
Mem. Carnegie Mus., 5: 271.
Holotype: FMNH 53597 (CM 2066). Guyana: Maripicru Creek.
Paratype. Same locality.
PROCHILODONTIDAE
- Profundulus candalarius** Hubbs, 1924
Misc. Publ. Mus. Zool. Univ. Mich., 13: 15.
Holotype: FMNH 9071. Guatemala: spring at Candalaria.
Allotype. Same locality.
CYPRINODONTIDAE
- Protomyzon aphelocheilus** Inger and Chin, 1962
Fieldiana Zool., 45: 110.
Holotype: FMNH 68166. Malaysia: Sabah, Sungei Kaingeran.
Paratypes: 9. Same locality.
HOMALOPTERIDAE
- Psalidodon gymnodontus** Eigenmann, 1911
Ann. Carnegie Mus., 8(1): 166.
Holotype: FMNH 54574 (CM 3204). Brazil: Rio Iguaçu at Pôrto União.
CHARACIDAE
- Psectrogaster curviventris** Eigenmann and Kennedy, 1903
Proc. Acad. Nat. Sci. Phila., 59: 509.
Paratype. Paraguay: Asunción.
CURIMATIDAE
- Pseudancistrus carnegiei** Eigenmann, 1916
Ann. Carnegie Mus., 10(1-2): 85.
Holotype: FMNH 58350 (CM 7346). Colombia: Río San Gil, Santander.
Paratypes: 5. Colombia: Quebrada de Honda.
LORICARIIDAE
- Pseudancistrus nigrescens** Eigenmann, 1912
Mem. Carnegie Mus., 5: 234.
Holotype: FMNH 53105 (CM 1539). Guyana: Amatuik.
LORICARIIDAE
- Pseudancistrus pediculatus** Eigenmann, 1917
Proc. Am. Philos. Soc., 56(7): 679.
Holotype: FMNH 58352 (CM 7348). Colombia: Río Negro at Villavicencio.
Paratypes: 7. Colombia: 2 localities.
LORICARIIDAE
- Pseudanos irinae** Winterbottom, 1980
Life Sci. Contrib. R. Ont. Mus., 123: 27.
Paratypes: 3. Guyana: 3 localities.
ANOSTOMIDAE
- Pseudaspius atrilatus** Jordan and Thompson, 1914
Mem. Carnegie Mus., 6(4): 231.
Holotype: FMNH 57075 (CM 6012). Japan: Lake Biwa at Matsubara.
CYPRINIDAE
- Pseudaspius bergi** Jordan and Metz, 1913
Mem. Carnegie Mus., 6(1): 22.
Holotype: FMNH 55800 (CM 4563). Korea: Chinnampo.
Paratypes: 2. Same locality.
CYPRINIDAE
- Pseudaspius modestus** Jordan and Metz, 1913
Mem. Carnegie Mus., 6(1): 23.
Holotype: FMNH 55799 (CM 4561). Korea: Chinnampo.
Paratypes: 4. Same locality.
CYPRINIDAE
- Pseudauchenipterus nigrolineatus** Fowler, 1911
Proc. Acad. Nat. Sci. Phila., 63: 434.
Paratype. Venezuela: Pedernales.
AUCHENIPTERIDAE
- Pseudobagrus adiposalis** Oshima, 1919
Ann. Carnegie Mus., 12(2-4): 181.
Holotype: FMNH 59079 (CM 8216). Taiwan: Tamusui River near Shinten.
BAGRIDAE
- Pseudobagrus taiwanensis** Oshima, 1919
Ann. Carnegie Mus., 12(2-4): 180.
Holotype: FMNH 59078 (CM 8215). Taiwan: Tozen River near Taichu.
BAGRIDAE
- Pseudocheiroidon affinis** Meek and Hildebrand, 1916
Field Mus. Nat. Hist. Zool. Ser., 10(15): 275.
Holotype: FMNH 8944. Panama: Río Gatún, Monte Liría.
Paratypes: 251. Panama: 13 localities.
Published number FMNH 8941 is an error.
CHARACIDAE

Pseudochromis aurea marshallensis Schultz, 1953
Bull. U.S. Natl. Mus., 202(1): 392.
Paratypes: 2. Marshall Islands: 2 localities.
PSEUDOCROMIDAE

Pseudocorynopoma heterandria Eigenmann, 1914.
Indiana Univ. Stud., 20: 39.
Holotype: FMNH 56407 (CM 5222). Brazil:
Xiririca.
Paratypes: 8. Same locality.
CHARACIDAE

Pseudogobio chinssuensis Nichols, 1926
Am. Mus. Nov., 214: 3.
Paratype. China: Shansi, Chin-ssu.
CYPRINIDAE

Pseudoperilampus hondae Jordan and Metz, 1913
Mem. Carnegie Mus., 6(1): 20.
Holotype: FMNH 55803 (CM 4566). Korea:
Suigen.
CYPRINIDAE

Pseudopimelodus albomarginatus Eigenmann,
1912
Mem. Carnegie Mus., 5: 153.
Holotype: FMNH 53221 (CM 1680). Guyana:
Tukeit.
Paratypes: 10. Guyana: 2 localities.
PIMELODIDAE

Pseudopimelodus villosus Eigenmann, 1912
Mem. Carnegie Mus., 5: 152.
Holotype: FMNH 53219 (CM 1677). Guyana:
Potaro Landing.
Paratypes: 2. Guyana: 2 localities.
PIMELODIDAE

Pseudoplesiops sargenti Schultz, 1953
Bull. U.S. Natl. Mus., 202(1): 405.
Paratype. Philippine Islands: Endeavor Straits.
PSEUDOCROMIDAE

Pseudoraja fischeri Bigelow and Schroeder, 1954
Breviora, 24: 4.
Paratype. Gulf of Mexico: near Campeche Bank.
RAJIDAE

Pseudorasbora altipinna Nichols, 1925
Am. Mus. Nov., 182: 5.
Paratypes: 2. China: Szechwan, Yen-ching-kao.
CYPRINIDAE

Pseudorasbora depressirostris Nichols, 1925
Am. Mus. Nov., 182: 5.
Paratypes: 2. China: Shansi, Chin-ssu.
CYPRINIDAE

Pseudotatia parva Mees, 1974
Zool. Verh. (Leiden), 132: 105.
Holotype: FMNH 70580. Brazil: Joazeiro.
Paratypes: 13. Same locality.
AUCHENIPTERIDAE

Pseudotolithus brunneolus Jordan and Richardson,
1909
Mem. Carnegie Mus., 4(4): 191.
Holotype: FMNH 52174 (CM 327). Taiwan:
Takao.
Paratype. Same locality.
SCIAENIDAE

Pseudupeneus chrysonemus Jordan and Evermann, 1903
Bull. U.S. Fish Comm. 1902, 22: 186.
Paratypes: 2. Hawaiian Islands: Hilo.
MULLIDAE

Psilorhynchus gracilis Rainboth, 1983
Proc. Calif. Acad. Sci., 43(6): 69.
Paratypes: 5. Bangladesh: Mahananda River at
Tetulia.
PSILORHYNCHIDAE

Psychichthys eidolon Jordan and Hubbs, 1925
Mem. Carnegie Mus., 10(2): 117.
Holotype: FMNH 59519 (CM 7779). Japan: Sagami Bay, Izu off Mishima.
CHIMAERIDAE

Pterobrycon landoni Eigenmann, 1913
Indiana Univ. Stud., 18: 3.
Holotype: FMNH 56250 (CM 5051). Colombia:
Boca de Raspadura.
CHARACIDAE

Pterois elliotti Meek, 1897
Field Mus. Nat. Hist. Zool. Ser., 1(8): 169.
Holotype: FMNH 484. Somali: Berbera.
SCORPAENIDAE

Pterolebias hoignei Thomerson, 1974
Copeia, 1974(1): 31.
Paratypes: 9. Venezuela: Río Pao near Caño Benito.
APLOCHEILIDAE

Pterolebias maculipinnis Weibezahn, 1964
In: Radda, Die Aquar. un Terr. Zeitschrift,
17: 39–41 (not seen).
Neotype: FMNH 85267. Venezuela: Río Pao
drainage, Cojedes.

Neotype designated by Taphorn and Thom-
erson (1978, Acta Biol. Venez., 9(4): 395);
perhaps authorship should be attributed
to Radda (1964).

APLOCHEILIDAE

Pterygoplichthys anisitsi Eigenmann and Kenne-
dy, 1903

Proc. Acad. Nat. Sci. Phila., 59: 503.

Paratype. Paraguay: Asunción.

Misspelled *Pteryophlichthys* in Eigenmann and
Kennedy (1903).

LORICARIIDAE

Puntius joaquinae Wood, 1968

Philipp. J. Sci., 95(4): 415.

Holotype: FMNH 73395. Philippine Islands:
Mindanao, Agus River.

Paratypes: 5. Philippine Islands: 4 localities.

CYPRINIDAE

Puntius snyderi Oshima, 1919

Ann. Carnegie Mus., 12(2–4): 216.

Holotype: FMNH 59093 (CM 8231). Taiwan:
Nanto, Rigyokutsu.

CYPRINIDAE

Puntius umalii Wood, 1968

Philipp. J. Sci., 95(4): 412.

Holotype: FMNH 73393. Philippine Islands:
Mindanao, Agus River.

Paratypes: 12. Philippine Islands: 5 localities.

CYPRINIDAE

Pygidium alternatum Eigenmann, 1917

Proc. Am. Philos. Soc., 56(8): 700.

Holotype: FMNH 58082 (CM 7079). Brazil: Rio
Doce.

Paratypes: 62. Same locality.

TRICHOMYCTERIDAE

Pygidium banneaui Eigenmann, 1912

Indiana Univ. Stud., 16: 19.

Holotype: FMNH 56025 (CM 4815). Colombia:
Bernal Creek near Honda.

Paratypes: 54. Same locality.

TRICHOMYCTERIDAE

Pygidium barbouri Eigenmann, 1911

Ann. Carnegie Mus., 7(2): 214.

Paratypes: 5. Bolivia: upper Beni River.

TRICHOMYCTERIDAE

Pygidium bogotense Eigenmann, 1912

Indiana Univ. Stud., 16: 18.

Holotype: FMNH 56030 (CM 4820). Colombia:
Chapinero, near Bogotá.

Paratypes: 131. Colombia: 2 localities.

TRICHOMYCTERIDAE

Pygidium caliense Eigenmann, 1912

Indiana Univ. Stud., 16: 18.

Holotype: FMNH 56029 (CM 4819). Colombia:
Cali.

Published number CM 6819 is an error.

TRICHOMYCTERIDAE

Pygidium chapmani Eigenmann, 1912

Indiana Univ. Stud., 16: 18.

Holotype: FMNH 56027 (CM 4817). Colombia:
Boquia.

Paratypes: 26. Same locality.

TRICHOMYCTERIDAE

Pygidium conradi Eigenmann, 1912

Mem. Carnegie Mus., 5: 212.

Holotype: FMNH 53721 (CM 2212). Guyana:
Amatuk.

TRICHOMYCTERIDAE

Pygidium davisii Haseman, 1911

Ann. Carnegie Mus., 7(3–4): 380.

Holotype: FMNH 60309 (CM 2862). Brazil:
Serrinha Paraná.

Paratypes: 9. Brazil: creek near Serrinha Paraná.

TRICHOMYCTERIDAE

Pygidium dorsostriatum Eigenmann, 1917

Proc. Am. Philos. Soc., 56(8): 695.

Holotype: FMNH 58096 (CM 7093). Colombia:
Villavicencio.

Paratype. Same locality.

Species name was spelled *dorsotriatum* in Ei-
genmann (1917), apparently a typo-
graphic error emended in subsequent pa-
pers by Eigenmann.

TRICHOMYCTERIDAE

Pygidium gracilior Eigenmann, 1912

Mem. Carnegie Mus., 5: 213.

Holotype: FMNH 53264 (CM 1730). Guyana:
Erukin.

TRICHOMYCTERIDAE

- Pygidium guianensis** Eigenmann, 1909
Ann. Carnegie Mus., 6(1): 11.
Holotype: FMNH 52676 (CM 1003). Guyana:
Aruataima Falls.
TRICHOMYCTERIDAE
- Pygidium hasemani** Eigenmann, 1914
Indiana Univ. Stud., 20: 48.
Holotype: FMNH 56424 (CM 5238). Brazil:
Santarém.
Paratypes: 312. Same locality.
TRICHOMYCTERIDAE
- Pygidium iheringi** Eigenmann, 1917
Proc. Am. Philos. Soc., 56(8): 697.
Holotype: FMNH 58074 (CM 7071). Brazil: São
Paulo, Sapina.
Paratypes: 2. Same locality.
TRICHOMYCTERIDAE
- Pygidium latistriatum** Eigenmann, 1917
Proc. Am. Philos. Soc., 56(8): 696.
Holotype: FMNH 58449 (CM 7450). Colombia:
Quebrada de Pinchote, Santander.
TRICHOMYCTERIDAE
- Pygidium paolence** Eigenmann, 1917
Proc. Am. Philos. Soc., 56(8): 698.
Holotype: FMNH 58085 (CM 7081). Brazil: Rio
Tietê at Alto da Serra.
Paratype. Brazil: Rio Parahyba.
TRICHOMYCTERIDAE
- Pygidium proops parahybae** Eigenmann, 1918
Mem. Carnegie Mus., 7(5): 332.
Holotype: FMNH 58576 (CM 7598). Brazil: Rio
Parahyba at São João da Barra.
TRICHOMYCTERIDAE
- Pygidium punctulatum piurae** Eigenmann, 1922
Mem. Carnegie Mus., 9(1): 63.
Paratypes: 6. Peru: 3 localities.
TRICHOMYCTERIDAE
- Pygidium reinhardti** Eigenmann, 1917
Proc. Am. Philos. Soc., 56(8): 699.
Holotype: FMNH 58081 (CM 7078). Brazil: Rio
Itabira at Burmier.
TRICHOMYCTERIDAE
- Pygidium santaeritae** Eigenmann, 1918
Mem. Carnegie Mus., 7(5): 341.
Holotype: FMNH 58577 (CM 7599). Brazil: Rio
Preto at Santa Rita.
TRICHOMYCTERIDAE
- Pygidium septentrionale** Behre, 1928
Ann. Carnegie Mus., 18(2): 309.
Holotype: FMNH 59522 (CM 8515). Panama:
Salao Creek into Río Chiriquí del Tire.
Paratype. Panama: Sombrero Creek.
TRICHOMYCTERIDAE
- Pygidium stellatum** Eigenmann, 1918
Mem. Carnegie Mus., 7(5): 308.
Holotype: FMNH 58101 (CM 7097). Colombia:
Quebrada Sarjento.
Paratypes: 3. Same locality.
TRICHOMYCTERIDAE
- Pygidium straminium** Eigenmann, 1917
Proc. Am. Philos. Soc., 56(8): 694.
Holotype: FMNH 58105 (CM 7101). Colombia:
Quebrada del Mango.
Paratypes: 4. Colombia: 2 localities.
TRICHOMYCTERIDAE
- Pygidium striatum** Meek and Hildebrand, 1913
Field Mus. Nat. Hist. Zool. Ser., 10(8): 78.
Holotype: FMNH 7579. Panama: Río Cana at
Cana.
Paratypes: 18. Same locality.
TRICHOMYCTERIDAE
- Pygidium triguttatum** Eigenmann, 1918
Mem. Carnegie Mus., 7(5): 339.
Holotype: FMNH 58670 (CM 7600). Brazil: Ja-
carehy.
Paratypes: 3. Same locality.
TRICHOMYCTERIDAE
- Pygidium vermiculatum** Eigenmann, 1917
Proc. Am. Philos. Soc., 56(8): 699.
Holotype: FMNH 58077 (CM 7074). Brazil: Juiz
de Fora.
TRICHOMYCTERIDAE
- Pygidium zonatum** Eigenmann, 1918
Mem. Carnegie Mus., 7(5): 330.
Holotype: FMNH 58573 (CM 7596). Brazil:
Água Quente.
Paratypes: 3. Brazil: 2 localities.
TRICHOMYCTERIDAE
- Pygocentrus bilineatus** Eigenmann, 1909
Ann. Carnegie Mus., 6(1): 47.
Holotype: FMNH 58075 (CM 1072). Guyana:
Aruka River.
Paratype. Same locality.
CHARACIDAE

- Pyrrhulina australe** Eigenmann and Kennedy, 1903
Proc. Acad. Nat. Sci. Phila., 59: 508.
Paratypes: 2. Paraguay: Arroyo Trementina.
LEBIASINIDAE
- Quisquilius eugenius** Jordan and Evermann, 1903
Bull. U.S. Fish Comm. 1902, 22: 203.
Paratype: MISSING. Hawaiian Islands: Honolulu.
GOBIIDAE
- Rabula fuscomaculata** Schultz, 1953
Bull. U.S. Natl. Mus., 202(1): 147.
Paratypes: 2. Marshall Islands: Bikini Atoll, 2 localities.
MURAENIDAE
- Rabula marshallensis** Schultz, 1953
Bull. U.S. Natl. Mus., 202(1): 149.
Paratype: Marshall Islands: Kwajalein Atoll, Ennylabegan Island.
MURAENIDAE
- Rachovia pyropunctata** Taphorn and Thomerson, 1978
Acta Biol. Venez., 9(4): 403.
Holotype: FMNH 85713. Venezuela: west of Lago de Maracaibo, near San Felipe.
Allotype and paratypes: 30. Same locality.
APLOCHEILIDAE
- Raja hollandi** Jordan and Richardson, 1909
Mem. Carnegie Mus., 4(4): 163.
Holotype: FMNH 52101 (CM 224). Taiwan: Takao.
RAJIDAE
- Rhamdella montana** Eigenmann, 1913
Ann. Carnegie Mus., 8(3-4): 421.
Holotype: FMNH 56067 (CM 4858). Peru: Queta, brook southeast of Tarma.
Paratype. Same locality.
PIMELODIDAE
- Rhamdia barbata** Meek, 1907
Field Mus. Nat. Hist. Zool. Ser., 7(4): 106.
Holotype: FMNH 5906. Nicaragua: lagoon at San Francisco.
Paratypes: 8. Nicaragua: 2 localities.
PIMELODIDAE
- Rhamdia branneri** Haseman, 1911
Ann. Carnegie Mus., 7(3-4): 377.
Holotype: FMNH 54235 (CM 2851). Brazil: creek near Serrinha Paraná.
Paratypes: 2. Same locality.
PIMELODIDAE
- Rhamdia branneri voulezi** Haseman, 1911
Ann. Carnegie Mus., 7(3-4): 378.
Holotype: FMNH 54238 (CM 2854). Brazil: Rio Iguaçu at Pôrto União da Victoria.
Paratypes: 2. Same locality.
PIMELODIDAE
- Rhamdia cabreræ** Meek, 1906
Field Mus. Nat. Hist. Zool. Ser., 7(3): 93.
Holotype: FMNH 5500. Guatemala: outlet of Lake Amatitlán.
Paratypes: 49. Same locality.
PIMELODIDAE
- Rhamdia laticauda typhla** Greenfield, Greenfield, and Woods, 1982
Brenesia, 19/20: 564.
Holotype: FMNH 71605. Belize: cave in Mountain Pine Ridge.
Paratypes: 12. Same locality.
PIMELODIDAE
- Rhamdia luigiana** Villa, 1977
Brenesia, 12/13: 133.
Paratypes: 4. Nicaragua: Lago Nicaragua.
Villa (1977) lists 2 FMNH paratypes without catalogue number; we have 4 specimens from the type locality that were labeled paratypes.
PIMELODIDAE
- Rhamdia microps** Eigenmann, 1917
Ann. Carnegie Mus., 11(3-4): 394.
Holotype: FMNH 58285 (CM 7283). Brazil: Uruguayana.
Paratypes: 3. Same locality.
PIMELODIDAE
- Rhamdia nasuta** Meek, 1909
Field Mus. Nat. Hist. Zool. Ser., 7(7): 207.
Holotype: FMNH 6480. Costa Rica: Buenos Aires de Térraba.
PIMELODIDAE
- Rhamdia oaxacæ** Meek, 1902
Field Mus. Nat. Hist. Zool. Ser., 3(6): 74.
Holotype: FMNH 3716. Mexico: Cuicatlán, Río Quitepec.
Paratypes: 9. Same locality.
PIMELODIDAE
- Rhamdia regani** Meek, 1907
Field Mus. Nat. Hist. Zool. Ser., 7(5): 144.
Holotype: FMNH 6019. Costa Rica: Turrialba.
Paratypes: 9. Costa Rica: 2 localities.
PIMELODIDAE

- Rhamdiopsis moreirai** Haseman, 1911
Ann. Carnegie Mus., 7(3–4): 375.
Holotype: FMNH 54372 (CM 2849). Brazil: Rio Iguaçu at Serrinha Paraná.
Paratype. Same locality.
PIMELODIDAE
- Rhaphiobelone robusta** Schultz, 1953
Bull. U.S. Natl. Mus., 202(1): 164.
Paratype. Philippine Islands: Luzon, Port Dupon.
BELONIDAE
- Rhegma bermudensis** Kanazawa, 1952
Fieldiana Zool., 34(7): 82.
Holotype: FMNH 48924. Bermuda: south side beaches.
Paratypes: 4. Bermuda and The Bahamas: 2 localities.
GRAMMISTIDAE
- Rhigophila dearborni** DeWitt, 1962
Copeia, 1962(4): 821.
Paratypes: 6. Antarctica: McMurdo Sound.
ZOARCIDAE
- Rhinodoras boehlkei** Glodek, Whitmire, and Orcés, 1976
Fieldiana Zool., 70(1): 3.
Holotype: FMNH 79203. Ecuador: Río Bobonaza.
DORADIDAE
- Rhinodoras thomersoni** Taphorn and Lilyestrom, 1984
Rev. UNELLEZ Cienc. Tec., 2: 87.
Paratypes: 3. Venezuela: mouth of Río Catatumbo, near El Congo.
DORADIDAE
- Rhinogobius formosanus** Oshima, 1919
Ann. Carnegie Mus., 12(2–4): 300.
Holotype: FMNH 59135 (CM 8273). Taiwan: Shinchiku.
GOBIIDAE
- Rhinogobius mowbrayi** Bean, 1906
Field Mus. Nat. Hist. Zool. Ser., 7(2): 81.
Holotype: FMNH 5030. Bermuda: Challenger Bank.
GOBIIDAE
- Rhinogobius taiwanus** Oshima, 1919
Ann. Carnegie Mus., 12(2–4): 298.
Holotype: FMNH 59134 (CM 8272). Taiwan: Shinchiku.
GOBIIDAE
- Rhinosardinia serrata** Eigenmann, 1912
Mem. Carnegie Mus., 5: 445.
Holotype: FMNH 53927 (CM 2443). Guyana: Morawhanna.
Paratypes: 215. Guyana: 2 localities.
CLUPEIDAE
- Rhizoprionodon oligolinx** Springer, 1964
Proc. U.S. Natl. Mus., 115(3493): 521.
Paratypes: 3. Singapore, Java and Sri Lanka: 3 localities.
CARCHARHINIDAE
- Rhoadsia altipinna** Fowler, 1911
Proc. Acad. Nat. Sci. Phila., 63: 498.
Paratype. Ecuador: tributary of Chimbo River.
CHARACIDAE
- Rhodeus chosenicus** Jordan and Metz, 1913
Mem. Carnegie Mus., 6(1): 19.
Holotype: FMNH 55804 (CM 4567). Korea: Suigen.
Paratypes: 2. Same locality.
CYPRINIDAE
- Rhodeus kurumeus** Jordan and Thompson, 1914
Mem. Carnegie Mus., 6(4): 229.
Paratype. Japan: Chikugo River at Kurume.
CYPRINIDAE
- Rhombatractus affinis** Weber, 1908
Nova Guinea, 5(2): 234.
Syntypes: 2, FMNH 52394 (CM 650). New Guinea: Timena River.
MELANOTAENIIDAE
- Rhombatractus sentaniensis** Weber, 1908
Nova Guinea, 5(2): 235.
Syntypes: 4, FMNH 52393 (CM 649). New Guinea: Lake Sentani.
MELANOTAENIIDAE
- Rhombosoma sepikensis** Herre, 1935
Field Mus. Nat. Hist. Zool. Ser., 18(12): 400.
Holotype: FMNH 17265. New Guinea: Sepik River at Marienberg.
Paratypes: 6. Same locality.
MELANOTAENIIDAE
- Rhynchodoras woodsi** Glodek, 1976
Copeia, 1976(1): 44.
Holotype: FMNH 77008. Ecuador: Río Bobonaza at Moreta.
Paratypes: 2 MISSING. Same locality.
DORADIDAE

Rineloricaria formosa Isbrucker and Nijssen, 1979
Bijdr. Dierkd., 48(2): 192.
Holotype: FMNH 83713. Colombia: Río Orinoco basin, lagoon near Puerto Inírida.
Paratypes: 30. Colombia: 2 localities.
LORICARIIDAE

Rivulus breviceps Eigenmann, 1909
Ann. Carnegie Mus., 6(1): 49.
Holotype: FMNH 52710 (CM 1075). Guyana: Shrimp Creek.
Paratype: MISSING. Same locality.
APLOCHEILIDAE

Rivulus brunneus Meek and Hildebrand, 1913
Field Mus. Nat. Hist. Zool. Ser., 10(8): 86.
Holotype: FMNH 7593. Panama: Canal Zone, Creek at Toro Point.
Paratypes: 7. Same locality.
APLOCHEILIDAE

Rivulus compressus Henn, 1916
Ann. Carnegie Mus., 10(1-2): 111.
Holotype: FMNH 57003 (CM 5819). Brazil: Manaus.
APLOCHEILIDAE

Rivulus frenatus Eigenmann, 1912
Ann. Carnegie Mus., 5: 51.
Holotype: FMNH 53538 (CM 1085). Guyana: Essequibo River at Gluck Island.
APLOCHEILIDAE

Rivulus holmiae Eigenmann, 1909
Ann. Carnegie Mus., 6(1): 50.
Holotype: FMNH 53535 (CM 1076). Guyana: creek near Holmia.
Paratypes: 5. Same locality.
APLOCHEILIDAE

Rivulus lanceolatus Eigenmann, 1909
Ann. Carnegie Mus., 6(1): 51.
Holotype: FMNH 53537 (CM 1084). Guyana: Rockstone.
APLOCHEILIDAE

Rivulus magdalenae Eigenmann and Henn, 1916
In: Henn, Ann. Carnegie Mus., 10(1-2): 109.
Holotype: FMNH 56997 (CM 5813). Colombia: Ibagué.
Paratypes: 32. Colombia: 5 localities.
APLOCHEILIDAE

Rivulus montium Hildebrand, 1938
Field Mus. Nat. Hist. Zool. Ser., 22(4): 319.
Paratypes: 2. Panama: between Río Boquerón and Río Pequeni.
APLOCHEILIDAE

Rivulus stagnatus Eigenmann, 1909
Ann. Carnegie Mus., 6(1): 50.
Holotype: FMNH 52715 (CM 1082). Guyana: Christianburg.
Paratypes: 4. Same locality.
APLOCHEILIDAE

Rivulus stellifer Thomerson and Turner, 1973
Copeia, 1973(4): 783.
Paratypes: 5. Venezuela: Cojedes near Caño Benito.
APLOCHEILIDAE

Rivulus volcanus Hildebrand, 1938
Field Mus. Nat. Hist. Zool. Ser., 22(4): 316.
Paratypes: 10. Panama: Laguna Grande (Davis), Chiriquí.
APLOCHEILIDAE

Rivulus waimacui Eigenmann, 1909
Ann. Carnegie Mus., 6(1): 50.
Holotype: FMNH 53536 (CM 1078). Guyana: Shrimp Creek.
Paratypes: 5. Guyana: 3 localities.
APLOCHEILIDAE

Roeboides cauae Eigenmann, 1922
Mem. Carnegie Mus., 9(1): 162.
Paratypes: 298. Colombia: 3 localities.
CHARACIDAE

Roeboides hildebrandi Eigenmann, 1922
Mem. Carnegie Mus., 9(1): 162.
Holotype: FMNH 56884 (CM 5186). Colombia: Istmina.
Paratypes: 44. Colombia: 6 localities.
CHARACIDAE

Roeboides macrolepis Meek and Hildebrand, 1913
Field Mus. Nat. Hist. Zool. Ser., 10(8): 84.
Holotype: FMNH 7590. Panama: Río Cupe, Boca de Cupe.
Paratypes: 10. Panama: 2 localities.
CHARACIDAE

Rueboides magdalenae Eigenmann, 1922

Mem. Carnegie Mus., 9(1): 161.

Syntypes: 201, FMNH 56386–56395 (CM 5202–5210). Colombia: Soplaviento.

Eigenmann (1922, p. 161) indicated that CM 5202 and IUM 12950 (now at California Academy of Sciences) were the “type and paratypes” but apparently never separated, labeled or figured the holotype.

CHARACIDAE

Rueboides meeki Eigenmann, 1922

Mem. Carnegie Mus., 9(1): 163.

Holotype: FMNH 55149 (CM 3849). Colombia: Quibdó.

Paratypes: 59. Colombia: 3 localities.

CHARACIDAE

Rueboides occidentalis Meek and Hildebrand, 1916

Field Mus. Nat. Hist. Zool. Ser., 10(15): 293.

Holotype: FMNH 8948 MISSING. Panama: Canal Zone, Río Cárdenas at Corozal.

Paratypes: 223. Panama: 21 localities.

CHARACIDAE

Rueboides salvadoris Hildebrand, 1925

Bull. Bur. Fish., 41: 246.

Paratypes: 21. El Salvador: 3 localities.

CHARACIDAE

Rueboides thurni Eigenmann, 1912

Mem. Carnegie Mus., 5: 399.

Holotype: FMNH 53676 (CM 2149). Guyana: Rockstone sandbank.

Paratypes: 7. Guyana: 2 localities.

CHARACIDAE

Sagamia russula Jordan and Snyder, 1901

Proc. U.S. Natl. Mus., 24(1244): 100.

Paratypes: 10. Japan. Nagasaki.

GOBIIDAE

Salarias martini Herre, 1942

Proc. Biol. Soc. Wash., 55: 2.

Paratypes: 7. Philippine Islands: Panay, Estancia.

BLENNIIDAE

Salarias walensis Herre, 1935

Field Mus. Nat. Hist. Zool. Ser., 18(12): 437.

Holotype: FMNH 17393. New Hebrides Islands: Wala Island.

Paratype. Same locality.

BLENNIIDAE

Salmo bathoecetor Meek, 1899

Field Mus. Nat. Hist. Zool. Ser., 1(12): 227.

Holotype: FMNH 2035. Washington: Lake Crescent.

Paratype. Same locality.

SALMONIDAE

Salmo clarki declivifrons Meek, 1899

Field Mus. Nat. Hist. Zool. Ser., 1(12): 230.

Holotype: FMNH 2006. Washington: Lake Southerland.

SALMONIDAE

Salmo clarki jordani Meek, 1899

Field Mus. Nat. Hist. Zool. Ser., 1(12): 229.

Holotype: FMNH 2012. Washington: Lake Southerland.

Paratypes: 18. Same locality.

SALMONIDAE

Salmo roosevelti Evermann, 1906

Bull. Bur. Fish. 1905, 25: 26.

Paratype. California: Volcáno Creek.

SALMONIDAE

Salvelinus imbrius Jordan and McGregor, 1925

In: Jordan and Hubbs, Mem. Carnegie Mus., 10(2): 142.

Holotype: FMNH 58692 (CM 7797). Japan: Iwami, stream near Hamada.

SALMONIDAE

Salvelinus timagamiensis Henn and Rinkenbach, 1925

Ann. Carnegie Mus., 16: 131.

Holotype: FMNH 70514 (CM 7969). Canada: Ontario, White Pine Lake.

Paratypes: 11. Same locality.

SALMONIDAE

Sarcocheilichthys morii Jordan and Hubbs, 1925

Mem. Carnegie Mus., 10(2): 175.

Holotype: FMNH 58718 (CM 7824). Korea: Ping-yang River.

CYPRINIDAE

Sardinella marquesensis Berry and Whitehead, 1968

Proc. Biol. Soc. Wash., 81: 209.

Paratypes: 2. Marquesas Islands: 2 localities.

CLUPEIDAE

Sargocentron macrosquamis Golani, 1984

Cybius, 8(2): 40.

Holotype: FMNH 86961. Amirantes Islands: eastern end of D'Arros Island.

HOLOCENTRIDAE

- Scaphesthes tamusuiensis** Oshima, 1919
Ann. Carnegie Mus., 12(2-4): 209.
Holotype: FMNH 59091 (CM 8228). Taiwan:
Tamusui River near Shinten.
CYPRINIDAE
- Scarus laui** Jordan and Evermann, 1903
Bull. U.S. Fish Comm. 1902, 22: 196.
Paratype. Hawaiian Islands: Honolulu.
SCARIDAE
- Scepterias fragilis** Jordan and Jordan, 1922
Mem. Carnegie Mus., 10(1): 45.
Holotype: FMNH 55204 (CM 3900). Hawaiian
Islands: Honolulu Market.
APOGONIDAE
- Schizodontopsis laticeps** Eigenmann, 1912
Mem. Carnegie Mus., 5: 229.
Holotype: FMNH 53359 (CM 1825). Guyana:
Crab Falls.
ANOSTOMIDAE
- Sciaena bathytatos** Chao and Miller, 1975
Bull. Mar. Sci., 25(2): 267.
Paratype. Caribbean Sea: off Venezuela.
SCIAENIDAE
- Sciaena iharae** Jordan and Metz, 1913
Mem. Carnegie Mus., 6(1): 37.
Holotype: FMNH 55629 (CM 4371). Korea:
Fusan.
SCIAENIDAE
- Sciaena trewavasae** Chao and Miller, 1975
Bull. Mar. Sci., 25(2): 262.
Paratype. Caribbean Sea: off Venezuela.
SCIAENIDAE
- Scleronema operculatum** Eigenmann, 1917
Proc. Am. Philos. Soc., 56(8): 691.
Holotype: FMNH 58080 (CM 7077). Brazil:
Cacequy.
Paratypes: 3. Same Locality.
TRICHOMYCTERIDAE
- Scolopsis eriomma** Jordan and Richardson, 1909
Mem. Carnegie Mus., 4(4): 188.
Holotype: FMNH 52247 (CM 317). Taiwan:
Takao.
NEMIPTERIDAE
- Scomberoides formosanus** Wakiya, 1924
Ann. Carnegie Mus., 15(2-3): 236.
Holotype: FMNH 59499 (CM 7765). Japan: Kii.
CARANGIDAE
- Scopelarchus beebe** Rofen, 1963
Aquatica, Aquatic Res. Inst. Stockton, 3: 3.
Paratype. Western North Atlantic: off Bermuda.
SCOPELARCHIDAE
- Scopelarchus sagax** Rofen, 1963
Aquatica, Aquatic Res. Inst. Stockton, 3: 2.
Holotype: FMNH 49852. Western North At-
lantic: off Bermuda.
SCOPELARCHIDAE
- Scopelengys clarkei** Butler and Ahlstrom, 1976
Fish. Bull., 74(1): 148.
Paratype. Central Pacific.
NEOSCOPELIDAE
- Scorpaenopsis catocala** Jordan and Evermann,
1903
Bull. U.S. Fish Comm. 1902, 22: 201.
Paratype. Hawaiian Islands: Hilo.
SCORPAENIDAE
- Sebastapistes badiorufus** Herre, 1935
Field Mus. Nat. Hist. Zool. Ser., 18(12): 409.
Holotype: FMNH 17337. Tuamotu Islands:
Takaroa Island.
SCORPAENIDAE
- Sebastodes clavilatus** Starks, 1911
Ann. Carnegie Mus., 7(2): 181.
Holotype: FMNH 52293 (CM 548). Washing-
ton: San Juan Islands.
SCORPAENIDAE
- Sebastodes deani** Starks, 1911
Ann. Carnegie Mus., 7(2): 178.
Holotype: FMNH 52290 (CM 545). Washing-
ton: San Juan Islands.
SCORPAENIDAE
- Sebastodes emphaeus** Starks, 1911
Ann. Carnegie Mus., 7(2): 182.
Holotype: FMNH 52294 (CM 549). Washing-
ton: San Juan Islands.
Paratypes: 5. Same locality.
SCORPAENIDAE

Sebastodes ijimae Jordan and Metz, 1913

Mem. Carnegie Mus., 6(1): 49.

Holotype: FMNH 55437 (CM 4172). Korea: Fusan.

SCORPAENIDAE

Sebastodes owstoni Jordan and Thompson, 1914

Mem. Carnegie Mus., 6(4): 270.

Holotype: FMNH 57088 (CM 6026). Japan: Aomori.

Paratypes: 3. Same locality.

SCORPAENIDAE

Sebastodes thompsoni Jordan and Hubbs, 1925

Mem. Carnegie Mus., 10(2): 265.

Paratype: Japan: Osaka.

SCORPAENIDAE

Serraniculus pumilio Ginsburg, 1952

J. Wash. Acad. Sci., 42(3): 88.

Paratype. Gulf of Mexico: off Texas.

SERRANIDAE

Serranus chionaraia Robins and Starck, 1961

Proc. Acad. Nat. Sci. Phila., 113(11): 270.

Paratype. Western Caribbean Sea: off Honduras.

SERRANIDAE

Serranus maytagi Robins and Starck, 1961

Proc. Acad. Nat. Sci. Phila., 113(11): 288.

Paratype. Western Caribbean Sea: off Honduras.

SERRANIDAE

Serrasalmo hollandi Eigenmann, 1915

Ann. Carnegie Mus., 9(3–4): 251.

Holotype: FMNH 56978 (CM 5792). Brazil: Rio Guaporé at Maciel.

CHARACIDAE

Serrasalmo humeralis gracilior Eigenmann, 1915

Ann. Carnegie Mus., 9(3–4): 257.

Syntypes: 3, FMNH 56977 (CM 5791). Brazil: Rio Guaporé at Maciel.

CHARACIDAE

Serrasalmus coccogenis Fowler, 1911

Proc. Acad. Nat. Sci. Phila., 63: 428.

Paratype. Venezuela: Caño Uracoa at La Pedrita.

CHARACIDAE

Sicydium altum Meek, 1907

Field Mus. Nat. Hist. Zool. Ser., 7(5): 149.

Holotype: FMNH 6034. Costa Rica: Turrialba.

Paratypes: 2. Costa Rica: El Guayabo.

GOBIIDAE

Sicydium hildebrandi Eigenmann, 1917

Proc. Am. Philos. Soc., 56(7): 685.

Holotype: FMNH 58465 (CM 7466). Colombia:

Río Dagua at Cisnero.

GOBIIDAE

Sicydium montanum Hubbs, 1920

Proc. Biol. Soc. Wash., 33: 89.

Holotype: FMNH 9053. Venezuela: Caracas, brook at Macuto.

Paratypes: 10. Same locality.

GOBIIDAE

Signalosa atchafalayae vanhyningi Weed, 1925

Field Mus. Nat. Hist. Zool. Ser., 12(11): 145.

Holotype: FMNH 11395. Florida: near Gainesville.

Paratypes: 3. Same locality.

CLUPEIDAE

Signalosa mexicana campi Weed, 1925

Field Mus. Nat. Hist. Zool. Ser., 12(11): 143.

Holotype: FMNH 11413. Texas: Brownsville, Resaca de la Guerra.

Paratypes: 493. Texas: 4 localities.

CLUPEIDAE

Siphamia cuprea Lachner, 1953

Bull. U.S. Natl. Mus., 202(1): 423.

Paratypes: 2. Sulu Archipelago: Pangasinan Island.

APOGONIDAE

Siphamia fuscolineata Lachner, 1953

Bull. U.S. Natl. Mus., 202(1): 425.

Paratypes: 2. Bikini Island.

APOGONIDAE

Siphostoma brevicaudum Meek, 1904

Field Mus. Nat. Hist. Zool. Ser., 5: 163.

Holotype: FMNH 4586. Mexico: Veracruz, Boca del Río.

Paratypes: 2. Same locality.

SYNGNATHIDAE

Siphostoma eigenmanni Wilson, 1916

Ann. Carnegie Mus., 10(1–2): 60.

Holotype: FMNH 56857 (CM 5672). Brazil: Bahia, Rio Vermelho.

SYNGNATHIDAE

Siphostoma elcapitanense Meek and Hildebrand, 1914

Field Mus. Nat. Hist. Zool. Ser., 10(10): 119.
Paratypes: 4. Panama and Costa Rica: 3 localities.

SYNGNATHIDAE

Skiffia lermac Meek, 1902

Field Mus. Nat. Hist. Zool. Ser., 3(6): 102.
Holotype: FMNH 3616. Mexico: Lago de Patzcuaro at Patzcuaro.

Paratypes: 131. Mexico: 2 localities.

GOODEIDAE

Skiffia variegata Meek, 1902

Field Mus. Nat. Hist. Zool. Ser., 3(6): 104.
Holotype: FMNH 3612. Mexico: Lago de Zirahuén at Zirahuén.

Allotype and paratypes: 99. Mexico: 2 localities.
GOODEIDAE

Solenostomus paegnius Jordan and Thompson, 1914

Mem. Carnegie Mus., 6(4): 235.

Holotype: FMNH 57083 (CM 6020). Japan: Misaki.

SOLENOSTOMIDAE

Soleonassus finis Eigenmann, 1912

Mem. Carnegie Mus., 5: 528.

Holotype: FMNH 53965 (CM 2487). Guyana: Tumatumari.

SOLEIDAE

Solivomer arenidens Miller, 1947

Proc. U.S. Natl. Mus., 97(3211): 84.

Paratypes: 2. Philippine Islands: off Balicasag Island.

NEOSCOPELIDAE

Sonoda megalophthalma Grey, 1959

Bull. Mus. Comp. Zool. Harv. Coll., 121(4): 181.

Holotype: FMNH 65981. Western Caribbean Sea.

Paratypes: 22. Western Caribbean Sea: 5 localities.

STERNOPTYCHIDAE

Sovichthys abuelo Schultz, 1944

Proc. U.S. Natl. Mus., 94(3172): 191.

Paratypes: 4. Venezuela: Río Socuy, above its mouth.

PIMELODIDAE

Sparisoma rhomaleum Meek and Hildebrand, 1928

Field Mus. Nat. Hist. Zool. Ser., 15(3): 754.

Paratype. Panama: Colón market.

SCARIDAE

Spectrunculus radcliffei Jordan and Thompson, 1914

Mem. Carnegie Mus., 6(4): 301.

Holotype: FMNH 57123 (CM 6061). Japan: Misaki.

OPHIDIIDAE

Sphagebranchus conklini Eigenmann, 1916

Ann. Carnegie Mus., 10(1-2): 55.

Holotype: FMNH 57731 (CM 6710). The Bahamas: New Providence.

OPHICHTHIDAE

Spheroides asterias Blosser, 1909

Ann. Carnegie Mus., 6(1): 300.

Holotype: FMNH 53050 (CM 1475). Lesser Antilles: Saint Croix Island.

TETRAODONTIDAE

Sphoeroides kendalli Meek and Hildebrand, 1928

Field Mus. Nat. Hist. Zool. Ser., 15(3): 819.

Paratype. Panama: Chame Point.

TETRAODONTIDAE

Sphoeroides parvus Shipp and Yerger, 1969

Proc. Biol. Soc. Wash., 82: 478.

Paratypes: 10. Alabama: Mobile Bay.

TETRAODONTIDAE

Spinibarbus hollandi Oshima, 1919

Ann. Carnegie Mus., 12(2-4): 218.

Holotype: FMNH 59095 (CM 8233). Taiwan: Sobun River near Tabani.

CYPRINIDAE

Spinoblennius spiniger Herre, 1935

Field Mus. Nat. Hist. Zool. Ser., 18(12): 435.

Holotype: FMNH 17395. Ecuador: Islas Galápagos, Isla Edén.

Paratype. Same locality.

BLENNIIDAE

Spintherobolus broccae Myers, 1925

Ann. Carnegie Mus., 16: 143.

Holotype: FMNH 58863 (CM 7979). Brazil: hills behind Rio de Janeiro.

Paratype. Same locality.

CHARACIDAE

Spintherobolus papilliferus Eigenmann, 1911
Ann. Carnegie Mus., 8(1): 167.
Holotype: (CM 3582) MISSING. Brazil: São Paulo, Alto da Serra.
Paratypes: 5. Same locality.
CHARACIDAE

Spirinchus dilatus Schultz and Chapman, 1934
Ann. Mag. Nat. Hist., Ser. 10, 13: 68.
Paratype. Washington: Nooksack River.
OSMERIDAE

Spirinchus verecundus Jordan and Metz, 1913
Mem. Carnegie Mus., 6(1): 11.
Holotype: FMNH 55807 (CM 4570). Korea: Chinnampo.
Paratype. Same locality.
OSMERIDAE

Squaliolus sarmenti Noronha, 1926
Ann. Carnegie Mus., 16(3–4): 385.
Holotype: FMNH 58862 (CM 7976). Portugal: Madeira Islands, Funchal.
SQUALIDAE

Squalogadus intermedius Grey, 1959
Fieldiana Zool., 39(29): 330.
Paratypes: 5. Gulf of Mexico.
MACROURIDAE

Starksia occidentalis Greenfield, 1979
Fieldiana Zool., 73(2): 27.
Holotype: FMNH 83716. Belize: Tarpon Cay.
Paratypes: 41. Belize and Honduras: 11 localities.
LABRISOMIDAE

Starksia variabilis Greenfield, 1979
Fieldiana Zool., 73(2): 33.
Holotype: FMNH 83717. Colombia: Santa Marta.
Paratypes: 7. Same locality.
LABRISOMIDAE

Starksia y-lineata Gilbert, 1965
Not. Nat. (Phila.), 379: 2.
Paratypes: 2. British West Indies: Grand Cayman Island.
LABRISOMIDAE

Stegastes baldwini Allen and Woods, 1980
Rec. West. Aust. Mus., 8(2): 183.
Paratypes: 54. Clipperton Island.
POMACENTRIDAE

Stellifer colonensis Meek and Hildebrand, 1925
Field Mus. Nat. Hist. Zool. Ser., 15(2): 623.
Paratypes: 21. Panama: 4 localities.
SCIAENIDAE

Stellifer melanocheir Eigenmann, 1917
Proc. Am. Philos. Soc., 56(7): 682.
Holotype: FMNH 58507 (CM 7520). Colombia: Tumaco.
SCIAENIDAE

Stellifer mindii Meek and Hildebrand, 1925
Field Mus. Nat. Hist. Zool. Ser., 15(2): 626.
Paratypes: 3. Panama: Canal Zone near Mindi Reef.
SCIAENIDAE

Stellistius katsukii Jordan and Tanaka, 1927
Ann. Carnegie Mus., 17(12): 389.
Holotype: FMNH 59186 (CM 8329). Japan: Hokkaido, off Mororan.
HEXAGRAMMIDAE

Sternarchus hasemani Ellis, 1913
Mem. Carnegie Mus., 6(3): 147.
Holotype: FMNH 54562 (CM 3191). Brazil: near Santarém.
Paratypes: 15. Same locality.
APTERONOTIDAE

Sternarchus leptorhynchus Ellis, 1912
In: Eigenmann, Mem. Carnegie Mus., 5: 439.
Holotype: FMNH 53294 (CM 1762). Guyana: Amatuk.
Paratype: Guyana: Warraputa.
Published number CM 1763 for the holotype is an error.
APTERONOTIDAE

Sternarchus mariae Eigenmann and Fisher, 1914
Indiana Univ. Stud., 25: 236.
Holotype: FMNH 56774 (CM 5594). Colombia: Girardot.
APTERONOTIDAE

Sternarchus rostratus Meek and Hildebrand, 1913
Field Mus. Nat. Hist. Zool. Ser., 10(8): 85.
Holotype: FMNH 7592. Panama: Río Grande, Cana.
APTERONOTIDAE

Sternopygus dariensis Meek and Hildebrand, 1916
Field Mus. Nat. Hist. Zool. Ser., 10(15): 309.
Holotype: FMNH 8949. Panama: Río Tuyra at Marriganti.
Paratypes: 12. Same locality.
STERNOPYGIDAE

- Sternopygus pejeraton** Schultz, 1949
Proc. U.S. Natl. Mus., 99(3235): 60.
Paratypes: 4. Venezuela: Lago Maracaibo basin,
Río Apón.
STERNOPYGIDAE
- Stethaprion crenatus** Eigenmann, 1916
Ann. Carnegie Mus., 10(1-2): 80.
Holotype: FMNH 56413 (CM 5228). Bolivia:
San Joaquin.
Paratypes: 3. Bolivia and Brazil: 2 localities.
CHARACIDAE
- Stevardia aliata** Eigenmann, 1914
Indiana Univ. Stud., 20: 37.
Holotype: FMNH 56400 (CM 5215). Colombia:
Río Negro at Villavicencio.
Paratypes: 13. Colombia: 2 localities.
CHARACIDAE
- Stizostedion glaucum** Hubbs, 1926
Misc. Publ. Mus. Zool. Univ. Mich., 15: 58.
Paratypes: 2. Pennsylvania: Lake Erie off Erie.
PERCIDAE
- Stolephorus branchiomelas** Eigenmann, 1917
Proc. Am. Philos. Soc., 56(7): 682.
Holotype: FMNH 58489 (CM 7491). Colombia:
mouth of Río Dagua.
Paratypes: 6. Colombia: 2 localities.
ENGRAULIDAE
- Stolephorus guianensis** Eigenmann, 1912
Mem. Carnegie Mus., 5: 447.
Holotype: FMNH 53932 (CM 2448). Guyana:
Bartica.
Paratype: Guyana: below Wismar.
ENGRAULIDAE
- Stolephorus lyolepis** Evermann and Marsh, 1902
Bull. U.S. Fish Comm. 1900, 20(1): 89.
Paratypes: 5. Puerto Rico: Culebra.
ENGRAULIDAE
- Suruga fundicola** Jordan and Snyder, 1901
Proc. U.S. Natl. Mus., 24(1244): 96.
Paratypes: 4. Japan: Suruga Bay.
GOBIIDAE
- Symphysanodon berryi** Anderson, 1970
U.S. Fish Wildl. Serv. Fish. Bull., 68(2): 338.
Paratypes: 20. Caribbean Sea: 5 localities.
SERRANIDAE
- Symphysanodon octoactinus** Anderson, 1970
U.S. Fish Wildl. Serv. Fish. Bull., 68(2): 342.
Paratype: Western Caribbean Sea.
SERRANIDAE
- Synaphobranchus oregoni** Castle, 1960
Fieldiana Zool., 39(35): 388.
Paratypes: 6. Gulf of Mexico.
SYNAPHOBRANCHIDAE
- Synchiropus ijimae** Jordan and Thompson, 1914
Mem. Carnegie Mus., 6(4): 295.
Holotype: FMNH 57078 (CM 6015). Japan:
Misaki.
CALLIONYMIDAE
- Synchiropus minutulus** Fricke, 1981
Theses Zoologicae, 1: 119.
Holotype: FMNH 78973. Maldives Archipelago:
southwest of Mafilifleuri.
CALLIONYMIDAE
- Syngnathus mackayi nesiotes** Herald, 1942
Stanford Ichthyol. Bull., 2(4): 128.
Paratypes: 10. Bermuda: 7 localities.
SYNGNATHIDAE
- Syngnathus scovelli makaxi** Herald and Dawson, 1972
Copeia, 1972(4): 781.
Paratypes: 4. Mexico: Isla Mujeres, Laguna
Makax.
SYNGNATHIDAE
- Synodus binotatus** Schultz, 1953
Bull. U.S. Natl. Mus., 202(1): 35.
Paratypes: 2. Marshall Islands: 2 localities.
SYNODONTIDAE
- Synodus ulae** Schultz, 1953
Bull. U.S. Natl. Mus., 202(1): 38.
Paratype: Hawaiian Islands: Honolulu Market.
SYNODONTIDAE
- Taenioides snyderi** Jordan and Hubbs, 1925
Mem. Carnegie Mus., 10(2): 310.
Holotype: FMNH 58831 (CM 7941). Japan:
Wakanoura.
GOBIOIDIDAE
- Tagusa delicata** Herre, 1935
Field Mus. Nat. Hist. Zool. Ser., 18(12): 435.
Holotype: FMNH 17399. Ecuador: Islas Galápagos,
Isla Albemarle, Tagus Cove.
Paratypes: 3. Same locality.
BLENNIIDAE

- Terapon habbemai** Weber, 1910
Notes Leiden Mus., 32: 234.
Syntype: FMNH 52496 (CM 754). New Guinea:
Lorentz River, Alkmaar.
Spelled *Therapon* in Weber (1910).
TERAPONIDAE
- Tetragonopterus alleni** Eigenmann and McAtee, 1907
In: Eigenmann, McAtee, and Ward, Ann.
Carnegie Mus., 4(2): 126.
Paratype. Paraguay: Corumbá.
CHARACIDAE
- Tetraodon kretamensis** Inger, 1953
Fieldiana Zool., 34(11): 149.
Holotype: FMNH 51558. Malaysia: Sabah, Pi-
nang River.
Paratypes: 56. Malaysia: Sabah, 4 localities.
TETRAODONTIDAE
- Teuthis atrimentatus** Jordan and Evermann, 1903
Bull. U.S. Fish Comm. 1902, 22: 198.
Paratype. Hawaiian Islands: Honolulu.
ACANTHURIDAE
- Thalassoma cranei** Herre, 1935
Field Mus. Nat. Hist. Zool. Ser., 18(12): 412.
Holotype: FMNH 17347. Solomon Islands: Is-
abel Island, Tenibuli.
LABRIDAE
- Thalassophryne quadrizonatus** Eigenmann, 1922
Mem. Carnegie Mus., 9(1): 217.
Holotype: (CM 3921) MISSING. Colombia: Río
Truandó.
BATRACHOIDIDAE
- Thoracocharax brevis** Eigenmann, 1912
Indiana Univ. Stud., 16: 25.
Holotype: FMNH 56055 (CM 4845). Colombia:
Raspadura.
GASTEROPELECIDAE
- Thoracocharax magdalenae** Eigenmann, 1912
Indiana Univ. Stud., 16: 25.
Holotype: FMNH 56056 (CM 4846). Colombia:
Girardot.
GASTEROPELECIDAE
- Thorichthys ellioti** Meek, 1904
Field Mus. Nat. Hist. Zool. Ser., 5: 223.
Holotype: FMNH 4627. Mexico: Veracruz,
Motzorongo.
Paratypes: 24. Same locality.
CICHLIDAE
- Thyrina colombiensis** Hubbs, 1920
Occas. Pap. Mus. Zool. Univ. Mich., 88: 4.
Paratypes: 49. Colombia: 3 localities.
ATHERINIDAE
- Thyrina guija** Hildebrand, 1925
Bull. Bur. Fish., 41: 264.
Paratypes: 13. El Salvador: 3 localities.
ATHERINIDAE
- Thyrina meeki** Miller, 1907
Bull. Am. Mus. Nat. Hist., 23(2): 110.
Paratypes: 23. Guatemala: 5 localities.
ATHERINIDAE
- Thysanichthys evides** Jordan and Thompson, 1914
Mem. Carnegie Mus., 6(4): 272.
Holotype: FMNH 57082 (CM 6019). Japan:
Misaki.
SCORPAENIDAE
- Tomeurus gracilis** Eigenmann, 1909
Ann. Carnegie Mus., 6(1): 53.
Holotype: FMNH 53541 (CM 1093). Guyana:
Mud Creek in Aruka River.
Paratypes: 2. Guyana: 2 localities.
POECILIIDAE
- Toxus riddlei** Eigenmann, 1903
Bull. U.S. Fish Comm. 1902, 22: 226.
Allotype: Cuba: San Cristóbal.
POECILIIDAE
- Trachelyichthys exilis** Greenfield and Glodek, 1977
Fieldiana Zool., 72(3): 49.
Holotype: FMNH 80476. Peru: tributary of Río
Nanay.
Paratype. Same locality.
AUCHENIPTERIDAE
- Trachinotus jordani** Wakiya, 1924
Ann. Carnegie Mus., 15(2-3): 221.
Holotype: FMNH 59374 (CM 7756). Japan:
Bonin Islands.
Spelled *Trachynotus* in Wakiya (1924).
CARANGIDAE
- Trachipterus fukuzakii** Fitch, 1964
Calif. Fish Game, 50(4): 236.
Paratype. Mexico: Baja California, southwest of
Cabo San Lucas.
TRACHIPTERIDAE

- Trachurus argenteus** Wakiya, 1924
Ann. Carnegie Mus., 15(2-3): 145.
Paratype. Japan: Amakusa, near Nagasaki.
CARANGIDAE
- Trachycorystes fisheri** Eigenmann, 1916
Ann. Carnegie Mus., 10(1-2): 82.
Holotype: FMNH 57695 (CM 6667). Colombia:
Río Sucio.
Paratypes: 16. Colombia: 3 localities.
CARANGIDAE
- Triathalassothia gloverensis** Greenfield and
Greenfield, 1973
Copeia, 1973(3): 560.
Holotype: FMNH 71575. Belize: Glover's Reef.
Paratypes: 7. Belize: 5 localities.
BATRACHOIDIDAE
- Trichopsetta caribbaea** Anderson and Guthertz,
1967
Bull. Mar. Sci., 17(4): 898.
Paratypes: 2. Caribbean Sea: off Colombia.
BOTHIDAE
- Tridentiger kuroiwaie** Jordan and Tanaka, 1927
Ann. Carnegie Mus., 17(2): 276.
Holotype: FMNH 59181 (CM 8324). Japan:
Amami-Oshima.
Paratypes: 9. Same locality.
GOBIIDAE
- Tylobronchus maculosus** Eigenmann, 1912
Mem. Carnegie Mus., 5: 272.
Holotype: FMNH 53449 (CM 1923). Guyana:
creek below Potaro Landing.
Paratypes: 54. Guyana: 6 localities.
CHILODONTIDAE
- Tympanopleura piperata** Eigenmann, 1912
Mem. Carnegie Mus., 5: 203.
Holotype: FMNH 53243 (CM 1708). Guyana:
Crab Falls.
Paratype. Same locality.
AGENEIOSIDAE
- Ulua richardsoni** Jordan and Snyder, 1909
Mem. Carnegie Mus., 4(2): 39.
Holotype: FMNH 55365 (CM 413). Taiwan:
Takao.
CARANGIDAE
- Umbrina tumacoensis** Wilson, 1916
Ann. Carnegie Mus., 10(1-2): 67.
Holotype: FMNH 56840 (CM 5654) MISSING.
Colombia: Tumaco.
Paratypes: 2. Same locality.
SCIAENIDAE
- Upeneus arge** Jordan and Evermann, 1903
Bull. U.S. Fish Comm. 1902, 22: 187.
Paratype. Hawaiian Islands: Honolulu.
MULLIDAE
- Uropterygius fuscoguttatus** Schultz, 1953
Bull. U.S. Natl. Mus., 202(1): 156.
Paratype. Marshall Islands: Bikini Atoll, Eman
Island.
MURAENIDAE
- Vaimosa balteata** Herre, 1935
Field Mus. Nat. Hist. Zool. Ser., 18(12): 419.
Holotype: FMNH 17386. Indonesia: Waigiu Is-
land, Majalibit Inlet.
GOBIIDAE
- Vaimosa horiae** Herre, 1936
Philipp. J. Sci., 59: 280.
Syntypes (?): 2, FMNH 47241. Palau Islands:
freshwater creek on Bab-el-Thaup Island.
These are from the type locality, but were not
explicitly cited in the original descrip-
tion; see comment in Böhlke (1953, Stan-
ford Ichthyol. Bull., 5(1): 119).
GOBIIDAE
- Vaimosa osgoodi** Herre, 1935
Field Mus. Nat. Hist. Zool. Ser., 18(12): 420.
Holotype: FMNH 17387. Fiji Islands: Viti Levu
Island, Suva Harbor.
GOBIIDAE
- Vandellia hasemani** Eigenmann, 1918
Mem. Carnegie Mus., 7(5): 363.
Holotype: FMNH 58523 (CM 7542). Bolivia:
Río Mamoré.
Paratypes: 2. Same locality.
TRICHOMYCTERIDAE
- Vandellia sanguinea** Eigenmann, 1917
Proc. Am. Philos. Soc., 56(8): 701.
Holotype: FMNH 58086 (CM 7082). Brazil:
São Antonio de Rio Madeira.
TRICHOMYCTERIDAE

Varicorhinus robustus Nichols, 1925

Am. Mus. Nov., 182: 2.

Paratypes: 2. China: Fukien, Yenping.

CYPRINIDAE

Varicus bucca Robins and Böhlke, 1961

Copeia, 1961(1): 47.

Paratype. Antilles Islands: north of Culebra Island.

GOBIIDAE

Varicus imswe Greenfield, 1981

Copeia, 1981(2): 269.

Holotype: FMNH 83894. Belize: Carrie-Bow Cay.

Paratype. Same locality.

GOBIIDAE

Vesicatrus tegatus Eigenmann, 1911

Ann. Carnegie Mus., 8(1): 174.

Holotype: (CM 3201) MISSING. Brazil: Upper Rio Paraguay basin, Jaurú.

Paratypes: 6. Same locality.

CHARACIDAE

Vinciguerria mabahiss Johnson and Feltes, 1984

Fieldiana Zool., New Ser., 22: 22.

Paratype. Red Sea: Gulf of Aqaba.

PHOTICHTHYIDAE

Wallago maculatus Inger and Chin, 1959

Fieldiana Zool., 39(27): 279.

Holotype: FMNH 68038. Malaysia: Sabah, Kinabatangan River at Deramakot.

Paratypes: 11. Malaysia: Sabah, 2 localities.

SILURIDAE

Xiliphius magdalenae Eigenmann, 1912

Indiana Univ. Stud., 16: 10.

Holotype: FMNH 56039 (CM 4829). Colombia: Girardot.

Spelled *Xyliphius* and *Xiliphius* in Eigenmann (1912), but spelled as above in subsequent usages by Eigenmann and others.

ASPREINIDAE

Xiphophorus jalapae Meek, 1902

Field Mus. Nat. Hist. Zool. Ser., 3(6): 107.

Holotype: FMNH 3724. Mexico: Veracruz, Jalapa.

Allotype and paratypes: 110. Same locality.

POECILIIDAE

Xyrichthys niveilatus Jordan and Evermann, 1903

Bull. U.S. Fish Comm. 1902, 22: 194.

Paratype. Hawaiian Islands: Honolulu.

LABRIDAE

Xyrichthys sciistius Jordan and Thompson, 1914

Mem. Carnegie Mus., 6(4): 263.

Holotype: FMNH 57090 (CM 6028). Japan: Sagami Bay.

LABRIDAE

Ypsigramma lineata Schultz, 1953

Bull. U.S. Natl. Mus., 202(1): 375.

Paratype: Marshall Islands: Bikini Atoll, Yuro Island.

SERRANIDAE

Zalescopus satsumae Jordan and Hubbs, 1925

Mem. Carnegie Mus., 10(2): 313.

Holotype: FMNH 58839 (CM 7949). Japan: Kagoshima Bay.

URANOSCOPIDAE

Zalescopus tosae Jordan and Hubbs, 1925

Mem. Carnegie Mus., 10(2): 312.

Holotype: FMNH 58835 (CM 7945). Japan: Tosa at Kochi.

URANOSCOPIDAE

Zenarchopterus robertsi Collette, 1982

Copeia, 1982(2): 266.

Paratype: Papua New Guinea: tributary of Kumusi River.

HEMIRAMPHIDAE

Zenarchopterus sepikensis Herre, 1935

Field Mus. Nat. Hist. Zool. Ser., 18(12): 321.

Holotype: FMNH 17213. New Guinea: Sepik River at Koragu.

Paratypes: 10. New Guinea: 3 localities.

HEMIRAMPHIDAE

Zestichthys tanakae Jordan and Hubbs, 1925

Mem. Carnegie Mus., 10(2): 321.

Holotype: FMNH 58841 (CM 7951). Japan: Kushiro.

ZOARCIDAE

Zingel streber nerensis Banareescu and Nalbant, 1979

Trav. Mus. D'Hist. Nat. "Grigore Antipa", 20(1): 353.

Paratype. Romania: Banat, Nera River at Sasca.

PERCIDAE

Zoogoneticus diazi Meek, 1902

Field Mus. Nat. Hist. Zool. Ser., 3(6): 93.

Holotype: FMNH 3618. Mexico: Lago de Patzcuaro at Patzcuaro.

Paratypes: 126. Mexico: 2 localities.

GOODEIDAE

Zoogoneticus miniatus Meek, 1902

Field Mus. Nat. Hist. Zool. Ser., 3(6): 94.

Holotype: FMNH 3680. Mexico: Lago de Chalco near Chalco.

Paratypes: 4. Same locality.

GOODEIDAE

Zygonectes pulvereus Evermann, 1892

Bull. U.S. Fish Comm. 1891, 11: 85.

Paratype. Texas: Corpus Christi.

CYPRINODONTIDAE

List of Taxa Represented by Types

Arranged Alphabetically by Family

ACANTHOCLINIDAE

Acanthoplesiops hiatti Schultz, 1953

ACANTHURIDAE

Acanthurus randalli Briggs and Caldwell, 1957

Acanthurus triostegus marquesensis Schultz and Woods, 1948

Teuthis atrimentatus Jordan and Evermann, 1903

ACIPENSERIDAE

Acipenser oxyrhynchus desotoi Vladikov, 1955

AGENEIOSIDAE

Ageneiosus guianensis Eigenmann, 1912

Ageneiosus madeirensis Fisher, 1917

Ageneiosus marmoratus Eigenmann, 1912

Tympanopleura piperata Eigenmann, 1912

AGONIDAE

Ihuriella kasawae Jordan and Hubbs, 1925

ALBULIDAE

Albula virgata Jordan and Jordan, 1922

ALEPOCEPHALIDAE

Alepocephalus owstoni Tanaka, 1908

Alepocephalus umbriceps Jordan and Thompson, 1914

Grimatrocetes bullisi Grey, 1958

ANABANTIDAE

Ctenopoma multispinis Peters, 1844

ANOSTOMIDAE

Anostomus plicatus Eigenmann, 1912

Leporinodus retropinnis Eigenmann, 1922

Leporinus alternus Eigenmann, 1912

Leporinus arcus Eigenmann, 1912

Leporinus brunneus Myers, 1950

Leporinus ecuadoriensis Eigenmann and Henn, 1916

Leporinus granti Eigenmann, 1912

Leporinus steyermarki Inger, 1956

Pseudanos irinae Winterbottom, 1980

Schizodontopsis laticeps Eigenmann, 1912

ANTENNARIIDAE

Antennarius bermudensis Schultz, 1957

Antennarius pauciradiatus Schultz, 1957

Antennarius verrucosus Bean, 1906

Phrynelox atra Schultz, 1957

Phrynelox zebrinus Schultz, 1957

APLOCHEILIDAE

Aplocheilus chobensis Fowler, 1935

Aplocheilus luzonensis Herre and Ablan, 1934

Cynodonichthys tenuis Meek, 1904

Pterolebias hoignei Thomerson, 1974

Pterolebias maculipinnis Weibezahn, 1964

Rachovia pyropunctata Taphorn and Thomerson, 1978

Rivulus breviceps Eigenmann, 1909

Rivulus brunneus Meek and Hildebrand, 1913

Rivulus compressus Henn, 1916

Rivulus frenatus Eigenmann, 1912

Rivulus holmiae Eigenmann, 1909

Rivulus lanceolatus Eigenmann, 1909

Rivulus magdalenae Eigenmann and Henn, 1916

Rivulus montium Hildebrand, 1938

Rivulus stagnatus Eigenmann, 1909

Rivulus stellifer Thomerson and Turner, 1973

Rivulus volcanus Hildebrand, 1938

Rivulus waimacui Eigenmann, 1909

APOGONIDAE

Amia sialis Jordan and Thompson, 1914

Apogonichthys melampodus Blosser, 1909

Apogon abo Herre, 1935

Apogon beauforti Weber, 1908

Apogon leptocaulus Gilbert, 1972

Apogon nigrofasciatus Lachner, 1953

Apogon snyderi Jordan and Evermann, 1903

Apogon wichmanni Weber, 1908
Gymnapogon gracilicauda Lachner, 1953
Gymnapogon urospilotus Lachner, 1953
Mionorus pacificus Herre, 1935
Scepterias fragilis Jordan and Jordan, 1922
Siphamia cuprea Lachner, 1953
Siphamia fuscolineata Lachner, 1953

APTERONOTIDAE

Adontosternarchus clarkae Mago-Leccia, Lundberg, and Baskin, 1985
Adontosternarchus devenanzii Mago-Leccia, Lundberg, and Baskin, 1985
Porotergus gimbeli Ellis, 1912
Porotergus gymnotus Ellis, 1912
Sternarchus hasemani Ellis, 1913
Sternarchus leptorhynchus Ellis, 1912
Sternarchus mariae Eigenmann and Fisher, 1914
Sternarchus rostratus Meek and Hildebrand, 1913

ARGENTINIDAE

Argentina alicae Cohen and Atsoides, 1969
Argentina brucei Cohen and Atsoides, 1969
Argentina georgei Cohen and Atsoides, 1969
Argentina stewarti Cohen and Atsoides, 1969
Glossanodon pygmaeus Cohen, 1958

ARIIDAE

Arius kanganamanensis Herre, 1935
Arius nox Herre, 1935
Arius solidus Herre, 1935
Arius tuyra Meek and Hildebrand, 1923
Felichthys amblops Meek and Hildebrand, 1913
Galeichthys aguadulce Meek, 1904
Hemipimelodus papillifer Herre, 1915
Hemipimelodus velutinus Weber, 1908
Netuma osakae Jordan and Kasawa, 1925
Netuma planifrons Higuchi, Reis, and Araújo, 1982

ASPREDINIDAE

Agmus lyriformis Eigenmann, 1912
Bunocephalus amaurus Eigenmann, 1912
Bunocephalus chamaizelus Eigenmann, 1912
Bunocephalus colombianus Eigenmann, 1912
Bunocephalus depressus Haseman, 1911
Ernstichthys intonsus Stewart, 1985
Hoplomyzon papillatus Stewart, 1985
Xiliphius magdalenae Eigenmann, 1912

ASTROBLEPIDAE

Astroblepus cyclopus santanderensis Eigenmann, 1917

Astroblepus frenatus Eigenmann, 1917
Astroblepus grixalvi micrescens Eigenmann, 1917
Astroblepus latidens Eigenmann, 1917
Astroblepus pholeter Collette, 1962
Astroblepus rosei Eigenmann, 1922
Cyclopium chapmani Eigenmann, 1912
Cyclopium pirrense Meek and Hildebrand, 1913
Cyclopium trifasciatum Eigenmann, 1912
Cyclopium unifasciatum Eigenmann, 1912
Cyclopium vanceae Eigenmann, 1913
Cyclopium ventrale Eigenmann, 1912

ASTRONESTHIDAE

Astronesthes ijimai Tanaka, 1908

ATHERINIDAE

Allanetta crenolepis Schultz, 1953
Atherina evermanni Eigenmann, 1903
Atherina insularum Jordan and Evermann, 1903
Atherina ovalaua Herre, 1935
Atherinichthys nouhuysi Weber, 1910
Atherinops affinis littoralis Hubbs, 1918
Atherinops insularum cedroscensis Hubbs, 1918
Atherinops insularum guadalupae Hubbs, 1918
Atherion elymus aphrozoicus Schultz, 1953
Atherion elymus freyi Schultz, 1953
Chirostoma attenuatum Meek, 1902
Chirostoma consocium Jordan and Hubbs, 1919
Chirostoma jordani Woolman, 1894
Chirostoma labarcae Meek, 1902
Chirostoma mezquital Meek, 1904
Chirostoma patzcuaro Meek, 1902
Chirostoma regani Jordan and Hubbs, 1919
Chirostoma zirahuen Meek, 1902
Hypoatherina barnesi Schultz, 1953
Ischnomembras gabunensis Fowler, 1903
Melaniris balsanus Meek, 1902
Melaniris sardina Meek, 1907
Menidia chagresi Meek and Hildebrand, 1914
Menidia lisa Meek, 1904
Menidia peninsulae atrimentis Kendall, 1902
Menidia starksi Meek and Hildebrand, 1923
Mugilops cyanellus Meek and Hildebrand, 1923
Poblana ferdebueni Solórzano and López, 1965
Thyrina colombiensis Hubbs, 1920
Thyrina guija Hildebrand, 1925
Thyrina meeki Miller, 1907

AUCHENIPTERIDAE

Auchenipterus brevior Eigenmann, 1912
Auchenipterus demerarae Eigenmann, 1912
Entomocorus benjamini Eigenmann, 1917

Epapterus blohmi Vari, Jewett, Taphorn, and Gilbert, 1984
Gelanoglanis stroudi Böhlke, 1980
Glanidium ribeiroi Haseman, 1911
Pseudauchenipterus nigrolineatus Fowler, 1911
Pseudotatia parva Mees, 1974
Trachelyichthys exilis Greenfield and Glodek, 1977
Trachycorystes fisheri Eigenmann, 1916

BAGRIDAE

Bathybagrus tetranema Bailey and Stewart, 1984
Leiocassis robustus Inger and Chin, 1959
Leiogaster melanopus Weber, 1913
Liobagrus anguillicauda Nichols, 1926
Liobagrus nantoenis Oshima, 1919
Lophiobagrus aquilus Bailey and Stewart, 1984
Lophiobagrus brevispinis Bailey and Stewart, 1984
Mystus sabanus Inger and Chin, 1959
Pseudobagrus adiposalis Oshima, 1919
Pseudobagrus taiwanensis Oshima, 1919

BALISTIDAE

Monacanthus tuckeri Bean, 1906

BATHYCLUPEIDAE

Bathyclupea schroederi Dick, 1962

BATHYLAGIDAE

Bathylagus greyae Cohen, 1958
Nansenia ardesiaca Jordan and Thompson, 1914

BATRACHOIDIDAE

Batrachoides waltersi Collette and Russo, 1981
Porichthys bathoiketes Gilbert, 1968
Porichthys kymosemeum Gilbert, 1968
Porichthys myriaster Hubbs and Schultz, 1939
Thalassophryne quadrizonatus Eigenmann, 1922
Triathalassothia gloverensis Greenfield and Greenfield, 1973

BELONIDAE

Belone svetovidovi Knudsen and Wolff, 1970
Rhaphiobelone robusta Schultz, 1953

BELONTHIDAE

Macropodus filamentosus Oshima, 1919

BLENNIDAE

Alticus novemmaculosus Snyder, 1908
Blenniulus proteus Krejsa, 1960
Blennius nicholsi Tavalga, 1954

Chasmodes bosquianus longimaxilla Williams, 1983
Entomacrodus randalli Springer, 1967
Hypsoblennius lignus Meek and Hildebrand, 1928
Petroscirtes kulambangrae Herre, 1935
Plagiotremus phenax Smith-Vaniz, 1976
Salarias martini Herre, 1942
Salarias walensis Herre, 1935
Spinoblennius spiniger Herre, 1935
Tagusa delicata Herre, 1935

BOTHIDAE

Ancylosetta antillarum Gutherz, 1966
Ancylosetta cycloidea Tyler, 1959
Ancylosetta kumperae Tyler, 1959
Ancylosetta microctenus Gutherz, 1966
Engyprosopon xenandrus Gilbert, 1905
Monolene megalepis Woods, 1961
Platophrys inermis Gilbert, 1905
Trichopsetta caribbaea Anderson and Gutherz, 1967

BRAMIDAE

Eumegistus illustris Jordan and Jordan, 1922

BRANCHIOSTOMATIDAE

Branchiostoma bermudae Hubbs, 1922
Branchiostoma floridae Hubbs, 1922

CALLICHTHYIDAE

Aspidoras maculosus Nijssen and Isbrücker, 1976
Cascadura maculocephala Ellis, 1913
Chaenothorax eigenmanni Ellis, 1913
Corydoras australe Eigenmann and Ward, 1907
Corydoras blochi Nijssen, 1971
Corydoras bondi coppenamensis Nijssen, 1970
Corydoras ellisae Gosline, 1940
Corydoras heteromorphus Nijssen, 1970
Corydoras schwartzi surinamensis Nijssen, 1970
Corydoras weitzmani Nijssen, 1971
Hoplosternum magdalenae Eigenmann, 1913
Hoplosternum punctatum Meek and Hildebrand, 1916

CALLIONYMIDAE

Callionymus distethommatus Fowler, 1941
Callionymus hudsoni Fowler, 1941
Callionymus tenuis Fricke, 1981
Callionymus wilburi Herre, 1935
Calliurichthys astrinius Jordan and Jordan, 1922
Calliurichthys zaneetes Jordan and Jordan, 1922

Calymmichthys xenicus Jordan and Thompson, 1914

Synchiropus ijimae Jordan and Thompson, 1914

Synchiropus minutulus Fricke, 1981

CAPROIDAE

Antigonia combatia Berry and Rathjen, 1958

Antigonia malayana Weber, 1913

CARANGIDAE

Alectis breviventralis Wakiya, 1924

Alectis temmincki Wakiya, 1924

Caranx formosanus Jordan and Snyder, 1909

Caranx ishikawai Wakiya, 1924

Caranx miyakamii Wakiya, 1924

Caranx oshimai Wakiya, 1924

Caranx rastrosus Jordan and Snyder, 1909

Caranx tanakai Wakiya, 1924

Caranx uii Wakiya, 1924

Decapterus dayi Wakiya, 1924

Decapterus tabl Berry, 1968

Hemicaranx rhomboides Meek and Hildebrand, 1925

Scomberoides formosanus Wakiya, 1924

Trachinotus jordani Wakiya, 1924

Trachurus argenteus Wakiya, 1924

Ulua richardsoni Jordan and Snyder, 1909

CARAPIDAE

Disparichthys fluviatilis Herre, 1935

Echiodon dawsoni Williams and Shipp, 1982

CARCHARINIDAE

Rhizoprionodon oligolinx Springer, 1964

CATOSTOMIDAE

Carpiodes elongatus Meek, 1904

Carpiodes labiosus Meek, 1904

Carpiodes microstomus Meek, 1904

Catostomus conchos Meek, 1902

Catostomus sonorensis Meek, 1902

Moxostoma rubriques Hubbs, 1930

Pantosteus jordani Evermann, 1893

Pantosteus santaanae Snyder, 1908

CENTRARCHIDAE

Lepomis occidentalis Meek, 1902

Micropterus pseudaplites Hubbs, 1927

CENTRISCIDAE

Centriscus capito Oshima, 1922

CENTROPOMIDAE

Ambassis lafa Jordan and Seale, 1906

Ambassis vaivasensis Jordan and Seale, 1906

CETOPSIDAE

Hemicetopsis amphioxus Eigenmann, 1914

Hemicetopsis macilentus Eigenmann, 1912

Hemicetopsis minutus Eigenmann, 1912

Hemicetopsis othonops Eigenmann, 1912

CHAENOPSIDAE

Acanthemblemaria greenfieldi Smith-Vaniz and Palacio, 1974

Chaenopsis alepidota californiensis Böhlke, 1957

Chaenopsis roseola Hastings and Shipp, 1980

Ekemblemaria myersi Stephens, 1963

Emblemaria angelica Böhlke and Mead, 1957

Emblemaria hyltoni Johnson and Greenfield, 1976

Emblemaria nigra Meek and Hildebrand, 1928

Emblemariopsis pricei Greenfield, 1975

CHARACIDAE

Acanthocharax microlepis Eigenmann, 1912

Acestrorhynchus altus Menezes, 1969

Acestrorhynchus guianensis Menezes, 1969

Acestrorhynchus nasutus Eigenmann, 1912

Agoniates anchovia Eigenmann, 1914

Alestes langi Fowler, 1935

Alestes thamalakanensis Fowler, 1935

Ammocryptocharax elegans Weitzman and Kanazawa, 1976

Aphyocharax anisitsi Eigenmann and Kennedy, 1903

Aphyocharax erythrurus Eigenmann, 1912

Aphyocharax melanotus Eigenmann, 1912

Aphyocharax paraguayensis Eigenmann, 1915

Aphyocheiroduon hemigrammus Eigenmann, 1915

Aphyodite grammica Eigenmann, 1912

Asiphonichthys hemigrammus Eigenmann, 1912

Astyanax abramoides Eigenmann, 1909

Astyanax aeneus costaricensis Meek, 1914

Astyanax albeolus Eigenmann, 1908

Astyanax aurocaudatus Eigenmann, 1913

Astyanax bimaculatus novae Eigenmann, 1911

Astyanax bimaculatus paraguayensis Eigenmann, 1921

Astyanax daguae Eigenmann, 1913

Astyanax essequibensis Eigenmann, 1909

Astyanax grandis Meek and Hildebrand, 1912

Astyanax guaporensis Eigenmann, 1911

Astyanax guianensis Eigenmann, 1909

Astyanax gymnogenys Eigenmann, 1911

Astyanax hasemani Eigenmann, 1914

Astyanax heterurus Eigenmann and Wilson, 1914

Astyanax kompi Hildebrand, 1938

- Astyanax magdalenae* Eigenmann and Henn, 1916
- Astyanax marionae* Eigenmann, 1911
- Astyanax metae* Eigenmann, 1914
- Astyanax microlepis* Eigenmann, 1913
- Astyanax mucronatus* Eigenmann, 1909
- Astyanax mutator* Eigenmann, 1909
- Astyanax nasutus* Meek, 1907
- Astyanax notemigonoides* Fowler, 1911
- Astyanax paranahybae* Eigenmann, 1911
- Astyanax potaroensis* Eigenmann, 1909
- Astyanax regani* Meek, 1909
- Astyanax ribeirae* Eigenmann, 1911
- Astyanax robustus* Meek, 1912
- Astyanax ruberrimus* Eigenmann, 1913
- Astyanax scabripinnis paranae* Eigenmann, 1914
- Astyanax sciurus* Fowler, 1911
- Bleptonema amazonae* Eigenmann, 1914
- Bleptonema paraguayensis* Eigenmann, 1914
- Brachyhalcinus guianensis* Boeseman, 1952
- Bramocharax elongatus* Meek, 1907
- Bryconamericus alfredae* Eigenmann, 1927
- Bryconamericus alpha* Eigenmann, 1914
- Bryconamericus beta* Eigenmann, 1914
- Bryconamericus cascajalensis* Meek and Hildebrand, 1916
- Bryconamericus caucanus* Eigenmann, 1913
- Bryconamericus chocoensis* Eigenmann, 1913
- Bryconamericus cismontanus* Eigenmann, 1914
- Bryconamericus conventus* Eigenmann, 1913
- Bryconamericus decurrens* Eigenmann, 1913
- Bryconamericus dentatus* Eigenmann, 1913
- Bryconamericus deuterodonoides* Eigenmann, 1914
- Bryconamericus deuterodonoides caudovitatus* Inger, 1956
- Bryconamericus diquensis* Eigenmann, 1913
- Bryconamericus grosvenori* Eigenmann, 1927
- Bryconamericus henni* Eigenmann, 1914
- Bryconamericus hyphesson* Eigenmann, 1909
- Bryconamericus magdalenensis* Eigenmann, 1913
- Bryconamericus novae* Eigenmann and Henn, 1914
- Bryconamericus ortholepis* Eigenmann, 1913
- Bryconamericus osgoodi* Eigenmann and Allen, 1942
- Bryconamericus pachacuti* Eigenmann, 1927
- Bryconamericus peruanus ricae* Eigenmann, 1908
- Bryconamericus scopiferus* Eigenmann, 1913
- Bryconamericus scopiferus guaytarae* Eigenmann and Henn, 1914
- Bryconamericus tolimae* Eigenmann, 1913
- Bryconamericus zeteki* Hildebrand, 1938
- Brycon argenteus* Meek and Hildebrand, 1913
- Brycon behreae* Hildebrand, 1938
- Brycon henni* Eigenmann, 1913
- Brycon obscurus* Hildebrand, 1938
- Brycon petrosus* Meek and Hildebrand, 1913
- Brycon siebenthalae* Eigenmann, 1912
- Ceratobranchia binghami* Eigenmann, 1927
- Characidium blennioides* Eigenmann, 1909
- Characidium catenatum* Eigenmann, 1909
- Characidium caucanus* Eigenmann, 1912
- Characidium laterale* Eigenmann, 1909
- Characidium pellucidum* Eigenmann, 1909
- Characidium phoxocephalum* Eigenmann, 1912
- Characidium pteroides* Eigenmann, 1909
- Characidium vintoni* Eigenmann, 1909
- Characidium zebra* Eigenmann, 1909
- Charax metae* Eigenmann, 1922
- Charax rupununi* Eigenmann, 1912
- Cheirodon annae* McAtee, 1903
- Cheirodon dialepturus* Fink and Weitzman, 1974
- Cheirodon eigenmanni* Meek, 1912
- Cheirodon galusdae* Eigenmann, 1927
- Cheirodon interruptus ibicuhiensis* Eigenmann, 1915
- Cheirodon madeirae* Eigenmann, 1915
- Cheirodon microdon* Eigenmann, 1915
- Cheirodon notomelas* Eigenmann, 1915
- Cheirodon parahybae* Eigenmann, 1915
- Cheirodon stenodon* Eigenmann, 1915
- Compsura heterura* Eigenmann, 1915
- Creagrutus beni* Eigenmann, 1911
- Creagrutus brevipinnis* Eigenmann, 1913
- Creagrutus caucanus* Eigenmann, 1913
- Creagrutus magdalenae* Eigenmann, 1913
- Creagrutus melanzonus* Eigenmann, 1909
- Creagrutus notropoides* Meek and Hildebrand, 1912
- Creagrutus simus* Meek and Hildebrand, 1913
- Cynopotamus essequebensis* Eigenmann, 1912
- Dermatocheir catablepta* Durbin, 1909
- Deuterodon acanthogaster* Eigenmann, 1911
- Deuterodon atracaudata* Meek and Hildebrand, 1912
- Deuterodon pinnatus* Eigenmann, 1909
- Deuterodon potaroensis* Eigenmann, 1909
- Fowlerina franciscensis* Eigenmann, 1914
- Genycharax tarpon* Eigenmann, 1912
- Gephyrocharax caucanus* Eigenmann, 1912
- Gephyrocharax chocoensis* Eigenmann, 1912
- Gephyrocharax intermedius* Meek and Hildebrand, 1916
- Gephyrocharax melanocheir* Eigenmann, 1912
- Gephyrocharax whaleri* Hildebrand, 1938

- Glandulocauda inequalis* Eigenmann, 1911
Glandulocauda melanogenys Eigenmann, 1911
Glandulocauda melanopleura Eigenmann, 1911
Hasemania bilineata Ellis, 1911
Hasemania maxillaris Ellis, 1911
Hasemania melanura Ellis, 1911
Hemibrycon boquiae Eigenmann, 1913
Hemibrycon colombianus Eigenmann, 1914
Hemibrycon dariensis Meek and Hildebrand, 1916
Hemibrycon helleri Eigenmann, 1927
Hemigrammus analis Durbin, 1909
Hemigrammus brevis Ellis, 1911
Hemigrammus compressus Meek, 1904
Hemigrammus cylindricus Durbin, 1909
Hemigrammus erythrozonus Durbin, 1909
Hemigrammus iota Durbin, 1909
Hemigrammus kennedyi Eigenmann, 1903
Hemigrammus marginatus Ellis, 1911
Hemigrammus melanopterus Eigenmann and Kennedy, 1903
Hemigrammus minutus Meek and Hildebrand, 1912
Hemigrammus orthus Durbin, 1909
Hemigrammus riddlei Meek, 1907
Hemigrammus rodwayi Durbin, 1909
Heterocharax macrolepis Eigenmann, 1912
Holesthes heterodon Eigenmann, 1915
Holopristis riddlei Meek, 1907
Hyphessobrycon bifasciatus Ellis, 1911
Hyphessobrycon duragenys Ellis, 1911
Hyphessobrycon ecuadoriensis Eigenmann and Henn, 1914
Hyphessobrycon eos Durbin, 1909
Hyphessobrycon melanopleurus Ellis, 1911
Hyphessobrycon minimus Durbin, 1909
Hyphessobrycon minor Durbin, 1909
Hyphessobrycon panamensis daguae Eigenmann, 1922
Hyphessobrycon parvellus Ellis, 1911
Hyphessobrycon poecilioides Eigenmann, 1913
Hyphessobrycon proteus Eigenmann, 1913
Hyphessobrycon reticulatus Ellis, 1911
Hyphessobrycon rosaceus Durbin, 1909
Hyphessobrycon stictus Durbin, 1909
Hyphessobrycon taurocephalus Ellis, 1911
Hysteronotus megalostomus Eigenmann, 1911
Landonia latidens Eigenmann and Henn, 1914
Macropsobrycon uruguayanae Eigenmann, 1915
Megalamphodus megalopterus Eigenmann, 1915
Megalamphodus micropterus Eigenmann, 1915
Metynnis guaporensis Eigenmann, 1915
Metynnis mola Eigenmann and Kennedy, 1903
Metynnis roosevelti Eigenmann, 1915
Microbrycon minutus Eigenmann and Wilson, 1914
Microgenys minutus Eigenmann, 1913
Microschemobrycon guaporensis Eigenmann, 1915
Moenkhausia browni Eigenmann, 1909
Moenkhausia lepidura hasemani Eigenmann, 1917
Moenkhausia metae Eigenmann, 1922
Moenkhausia phaeonota Fink, 1979
Moenkhausia profunda Eigenmann, 1912
Moenkhausia shideleri Eigenmann, 1909
Mylossoma ocellatum Eigenmann, 1915
Nematobrycon amphiloxus Eigenmann and Wilson, 1914
Nematobrycon lacortei Weitzman and Fink, 1971
Odontostilbe hastatus Eigenmann, 1913
Odontostilbe melandetus Eigenmann, 1912
Odontostilbe paraguayensis Eigenmann and Kennedy, 1903
Odontostilbe trementinae Eigenmann and Kennedy, 1903
Oligobrycon microstomus Eigenmann, 1915
Parastremma sadina Eigenmann, 1912
Parecbasis cyclolepis Eigenmann, 1914
Petersius barnardi Herre, 1936
Petersius conserialis Hilgendorf, 1894
Phenacogaster beni Eigenmann, 1911
Phenacogaster franciscoensis Eigenmann, 1911
Phenacogaster megalostictus Eigenmann, 1909
Phenacogaster microstictus Eigenmann, 1909
Phenagoniates wilsoni Eigenmann, 1914
Piabina analis Eigenmann, 1914
Poecilocharax bovalii Eigenmann, 1909
Poecilurichthys moenkhausii Eigenmann and Kennedy, 1903
Pristella aubynei Eigenmann, 1909
Probolodus heterostomus Eigenmann, 1911
Psalidodon gymnodontus Eigenmann, 1911
Pseudocheiroidon affinis Meek and Hildebrand, 1916
Pseudocorynopoma heterandria Eigenmann, 1914
Pterobrycon landoni Eigenmann, 1913
Pygocentrus bilineatus Eigenmann, 1909
Rhoadsia altipinna Fowler, 1911
Roeboides cauae Eigenmann, 1922
Roeboides hildebrandi Eigenmann, 1922
Roeboides macrolepis Meek and Hildebrand, 1913
Roeboides magdalenae Eigenmann, 1922
Roeboides meeki Eigenmann, 1922

Roeboides occidentalis Meek and Hildebrand, 1916
Roeboides salvadoris Hildebrand, 1925
Roeboides thurni Eigenmann, 1912
Serrasalmo hollandi Eigenmann, 1915
Serrasalmo humeralis gracilior Eigenmann, 1915
Serrasalmus coccogenis Fowler, 1911
Spintherobolus broccae Myers, 1925
Spintherobolus papilliferus Eigenmann, 1911
Stethaprion crenatus Eigenmann, 1916
Stevardia aliata Eigenmann, 1914
Tetragonopterus alleni Eigenmann and McAtee, 1907
Vesicatrux tegatus Eigenmann, 1911

CHILODONTIDAE

Tylobronchus maculosus Eigenmann, 1912

CHIMAERIDAE

Psychichthys eidolon Jordan and Hubbs, 1925

CHLOROPHTHALMIDAE

Bathypterois bigelowi Mead, 1958
Bathypterois dubius Vaillant, 1888
Bathypterois pectoralis Garman, 1899
Bathytrophops marionae Mead, 1958

CICHLIDAE

Aequidens awani Haseman, 1911
Aequidens azurifer Fowler, 1911
Aequidens chimantanus Inger, 1956
Aequidens duopunctata Haseman, 1911
Aequidens guaporensis Haseman, 1911
Aequidens mariae Eigenmann, 1922
Aequidens plagiozonatus Kullander, 1984
Aequidens potaroensis Eigenmann, 1912
Apistogramma caetei Kullander, 1980
Apistogramma inconspicua Kullander, 1982
Apistogramma macmasteri Kullander, 1979
Astronotus orbiculatus Haseman, 1911
Chromis mossambicus Peters, 1852
Chromis sparsidens Hilgendorf, 1905
Cichlasoma acutum Miller, 1907
Cichlasoma alfari Meek, 1907
Cichlasoma boliviense Kullander, 1983
Cichlasoma calobrense Meek and Hildebrand, 1913
Cichlasoma centrale Meek, 1906
Cichlasoma dorsatum Meek, 1907
Cichlasoma eigenmanni Meek, 1902
Cichlasoma evermanni Meek, 1904
Cichlasoma frontale Meek, 1909
Cichlasoma grammodes Taylor and Miller, 1980
Cichlasoma granadense Meek, 1907

Cichlasoma hartwegi Taylor and Miller, 1980
Cichlasoma hedricki Meek, 1904
Cichlasoma meeki Hildebrand, 1925
Cichlasoma milleri Meek, 1907
Cichlasoma mojarra Meek, 1904
Cichlasoma nigratum Meek, 1907
Cichlasoma ornatum gephyrum Eigenmann, 1922
Cichlasoma paranaense Kullander, 1983
Cichlasoma punctatum Meek, 1909
Cichlasoma spilatum Meek, 1912
Cichlasoma tenue Meek, 1906
Cichlasoma tuba Meek, 1912
Cichlasoma tuyrense Meek and Hildebrand, 1913
Cichlasoma umbriferum Meek and Hildebrand, 1913
Cichlasoma urophthalmus mayorum Hubbs, 1936
Cichlasoma zonatum Meek, 1905
Crenicara altispinosa Haseman, 1911
Crenicichla alta Eigenmann, 1912
Crenicichla dorsocellata Haseman, 1911
Crenicichla iguassuensis Haseman, 1911
Crenicichla jaguarensis Haseman, 1911
Crenicichla santaremensis Haseman, 1911
Crenicichla simoni Haseman, 1911
Geophagus brasiliensis iporangensis Haseman, 1911
Geophagus brasiliensis itapicuruensis Haseman, 1911
Gobiocichla ethelwynnae Roberts, 1982
Gobiocichla wonderi Kanazawa, 1951
Heros beani Jordan, 1889
Heterogramma corumbae Eigenmann and Ward, 1907
Heterogramma ortmanni Eigenmann, 1912
Heterogramma ritense Haseman, 1911
Heterogramma taeniatum pertense Haseman, 1911
Heterogramma trifasciatum maciliense Haseman, 1911
Lamprologus obliquus Nichols and Griscom, 1917
Nannacara bimaculata Eigenmann, 1912
Nanochromis transvestitus Stewart and Roberts, 1984
Neotrophus carpintis Jordan and Snyder, 1900
Neotrophus fluviatilis Meek, 1912
Neotrophus panamensis Meek and Hildebrand, 1913
Thorichthys ellioti Meek, 1904

CITHARINIDAE

Distichodus stigmaturus Fowler, 1935

CLARIIDAE

Clarias malaris Nichols and Griscom, 1917

Clarias mossambicus Peters, 1852

CLINIDAE

Gibbonsia elegans erroli Hubbs, 1952

Gibbonsia elegans montereyensis Hubbs, 1927

Gibbonsia elegans velifera Hubbs, 1952

Gibbonsia metzi Hubbs, 1927

Gibbonsia norae Hubbs, 1952

CLUPEIDAE

Alosa ohiensis Evermann, 1902

Dorosoma anale Meek, 1904

Dorosoma chavesi Meek, 1907

Gonostoma ebelingi Grey, 1960

Ilisha argentata Meek and Hildebrand, 1923

Jenkinsia majua majua Whitehead, 1963

Jenkinsia majua woodsi Whitehead, 1963

Odontognathus compressus Meek and Hildebrand, 1923

Opisthonema berlangai Berry and Barrett, 1963

Opisthonema medirastre Berry and Barrett, 1963

Pellonula tenuis Nichols and Griscom, 1917

Rhinosardinia serrata Eigenmann, 1912

Sardinella marquesensis Berry and Whitehead, 1968

Signalosa atchafalayae vanhyningi Weed, 1925

Signalosa mexicana campi Weed, 1925

COBITIDAE

Acanthophtalmus mariae Inger and Chin, 1962
Acanthophtalmus sandakanensis Inger and Chin, 1962

Barbatula toni posteroventralis Nichols, 1925

Barbatula yarkandensis sellaefer Nichols, 1925

Botia purpurea Nichols, 1925

Cobitis aurata vallachica Nalbant, 1957

Cobitis taenia melanoleuca Nichols, 1925

Crossostoma fascicauda Nichols, 1926

Misgurnus anguillicaudatus tungting Nichols, 1925

Misgurnus mizolepis fukien Nichols, 1925

Misgurnus mizolepis grangeri Nichols, 1925

Misgurnus mohoity yunnan Nichols, 1925

Oreias furcatus Banareescu and Nalbant, 1976

CONGRIDAE

Conger jordani Kanazawa, 1958

Leptocephalus ectenurus Jordan and Richardson, 1909

Leptocephalus harringtonensis Mowbray, 1931

CORYPHAENIDAE

Ectenias brunneus Jordan and Thompson, 1914

COTTIDAE

Clinocottus analis australis Hubbs, 1926

Leptocottus armatus australis Hubbs, 1921

CURIMATIDAE

Curimata cerasina Vari, 1983

Curimata incompta Vari, 1983

Curimatus atratoensis Eigenmann, 1912

Curimatus gillii Eigenmann and Kennedy, 1903

Curimatus issororoensis Eigenmann, 1912

Curimatus metae Eigenmann, 1922

Curimatus morawhannae Eigenmann, 1912

Curimatus patiae Eigenmann, 1914

Curimatus peruanus Eigenmann, 1922

Psectrogaster curviventris Eigenmann and Kennedy, 1903

CYCLOPTERIDAE

Careproctus burkei Jordan and Thompson, 1914

Careproctus gilberti Jordan and Thompson, 1914

CYNOGLOSSIDAE

Areliscus hollandi Jordan and Metz, 1913

Cynoglossus heterolepis Weber, 1910

Dollfusichthys sinusarabici Chabanaud, 1931

CYPRINIDAE

Achahara jusanensis Jordan and Hubbs, 1925

Acheilognathus barbatus Nichols, 1926

Acheilognathus gracilis Nichols, 1926

Acheilognathus tabira Jordan and Thompson, 1914

Agosia adobe Jordan and Evermann, 1891

Agosia yarrowi Jordan and Evermann, 1891

Alburnops illecebrosus Girard, 1857

Algansea aphanea Barbour and Miller, 1978

Algansea monticola Barbour and Contreras, 1968

Algansea rubescens Meek, 1902

Amblypharyngodon grandisquamis Jordan and Starks, 1917

Aztecula mexicana Meek, 1902

Barbus akeleyi Hubbs, 1918

Barbus anoplus Weber, 1897

Barbus athi Hubbs, 1918

Barbus bantolanensis Day, 1914

Barbus bifrenatus Fowler, 1935

Barbus carens Boulenger, 1912

Barbus fitzsimonsi Fowler, 1935

- Barbus helleri* Hubbs, 1918
Barbus oatesii Boulenger, 1893
Barbus tsotsorogensis Fowler, 1935
Beirabarbuis palustris Herre, 1936
Belligobio eristigma Jordan and Hubbs, 1925
Chondrostoma cyri orientalis Bianco and Ban-arescu, 1982
Coripareius septentrionalis Nichols, 1925
Culter aokii Oshima, 1919
Dangila sabana Inger and Chin, 1962
Dionda diaboli Hubbs and Brown, 1956
Dionda erimyzonops Hubbs and Miller, 1974
Evarra tlahuacensis Meek, 1902
Formosania gilberti Oshima, 1919
Gila minacae Meek, 1902
Gnathopogon iijimae Oshima, 1919
Gnathopogon ishiwakae Jordan and Thompson, 1914
Gnathopogon longifilis Jordan and Hubbs, 1925
Gnathopogon majimae Jordan and Hubbs, 1925
Gnathopogon suwae Jordan and Hubbs, 1925
Gnathopogon tsuchigae Jordan and Hubbs, 1925
Gobio coriparoides Nichols, 1925
Gobio kesleri banaticus Banarescu, 1953
Gobio rivuloides Nichols, 1925
Gobio soldatovi minus Nichols, 1925
Hampala macrolepidota sabana Inger and Chin, 1962
Hemiculterella engraulis Nichols, 1925
Hybognathus hankinsoni Hubbs, 1929
Hybopsis x-punctata trautmani Hubbs and Crowe, 1956
Hybopsis x-punctata x-punctata Hubbs and Crowe, 1956
Labeo fisheri Jordan and Starks, 1917
Labeo jordani Oshima, 1919
Lepidomeda mollispinis mollispinis Miller and Hubbs, 1960
Leuciscus carletoni Kendall, 1903
Leucogobio polytaenia Nichols, 1925
Moroco yamamotis Jordan and Hubbs, 1925
Nematabramis alestes borneensis Inger and Chin, 1962
Nototropis forlonensis Meek, 1904
Nototropis nazas Meek, 1904
Notropis aztecus Woolman, 1894
Notropis chihuahua Woolman, 1894
Notropis edwardraneyi Suttkus and Clemmer, 1968
Notropis greeni Hubbs and Ortenburger, 1929
Notropis muskoka Meek, 1899
Notropis panarcys Hubbs and Miller, 1978
Notropis robustus Meek, 1902
Notropis saladonis Hubbs and Hubbs, 1958
Notropis santarosaliae Meek, 1902
Paracrossochilus aceris Inger and Chin, 1962
Parapelecus eigenmanni Jordan and Metz, 1913
Phoxinus cumberlandensis Starnes and Starnes, 1978
Phoxinus orcuttii Eigenmann and Eigenmann, 1890
Phoxiscus kikuchii Oshima, 1919
Pimephales anuli Kendall, 1903
Pseudaspius atrilatus Jordan and Thompson, 1914
Pseudaspius bergi Jordan and Metz, 1913
Pseudaspius modestus Jordan and Metz, 1913
Pseudogobio chinssuensis Nichols, 1926
Pseudoperilampus honda Jordan and Metz, 1913
Pseudorasbora altipinna Nichols, 1925
Pseudorasbora depressirostris Nichols, 1925
Puntius joaquinae Wood, 1968
Puntius snyderi Oshima, 1919
Puntius umali Wood, 1968
Rhodeus chosonicus Jordan and Metz, 1913
Rhodeus kurumeus Jordan and Thompson, 1914
Sarcocheilichthys morii Jordan and Hubbs, 1925
Scaphesthes tamusuiensis Oshima, 1919
Spinibarbus hollandi Oshima, 1919
Varicorhinus robustus Nichols, 1925
- CYPRINODONTIDAE**
- Cyprinodon bifasciatus* Miller, 1968
Cyprinodon dearborni Meek, 1909
Cyprinodon fontinalis Smith and Miller, 1980
Cyprinodon meeki Miller, 1976
Cyprinodon orthonotus Peters, 1844
Fundulus cubensis Eigenmann, 1903
Fundulus neumanni Hilgendorf, 1905
Fundulus oaxacae Meek, 1902
Orestias mundus Parenti, 1984
Profundulus candalaria Hubbs, 1924
Zygonectes pulvereus Evermann, 1892
- DACTYLOSCOPIDAE**
- Gillellus greyae* Kanazawa, 1952
- DASYATIDAE**
- Dasyatis ushieii* Jordan and Hubbs, 1925
- DICERATIIDAE**
- Diceratias pileatus* Uwate, 1979
- DIRETMIDAE**
- Diretmus pauciradiatus* Woods, 1973

DORADIDAE

- Anduzedoras arleoi* Fernández-Yépez, 1968
Anduzedoras copei Fernández-Yépez, 1968
Apuredoras rivasi Fernández-Yépez, 1950
Doraops zuloagai Schultz, 1944
Doras lentiginosus Eigenmann, 1917
Hemidoras leporhinus Eigenmann, 1912
Hemidoras micropoetus Eigenmann, 1912
Hemidoras microstomus Eigenmann, 1912
Hemidoras notospilus Eigenmann, 1912
Hildadoras bolivarensis Fernández-Yépez, 1968
Leptodoras linnelli Eigenmann, 1912
Rhinodoras boehlkei Glodek, Whitmire, and Orcés, 1976
Rhinodoras thomersoni Taphorn and Lilystrom, 1984
Rhynchodoras woodsi Glodek, 1976

DRACONETTIDAE

- Draconetta oregona* Briggs and Berry, 1959

ELEOTRIDAE

- Alepideleotris tigris* Herre, 1935
Boroda malua Herre, 1935
Bostrychus zonatus Weber, 1908
Dormitator gymnocephalus Eigenmann, 1912
Eleotris herwerdenii Weber, 1910
Eleotris heterodon Weber, 1908
Eleotris isthmensis Meek and Hildebrand, 1916
Eleotris latifasciatus Meek and Hildebrand, 1912
Eleotris macrolepis Meek, 1914
Encaeura evides Jordan and Hubbs, 1925
Euleptoeleotris clarki Hildebrand, 1938
Euleptoeleotris shropshirei Hildebrand, 1938
Hemieleotris levis Eigenmann, 1917
Leptophilypnus fluviatilis Meek and Hildebrand, 1916
Microeleotris mindii Meek and Hildebrand, 1916
Microeleotris panamensis Meek and Hildebrand, 1916
Pogoneleotris microps Weber, 1908

EMBIOTOCIDAE

- Crossochir koelzi* Hubbs, 1933

ENGRAULIDAE

- Anchoa hepsetus colonensis* Hildebrand, 1943
Anchoa lamprotaenia Hildebrand, 1943
Anchoa mitchilli diaphana Hildebrand, 1943
Anchovia arenicola Meek and Hildebrand, 1923
Anchovia brevirostra Meek and Hildebrand, 1923
Anchovia eigenmannia Meek and Hildebrand, 1923
Anchovia elongata Meek and Hildebrand, 1923

- Anchovia parva* Meek and Hildebrand, 1923
Anchoviella belizensis Thomerson and Greenfield, 1975
Anchoviella helleri Hubbs, 1921
Stolephorus branchiomelas Eigenmann, 1917
Stolephorus guianensis Eigenmann, 1912
Stolephorus lyolepis Evermann and Marsh, 1902

EVERMANNELLIDAE

- Evermannella ahlstromi* Johnson and Glodek, 1975
Evermannella sicaria Rofen, 1963

EXOCOETIDAE

- Cypselurus alienus* Herre, 1935
Cypselurus antoncichi Woods and Schultz, 1953

GASTEROPELECIDAE

- Thoracocharax brevis* Eigenmann, 1912
Thoracocharax magdalenae Eigenmann, 1912

GEMPYLIDAE

- Diplogonurus maderensis* Noronha, 1926
Epinnula orientalis americana Grey, 1953
Epinnula orientalis pacifica Grey, 1953

GERREIDAE

- Diapterus evermanni* Meek and Hildebrand, 1925
Eucinostomus californiensis elongatus Meek and Hildebrand, 1925

GOBIESOCIDAE

- Cotylin nigripinnis woodsi* Schultz, 1944
Gobiesox costaricensis Meek, 1912
Gobiesox fluviatilis Briggs and Miller, 1960
Gobiesox fulvus Meek, 1907
Gobiesox marijeanae Briggs, 1960
Gobiesox mexicanus Briggs and Miller, 1960
Gobiesox paradiseus Herre, 1935
Gobiesox potamius Briggs, 1955

GOBIIDAE

- Amblygobius myersi* Herre, 1935
Aparrius aurocingulus Herre, 1935
Atuona tricuspidata Herre, 1935
Awaous decemlineatus Eigenmann, 1917
Barbuligobius boehlkei Lachner and McKinney, 1974
Brachygobius kabiliensis Inger, 1958
Brachygobius sabanus Inger, 1958
Brachygobius xanthomelas Herre, 1937
Callogobius ocellatus Herre, 1935
Chasmias misakius Jordan and Snyder, 1901

Coryphopterus punctipectophorus Springer, 1960
Cryptocentrus geniornatus Herre, 1935
Ctenogobius aterrimus Herre, 1935
Ctenogobius leavelli Herre, 1935
Ctenogobius malekulae Herre, 1935
Ctenogobius myxodermus Herre, 1935
Ctenogobius opalescens Herre, 1936
Ctenogobius villadolidi Herre, 1936
Ctenogobius virgatulus Jordan and Snyder, 1901
Ctenogobius waigiensis Herre, 1935
Euctenogobius liolepis Meek and Hildebrand, 1928
Euctenogobius panamensis Meek and Hildebrand, 1928
Eviota japonica Jewett and Lachner, 1983
Evorthodus minutus Meek and Hildebrand, 1928
Gerhardinus nudus Meek and Hildebrand, 1928
Gladiogobius ensifer Herre, 1933
Glossogobius abacopus Jordan and Richardson, 1909
Glossogobius koragensis Herre, 1935
Glossogobius parvus Oshima, 1919
Glossogobius sandakanensis Inger, 1957
Gnatholepis corlettei Herre, 1935
Gnatholepis knighti Jordan and Evermann, 1903
Gobiella birtwistlei Herre, 1934
Gobionellus saepepallens Gilbert and Randall, 1968
Gobiopsis aporia Lachner and McKinney, 1978
Gobiopsis woodsi Lachner and McKinney, 1978
Gobius claytonii Meek, 1902
Gobius cliffordpopei Nichols, 1925
Gobius daguae Eigenmann, 1917
Gobius parvus Meek, 1902
Macgregorella badia Herre, 1935
Macgregorella santa Herre, 1935
Macrodonotogobius wilburi Herre, 1936
Quisquilius eugenius Jordan and Evermann, 1903
Rhinogobius formosanus Oshima, 1919
Rhinogobius mowbrayi Bean, 1906
Rhinogobius taiwanus Oshima, 1919
Sagamia russula Jordan and Snyder, 1901
Sicydium altum Meek, 1907
Sicydium hildebrandi Eigenmann, 1917
Sicydium montanum Hubbs, 1920
Suruga fundicola Jordan and Snyder, 1901
Tridentiger kuroi Jordan and Tanaka, 1927
Vaimosa balteata Herre, 1935
Vaimosa horiae Herre, 1936
Vaimosa osgoodi Herre, 1935
Varicus bucca Robins and Böhlke, 1961
Varicus imswae Greenfield, 1981

GOBIOIDIDAE

Taenioides snyderi Jordan and Hubbs, 1925

GOODEIDAE

Characodon multiradiatus Meek, 1904
Goodea captiva Hubbs, 1924
Goodea toweri Meek, 1904
Goodea withei Meek, 1904
Skiffia lermiae Meek, 1902
Skiffia variegata Meek, 1902
Zoogoneticus diazi Meek, 1902
Zoogoneticus miniatus Meek, 1902

GRAMMISTIDAE

Rhagma bermudensis Kanazawa, 1952

HAEMULIDAE

Anisotremus williamsi Meek, 1905
Pomadasys grandis Meek, 1907
Pomadasys starri Meek, 1904
Pomadasys templei Meek, 1904

HELOGENIDAE

Helogenes unidorsalis Glodek and Carter, 1978

HEMIODONTIDAE

Bivibranchia protractila Eigenmann, 1912
Hemiodus orthonops Eigenmann and Kennedy, 1903
Hemiodus parnaguae Eigenmann and Henn, 1916
Hemiodus quadrimaculatus Pellegrin, 1908

HEMIRAMPHIDAE

Hemiramphus archipelagicus Collette, 1978
Hemiramphus bermudensis Collette, 1962
Hyporhamphus hildebrandi Jordan and Evermann, 1927
Hyporhamphus melanopterus Collette and Parin, 1978
Hyporhamphus snyderi Meek and Hildebrand, 1923
Odontorhamphus cancellori Weed, 1933
Zenarchopterus robertsi Collette, 1982
Zenarchopterus sepikensis Herre, 1935

HETERENCHELYIDAE

Heterenchelys biaggii Böhlke, 1956

HEXAGRAMMIDAE

Pleurogrammus azonus Jordan and Metz, 1913
Stellistius katsukii Jordan and Tanaka, 1927

HOLOCENTRIDAE

- Adioryx poco* Woods, 1965
Flammeo scythrops Jordan and Evermann, 1903
Holocentrus bullisi Woods, 1955
Holocentrus ensifer Jordan and Evermann, 1903
Holocentrus meeki Bean, 1906
Holocentrus xantherythrus Jordan and Evermann, 1903
Myripristis argyromus Jordan and Evermann, 1903
Myripristis berndti Jordan and Evermann, 1903
Myripristis bowditchae Woods, 1953
Myripristis chryseres Jordan and Evermann, 1903
Myripristis mooreanus Herre, 1935
Myripristis randalli Greenfield, 1974
Myripristis symmetricus Jordan and Evermann, 1903
Myripristis tiki Greenfield, 1974
Myripristis undecimalis Herre, 1935
Myripristis woodsi Greenfield, 1974
Ostichthys acanthorhinus Randall, Shimizu, and Yamakawa, 1982
Sargocentron macrosquamis Golani, 1984

HOMALOPTERIDAE

- Gastromyzon fasciatus* Inger and Chin, 1961
Gastromyzon lepidogaster Roberts, 1982
Gastromyzon megalepis Roberts, 1982
Gastromyzon pauciradiatus Inger and Chin, 1961
Gastromyzon punctulatus Inger and Chin, 1961
Gastromyzon ridens Roberts, 1982
Homaloptera hingi Herre, 1934
Protomyzon aphelocheilus Inger and Chin, 1962

ICTALURIDAE

- Ameiurus australis* Meek, 1904
Ameiurus mexicanus Meek, 1904
Noturus albatel Taylor, 1969
Noturus flavater Taylor, 1969
Noturus munitus Suttkus and Taylor, 1965

JENYNSHIDAE

- Fitzroyia eigenmanni* Haseman, 1911

KNERIIDAE

- Grasseichthys gabonensis* Géry, 1964

KUHLIIDAE

- Kuhlia bilunulata* Herre, 1935

LABRIDAE

- Halichoeres fijiensis* Herre, 1935
Hemipteronotus baldwini Jordan and Evermann, 1903

- Hemipteronotus evides* Jordan and Richardson, 1909
Iridio bimaculata Wilson, 1916
Iridio decoratus Bean, 1906
Iridio elegans Bean, 1906
Iridio meyeri Bean, 1906
Iridio microstomus Bean, 1906
Lepidaplois strophodes Jordan and Evermann, 1903
Thalassoma cranei Herre, 1935
Xyrichthys niveilatus Jordan and Evermann, 1903
Xyrichthys sciistius Jordan and Thompson, 1914

LABRISOMIDAE

- Exerpes asper earli* Hubbs, 1952
Labrisomus lentiginosus Bean, 1906
Labrisomus multiporosus Hubbs, 1953
Malacotenus afueriae multipunctatus Springer, 1958
Malacotenus ebisui Springer, 1958
Malacotenus gigas Springer, 1958
Malacotenus hubbsi hubbsi Springer, 1958
Malacotenus hubbsi polyporosus Springer, 1958
Malacotenus margaritae mexicanus Springer, 1958
Malacotenus puertoricensis Evermann and Marsh, 1900
Malacotenus triangulatus Springer, 1958
Malacotenus zaca Springer, 1958
Malacotenus zonifer sudensis Springer, 1958
Mnierpes macrocephalus catherinae Hubbs, 1952
Nemaclinus atelestos Böhlke and Springer, 1975
Paraclinus integripinnis zaca Hubbs, 1952
Starksia occidentalis Greenfield, 1979
Starksia variabilis Greenfield, 1979
Starksia y-lineata Gilbert, 1965

LEBIASINIDAE

- Archicheir minutus* Eigenmann, 1909
Copeina osgoodi Eigenmann, 1922
Nannostomus marginatus Eigenmann, 1909
Nannostomus minimus Eigenmann, 1909
Nannostomus simplex Eigenmann, 1909
Poecilobrycon auratus Eigenmann, 1909
Poecilobrycon erythrurus Eigenmann, 1909
Poecilobrycon harrisoni Eigenmann, 1909
Poecilobrycon ocellatus Eigenmann, 1909
Pyrrhulina australe Eigenmann and Kennedy, 1903

LEIOGNATHIDAE

- Gazza achlamys* Jordan and Starks, 1917

LETHRINIDAE

Gnathodentex oculomaculatus Herre, 1935

LOPHIIDAE

Lophiodes beroe Caruso, 1981

Lophiodes reticulatus Caruso and Suttkus, 1979

LORICARIIDAE

Acanthicus canensis Meek and Hildebrand, 1913

Acestridium discus Haseman, 1911

Ancistrus lithurgicus Eigenmann, 1912

Ancistrus melas Eigenmann, 1916

Ancistrus planiceps Meek and Hildebrand, 1913

Ancistrus spinosus Meek and Hildebrand, 1916

Chaetostomus leucomelas Eigenmann, 1917

Corymbophanes andersoni Eigenmann, 1909

Farlowella azygia Eigenmann and Vance, 1917

Farlowella hasemani Eigenmann and Vance, 1917

Farlowella jauruensis Eigenmann and Vance, 1917

Hemiancistrus daguae Eigenmann, 1912

Hemiancistrus mayoloi Eigenmann, 1912

Hemiancistrus wilsoni Eigenmann, 1917

Lasiancistrus brevispinis Heitmans, Nijssen, and Isbrücker, 1983

Lasiancistrus caucanus Eigenmann, 1912

Lithogenes villosus Eigenmann, 1909

Lithoxus lithoides Eigenmann, 1912

Loricaria capetensis Meek and Hildebrand, 1913

Loricaria filamentosa latiura Eigenmann and Vance, 1912

Loricaria filamentosa seminuda Eigenmann and Vance, 1912

Loricaria fimbriata Eigenmann and Vance, 1912

Loricaria griseus Eigenmann, 1909

Loricaria gymnogaster Eigenmann and Vance, 1912

Loricaria gymnogaster lagoichthys Schultz, 1944

Loricaria microdon Eigenmann, 1909

Loricaria stewarti Eigenmann, 1909

Loricaria submarginatus Eigenmann, 1909

Loricaria tuyrensis Meek and Hildebrand, 1913

Loricaria uracantha rupestre Schultz, 1944

Loricaria variegata venezuelae Schultz, 1944

Oxyloricaria citurensis Meek and Hildebrand, 1913

Oxyloricaria dariensis Meek and Hildebrand, 1913

Parotocinclus jimi Garavello, 1976

Plecostomus derbyi Haseman, 1911

Plecostomus hemiurus Eigenmann, 1912

Pseudancistrus carnegiei Eigenmann, 1916

Pseudancistrus nigrescens Eigenmann, 1912

Pseudancistrus pediculatus Eigenmann, 1917

Pterygoplichthys anisitsi Eigenmann and Kennedy, 1903

Rineloricaria formosa Isbrücker and Nijssen, 1979

LUTJANIDAE

Etelis evurus Jordan and Evermann, 1903

Pristipomoides freemani Anderson, 1966

MACROURIDAE

Coelorhynchus gilberti Jordan and Hubbs, 1925

Coryphaenoides bonanox Jordan and Thompson, 1914

Hymenocephalus aterrimus Gilbert, 1905

Hymenocephalus grimaldii Weber, 1913

Macrurus richardi Weber, 1913

Optonurus atherodon Gilbert and Cramer, 1896

Squalogadus intermedius Grey, 1959

MASTACEMBELIDAE

Mastacembelus keithi Herre, 1940

Mastacembelus oatesii Boulenger, 1893

MELAMPHAIDAE

Melamphaes pumilus Ebeling, 1962

MELANOTAENIIDAE

Melanotaenia kabia Herre, 1935

Melanotaenia rosacea Herre, 1935

Rhombatractus affinis Weber, 1908

Rhombatractus sentaniensis Weber, 1908

Rhombosoma sepikensis Herre, 1935

MICRODESMIDAE

Gunnellichthys viridesens Dawson, 1968

Microdesmus carri Gilbert, 1966

Microdesmus woodsi Kanazawa, 1952

MORIDAE

Physiculus grinnelli Jordan and Jordan, 1922

MUGILIDAE

Joturus daguae Eigenmann, 1917

Liza formosae Oshima, 1922

Liza parva Oshima, 1922

Liza pescadorensis Oshima, 1922

Mugil anpinensis Oshima, 1922

Myxus malayanus Herre, 1936

MULLIDAE

Mulloidies flammeus Jordan and Evermann, 1903

Pseudupeneus chrysonemus Jordan and Evermann, 1903

Upeneus arge Jordan and Evermann, 1903

MURAENIDAE

Anarchias yoshiae Kanazawa, 1952

Echidna unicolor Schultz, 1953

Echidna vineta Jenkins, 1903

Echidna zonophaea Jordan and Evermann, 1903

Enchelycore carychroa Böhlke and Böhlke, 1976

Gymnothorax bayeri Schultz, 1953

Gymnothorax bikiniensis Schultz, 1953

Gymnothorax efatensis Herre, 1935

Gymnothorax leucostigma Jordan and Richardson, 1909

Rabula fuscomaculata Schultz, 1951

Rabula marshallensis Schultz, 1953

Uropterygius fuscoguttatus Schultz, 1953

MYCTOPHIDAE

Dasyscopelus orientalis Gilbert, 1913

Diaphus gigas Gilbert, 1913

Diaphus glandulifer Gilbert, 1913

Diaphus latus Gilbert, 1913

Diaphus sagamiensis Gilbert, 1913

Lampanyctus jordani Gilbert, 1913

Lampanyctus micropunctatus Chapman, 1939

Macrostoma quercinum japonicum Tanaka, 1908

Myctophum hollandi Jordan and Jordan, 1922

MYXINIDAE

Myxine biniplicata Richardson and Jowett, 1951

NEMIPTERIDAE

Parascolopsis townsendi Boulenger, 1901

Scolopsis eriomma Jordan and Richardson, 1909

NEOSCOPELIDAE

Scopelengys clarkei Butler and Ahlstrom, 1976

Solivomer arenidens Miller, 1947

NEOSTETHIDAE

Neostethus borneensis Herre, 1939

NOMEIDAE

Icticus ischanus Jordan and Thompson, 1914

NOTACANTHIDAE

Polyacanthonotus merretti Sulak, Crabtree, and Hureau, 1984

OGCOCEPHALIDAE

Ogcocephalus corniger Bradbury, 1980

Ogcocephalus declivirostris Bradbury, 1980

Ogcocephalus pantostictus Bradbury, 1980

Ogcocephalus pumilus Bradbury, 1980

Ogcocephalus rostellum Bradbury, 1980

OPHICHTHIDAE

Aplatophis chauliodus Böhlke, 1956

Callechelys bilinearis Kanazawa, 1952

Cirriacaecula johnsoni Schultz, 1953

Ophichthus evermanni Jordan and Richardson, 1909

Ophichthus hyposagmatus McCosker and Böhlke, 1984

Sphagebranchus conklini Eigenmann, 1916

OPHIDIIDAE

Calamopteryx jeb Cohen, 1973

Dicrolene kanazawai Grey, 1958

Grammonus mowbrayi Grey, 1951

Lepophidium aporrhox Robins, 1961

Lepophidium pheromystax Robins, 1960

Lepophidium staurophor Robins, 1958

Luciobrotula corethromycter Cohen, 1954

Monomitopus kumae Jordan and Hubbs, 1925

Othophidium schmidtii Woods and Kanazawa, 1951

Porogadus silus Carter and Sulak, 1984

Spectrunculus radcliffei Jordan and Thompson, 1914

OPISTOGNATHIDAE

Lonchopisthus higmani Mead, 1959

OSMERIDAE

Spirinchus dilatus Schultz and Chapman, 1934

Spirinchus verecundus Jordan and Metz, 1913

PANGASIIDAE

Pangasius tubbi Inger and Chin, 1959

PARALEPIDAE

Paralepis philippinus Fowler, 1934

PARODONTIDAE

Apareiodon hasemani Eigenmann, 1916

Apareiodon itapicuruensis Eigenmann and Henn, 1916

Parodon bifasciatus Eigenmann, 1912

Parodon dariensis Meek and Hildebrand, 1913

Parodon ecuadoriensis Eigenmann and Henn, 1914

Parodon terminalis Eigenmann and Henn, 1914

PEMPHERIDAE

Liopempheris sasakii Jordan and Hubbs, 1925

Pempheris dispar Herre, 1935

PERCICHTHYIDAE

Malakichthys wakiyai Jordan and Hubbs, 1925

PERCIDAE

Etheostoma obeyense Kirsch, 1892

Hadropterus scierus apristis Hubbs and Hubbs, 1954

Hololepis saludae Hubbs and Cannon, 1935

Stizostedion glaucum Hubbs, 1926

Zingel streber nerensis Banarescu and Nalbant, 1979

PERCOPHIDAE

Bembrops greyi Poll, 1959

PETROMYZONTIDAE

Ichthyomyzon unicuspis Hubbs and Trautman, 1937

PHOTICHTHYIDAE

Vinciguerria mabahiss Johnson and Feltes, 1984

PIMELODIDAE

Brachyglanis frenata Eigenmann, 1912

Brachyglanis melas Eigenmann, 1912

Brachyglanis phalacra Eigenmann, 1912

Cephalosilurus fowleri Haseman, 1911

Cetopsorhamdia boquillae Eigenmann, 1922

Cetopsorhamdia nasus Eigenmann and Fisher, 1916

Cetopsorhamdia phantasia Stewart, 1985

Chasmocranus brevior Eigenmann, 1912

Chasmocranus chimantanus Inger, 1956

Chasmocranus longior Eigenmann, 1912

Chasmocranus rosae Eigenmann, 1922

Cheirocerus eques Eigenmann, 1917

Heptapterus stewarti Haseman, 1911

Imparfinis hollandi Haseman, 1911

Imparfinis microps Eigenmann and Fisher, 1916

Imparfinis mirini Haseman, 1911

Leptoglanis essequibensis Eigenmann, 1912

Megalonema platycephalum Eigenmann, 1912

Megalonema punctatum Meek and Hildebrand, 1913

Megalonema robustum Meek and Hildebrand, 1913

Megalonema xanthum Eigenmann, 1912

Microglanis iheringi Gomes, 1946

Microglanis poecilus Eigenmann, 1912

Microglanis secundus Mees, 1974

Microglanis variegatus Eigenmann, 1914

Myoglanis potaroensis Eigenmann, 1912

Nannorhamdia nemacheir Eigenmann and Fisher, 1916

Pimelodella avanhandavae Eigenmann, 1917

Pimelodella boliviana Eigenmann, 1917

Pimelodella eigenmanni Meek, 1905

Pimelodella griffini Eigenmann, 1917

Pimelodella hasemani Eigenmann, 1917

Pimelodella itapicuruiensis Eigenmann, 1917

Pimelodella laticeps Eigenmann, 1917

Pimelodella laticeps australis Eigenmann, 1917

Pimelodella macturki Eigenmann, 1912

Pimelodella megalops Eigenmann, 1912

Pimelodella metae Eigenmann, 1917

Pimelodella notomelas Eigenmann, 1917

Pimelodella serrata Eigenmann, 1917

Pimelodus clarias coprophagus Schultz, 1944

Pimelodus heteropleurus Eigenmann, 1912

Pimelodus ortmanni Eigenmann, 1911

Platysilurus barbatus Haseman, 1911

Pseudopimelodus albomarginatus Eigenmann, 1912

Pseudopimelodus villosus Eigenmann, 1912

Rhamdella montana Eigenmann, 1913

Rhamdia barbata Meek, 1907

Rhamdia branneri Haseman, 1911

Rhamdia branneri voulezi Haseman, 1911

Rhamdia cabreræ Meek, 1906

Rhamdia laticauda typhla Greenfield, Greenfield, and Woods, 1982

Rhamdia luigiana Villa, 1977

Rhamdia microps Eigenmann, 1917

Rhamdia nasuta Meek, 1909

Rhamdia oaxacæ Meek, 1902

Rhamdia regani Meek, 1907

Rhamdiopsis moreirai Haseman, 1911

Sovichthys abuelo Schultz, 1944

PLESIOPIDAE

Plesiops cephalotaenia Inger, 1955

PLOTOSIDAE

Copidoglanis novaeguineae Weber, 1908

Plotosus papuensis Weber, 1910

POECHILIIDAE

Acanthophaelus bifurcus Eigenmann, 1909

Acanthophaelus melanzonus Eigenmann, 1909

Alfaro acutiventralis Meek, 1912

Cnesterodon carnegiei Haseman, 1911

Diphyacantha chochoensis Henn, 1916

Gambusia alvarezi Hubbs and Springer, 1957

Gambusia bonita Meek, 1904

Gambusia caliensis Eigenmann and Henn, 1916 in Henn, 1916

Gambusia cana Meek and Hildebrand, 1913

Gambusia cascajalensis Meek and Hildebrand, 1913
Gambusia darienensis Meek and Hildebrand, 1913
Gambusia fasciata Meek, 1904
Gambusia heterochir Hubbs, 1957
Gambusia hurtadoi Hubbs and Springer, 1957
Gambusia infans Woolman, 1894
Gambusia latipunctata Meek and Hildebrand, 1913
Gambusia nigroventralis Eigenmann and Henn, 1912
Gambusia panuco Hubbs, 1926
Gambusia parismina Meek, 1912
Gambusia regani Hubbs, 1926
Gambusia turrubarensis Meek, 1912
Gambusia umbratilis Meek, 1912
Gambusia vittata Hubbs, 1926
Gambusia xanthosoma Greenfield, 1983
Gambusia yucatanica australis Greenfield, 1985
Girardinus garmani Eigenmann, 1903
Glaridichthys falcatus Eigenmann, 1903
Heterandria colombianus Eigenmann and Henn, 1912
Heterandria cubensis Eigenmann, 1903
Heterandria hasemani Henn, 1916
Heterandria lutzi Meek, 1902
Limia hollandi Henn, 1916
Limia rivasi Franz and Burgess, 1983
Neoheterandria elegans Henn, 1916
Phalloptychus eigenmanni Henn, 1916
Phallotorynus fasciolatus Henn, 1916
Platypoecilus tropicus Meek, 1907
Platypoecilus variatus Meek, 1904
Poecilia amates Miller, 1907
Poecilia caudata Meek, 1909
Poecilia fria Eigenmann and Henn, 1914
Poecilia latipunctata Meek, 1904
Poecilia pittieri Meek, 1912
Poecilia tenuis Meek, 1907
Poeciliopsis balsas Hubbs, 1926
Priapichthys annectens hesperis Hubbs, 1924
Priapichthys fosteri Hildebrand, 1925
Priapichthys letonai Hildebrand, 1925
Priapichthys olomina Meek, 1914
Priapichthys panamensis Meek and Hildebrand, 1916
Tomeurus gracilis Eigenmann, 1909
Toxus riddlei Eigenmann, 1903
Xiphophorus jalapae Meek, 1902

POMACANTHIDAE

Centropyge argi Woods and Kanazawa, 1951
Centropyge heraldi Woods and Schultz, 1953

Centropyge tutuila Jordan and Jordan, 1922
Holacanthus lunatus Blosser, 1909

POMACENTRIDAE

Abudefduf lorenzi Hensley and Allen, 1977
Chromis alta Greenfield and Woods, 1980
Chromis atripectoralis Welander and Schultz, 1951
Chromis limbaughi Greenfield and Woods, 1980
Chromis meridiana Greenfield and Woods, 1980
Chromis randalli Greenfield and Hensley, 1970
Chromis villadolidi Jordan and Tanaka, 1927
Chromis weberi Fowler and Bean, 1928
Chromis woodsi Bruner and Arnam, 1979
Eupomacentrus chrysus Bean, 1906
Eupomacentrus rocasensis Emery, 1972
Glyphisodon troscheli Gill, 1863
Microspathodon frontatus Emery, 1970
Pomacentrus cranei Herre, 1935
Pomacentrus jenkinsi Jordan and Evermann, 1903
Pomacentrus reidi Fowler and Bean, 1928
Pomacentrus stigma Fowler and Bean, 1928
Stegastes baldwini Allen and Woods, 1980

PROCHILODONTIDAE

Prochilodus maripicru Eigenmann, 1912

PSEUDOCROMIDAE

Nesiotes purpurascens De Vis, 1884
Pseudochromis aurea marshallensis Schultz, 1953
Pseudoplesiops sargenti Schultz, 1953

PSILORHYNCHIDAE

Psilorhynchus gracilis Rainboth, 1983

RAJIDAE

Pseudoraja fischeri Bigelow and Schroeder, 1954
Raja hollandi Jordan and Richardson, 1909

RHAMPHICHTHYIDAE

Gymnorhamphichthys hypostomus Ellis, 1912

RHINOCHIMAERIDAE

Neoharriotta carri Bullis and Carpenter, 1966

SALMONIDAE

Coregonus stanleyi Kendall, 1903
Leucichthys alpenae Koelz, 1924
Leucichthys kiya Koelz, 1921
Leucichthys reighardi Koelz, 1924
Oncorhynchus adonis Jordan and McGregor, 1925

Oncorhynchus ishikawae Jordan and McGregor, 1925
Oncorhynchus kawamurae Jordan and McGregor, 1925
Oncorhynchus rhodurus Jordan and McGregor, 1925
Salmo bathoecetor Meek, 1899
Salmo clarki declivifrons Meek, 1899
Salmo clarki jordani Meek, 1899
Salmo roosevelti Evermann, 1906
Salvelinus imbrius Jordan and McGregor, 1925
Salvelinus timagamiensis Henn and Rinkensbach, 1925

SCARIDAE

Cryptotomus crassiceps Bean, 1906
Scarus laevis Jordan and Evermann, 1903
Sparisoma rhomaleum Meek and Hildebrand, 1928

SCHILBIDAE

Physalia villiersi Boulenger, 1912

SCIAENIDAE

Menticirrhus focaliger Ginsburg, 1952
Ophioscion punctatissimus Meek and Hildebrand, 1925
Ophioscion venezuelae Schultz, 1945
Otolithes lini Herre, 1935
Pseudotolithus brunneolus Jordan and Richardson, 1909
Sciaena bathytatos Chao and Miller, 1975
Sciaena iharae Jordan and Metz, 1913
Sciaena trewavasae Chao and Miller, 1975
Stellifer colonensis Meek and Hildebrand, 1925
Stellifer melanocheir Eigenmann, 1917
Stellifer mindii Meek and Hildebrand, 1925
Umbrina tumacoensis Wilson, 1916

SCOMBRIDAE

Pneumatophorus peruanus Jordan and Hubbs, 1925

SCOPELARCHIDAE

Scopelarchus beebe Rofen, 1963
Scopelarchus sagax Rofen, 1963

SCORPAENIDAE

Brachirus bellus Jordan and Hubbs, 1925
Pterois ellioti Meek, 1897
Scorpaenopsis catocala Jordan and Evermann, 1903
Sebastapistes badiorufus Herre, 1935
Sebastodes clavilatus Starks, 1911

Sebastodes deani Starks, 1911
Sebastodes emphaeus Starks, 1911
Sebastodes ijimae Jordan and Metz, 1913
Sebastodes owstoni Jordan and Thompson, 1914
Sebastodes thompsoni Jordan and Hubbs, 1925
Thysanichthys evides Jordan and Thompson, 1914

SCYLIORHINIDAE

Pristiurus sauteri Jordan and Richardson, 1909

SERRANIDAE

Anthias conspicuus Heemstra, 1973
Anthias mooreanus Herre, 1935
Anthias woodsi Anderson and Heemstra, 1980
Bodianus stellatus Blosser, 1909
Centropristes melanus Ginsburg, 1952
Cephalopholis boninius Jordan and Thompson, 1914
Diplectrum labarum Rosenblatt and Johnson, 1974
Diplectrum pacificum Meek and Hildebrand, 1925
Entonanthias pascualis Jordan and Tanaka, 1927
Epinephelus elongatus Schultz, 1953
Epinephelus emoryi Schultz, 1953
Epinephelus ionthas Jordan and Metz, 1913
Epinephelus melanostigma Schultz, 1953
Epinephelus spilotoceps Schultz, 1953
Franzia ardens Jordan and Thompson, 1914
Liopropoma mowbrayi Woods and Kanazawa, 1951
Pikea aurora Jordan and Evermann, 1903
Pikea mexicanus atlanticus Schultz, 1958
Pikea mexicanus mexicanus Schultz, 1958
Plectranthias garrupellus Robins and Starck, 1961
Prionodes baldwini Evermann and Marsh, 1899
Serraniculus pumilio Ginsburg, 1952
Serranus chionaraia Robins and Starck, 1961
Serranus maytagi Robins and Starck, 1961
Symphysanodon berryi Anderson, 1970
Symphysanodon octoactinus Anderson, 1970
Ypsigramma lineata Schultz, 1953

SILURIDAE

Kryptopterus parvanalis Inger and Chin, 1959
Ompok sabanus Inger and Chin, 1959
Wallago maculatus Inger and Chin, 1959

SISORIDAE

Glyptothorax housei Herre, 1942

SOLEIDAE

Achirus fluviatilis Meek and Hildebrand, 1928
Soleonassus finis Eigenmann, 1912

SOLENOTOMIDAE

Solenostomus paegnius Jordan and Thompson, 1914

SPARIDAE

Archosargus oviceps Ginsburg, 1952
Diplodus bermudensis Caldwell, 1965

SQUALIDAE

Squaliolus sarmenti Noronha, 1926

STERNOPTYCHIDAE

Araiphos gracilis Grey, 1961
Polyipnus nuttingii Gilbert, 1905
Sonoda megalophthalma Grey, 1959

STERNOPYGIDAE

Sternopygus dariensis Meek and Hildebrand, 1916
Sternopygus pejateron Schultz, 1949

STICHAELIDAE

Anoplarchus purpureus archolepis Hubbs, 1926
Lumpenus longirostris Evermann and Goldsborough, 1907

SYNAPHOBANCHIDAE

Dysomma abyssale Kamohara, 1938
Synaphobranchus oregoni Castle, 1960

SYNGNATHIDAE

Bombonia uxorius Herre, 1935
Corythoichthys elerae Evermann and Seale, 1907
Corythoichthys polynotatus Dawson, 1977
Doryrhamphus melanopleura cooki Weed and Woods, 1953
Hippocampus brunneus Bean, 1906
Hippocampus fisheri Jordan and Evermann, 1903
Micrognathus suvensis Herre, 1935
Siphostoma brevicaudum Meek, 1904
Siphostoma eigenmanni Wilson, 1916
Siphostoma elcapitanense Meek and Hildebrand, 1914
Syngnathus mackayi nesiotus Herald, 1942
Syngnathus scovelli makaxi Herald and Dawson, 1972

SYNODONTIDAE

Synodus binotatus Schultz, 1953
Synodus ulae Schultz, 1953

TERAPONIDAE

Helotes lorentzi Weber, 1910
Terapon habbema Weber, 1910

TETRAODONTIDAE

Chonerhinus amabilis Roberts, 1982
Chonerhinus nefastus Roberts, 1982
Chonerhinus remotus Roberts, 1982
Chonerhinus silus Roberts, 1982
Spheroides asterias Blosser, 1909
Sphoeroides kendalli Meek and Hildebrand, 1928
Sphoeroides parvus Shipp and Yerger, 1969
Tetraodon kretamensis Inger, 1953

TRACHICHTHYIDAE

Hoplostethus occidentalis Woods, 1973
Paratrachichthys argyrophanus Woods, 1961

TRACHIPTERIDAE

Trachipterus fukuzakii Fitch, 1964

TRIACANTHODIDAE

Parahollardia schmidtii Woods, 1959

TRICHOMYCTERIDAE

Ochmacanthus flabelliferus Eigenmann, 1912
Pygidium alternatum Eigenmann, 1917
Pygidium banneai Eigenmann, 1912
Pygidium barbouri Eigenmann, 1911
Pygidium bogotense Eigenmann, 1912
Pygidium caliense Eigenmann, 1912
Pygidium chapmani Eigenmann, 1912
Pygidium conradi Eigenmann, 1912
Pygidium davisii Haseman, 1911
Pygidium dorsostriatum Eigenmann, 1917
Pygidium gracilior Eigenmann, 1912
Pygidium guianensis Eigenmann, 1909
Pygidium hasemani Eigenmann, 1914
Pygidium iheringi Eigenmann, 1917
Pygidium latistriatum Eigenmann, 1917
Pygidium paolence Eigenmann, 1917
Pygidium proops parahybae Eigenmann, 1918
Pygidium punctulatum piurae Eigenmann, 1922
Pygidium reinhardti Eigenmann, 1917
Pygidium santaeritae Eigenmann, 1918
Pygidium septentrionale Behre, 1928
Pygidium stellatum Eigenmann, 1918
Pygidium straminium Eigenmann, 1917
Pygidium striatum Meek and Hildebrand, 1913
Pygidium triguttatum Eigenmann, 1918
Pygidium verruculatum Eigenmann, 1917
Pygidium zonatum Eigenmann, 1918
Scleronema operculatum Eigenmann, 1917

Vandellia hasemani Eigenmann, 1918
Vandellia sanguinea Eigenmann, 1917

TRIGLIDAE

Bellator ribeiroi Miller, 1965
Chelidonichthys ischyryus Jordan and Thompson, 1914
Peristedion gilberti Jordan, 1922
Peristedion greyae Miller, 1967
Prionotus aspersus Meek and Hildebrand, 1928

TRIPTERYGIIDAE

Enneapterygius punctulatus Herre, 1935
Enneapterygius waigiensis Herre, 1935

UMBRIDAE

Novumbra hubbsi Schultz, 1929

URANOSCOPIDAE

Zalescopus satsumae Jordan and Hubbs, 1925
Zalescopus tosae Jordan and Hubbs, 1925

XENCONGRIDAE

Kaupichthys atlanticus Böhlke, 1956
Kaupichthys brachychirus Schultz, 1953

ZOARCIDAE

Allolepis hollandi Jordan and Hubbs, 1925
Lycodes tanakae Jordan and Thompson, 1914
Rhigophila dearborni DeWitt, 1962
Zestichthys tanakae Jordan and Hubbs, 1925

Acknowledgments

This catalogue represents part of the junior author's matching commitment to National Science Foundation Grant DEB 82-04071 that financed in part the technical help needed for curation of

Field Museum's type collection; Terry Grande and Robert Schmitz (both at FMNH) assisted in that regard. We would especially like to thank Ian Chavez (Antioch College, Yellow Springs, Ohio) and Roy Yanong (Yale University, New Haven, Connecticut) for their assistance with library research and diverse other tasks. Rosetta Arrigo (FMNH) wrote the computer program that was used to print the type list and generously assisted with various other computer-related needs. Lynn Hobbs (FMNH) drafted the illustration. We also thank Robert K. Johnson and Robert M. Timm (FMNH) for constructive criticism of the manuscript. Numerous colleagues assisted by providing advice, information on manuscript names, and help with other problems.

Literature Cited

- BARTHOLOMEW, J. C., ET AL., EDs. 1980. The Times Atlas of the World. Comprehensive Edition, 6th ed. Times Books, New York.
- BIOSIS. 1981. Serial Sources for the BIOSIS Data Base, 1980. BioSciences Informative Service, Philadelphia.
- COLLETTE, B. B., AND E. A. LACHNER. 1976. Fish collections in the United States and Canada. *Copeia*, 1976(3): 625-644.
- GREY, M. 1947. Catalogue of the type specimens of fishes in Chicago Natural History Museum. *Fieldiana: Zool.*, 32(3): 109-205.
- HENN, A. W. 1928. II. List of types of fishes in the collection of the Carnegie Museum on September 1, 1928. *Ann. Carnegie Mus.*, 19(4): 38-39.
- NELSON, J. S. 1984. *Fishes of the World*, 2nd ed. John Wiley and Sons, New York.
- NITECKI, M. H. 1980. Featured institution—Field Museum of Natural History. *ASC Newsletter*, 8(5): 61-70.
- ROBINS, C. R., ET AL. 1980. A List of Common and Scientific Names of Fishes from the United States and Canada, 4th ed. Special Publication No. 12, American Fisheries Society, Bethesda, Md.

Species Index

- abacopus*, *Glossogobius* 40, 96
abo, *Apogon* 10, 86
abramoides, *Astyanax* 12, 89
abelo, *Sovichthys* 80, 100
abyssale, *Dysomma* 33, 103
acanthogaster, *Deuterodon* 31, 90
acanthorhinus, *Ostichthys* 63, 97
acerus, *Paracrossochilus* 63, 94
achlamys, *Gazza* 38, 97
acutiventralis, *Alfaro* 7, 100
acutum, *Cichlasoma* 25, 92
adiposalis, *Pseudobagrus* 70, 88
adobe, *Agosia* 6, 93
adonis, *Oncorhynchus* 62, 101
aeneus costaricensis, *Astyanax* 12, 89
affinis littoralis, *Atherinops* 14, 87
 Pseudocheirodon 70, 91
 Rhombatractus 75, 98
afuerae multipunctatus, *Malacoctenus* 54, 97
aguadulce, *Galeichthys* 36, 87
ahlstromi, *Evermannella* 35, 95
akeleyi, *Barbus* 15, 93
albater, *Noturus* 61, 97
albeolus, *Astyanax* 12, 89
albomarginatus, *Pseudopimelodus* 71, 100
alepidota californiensis, *Chaenopsis* 21, 89
alestes borneensis, *Nematabramis* 60, 94
alfari, *Cichlasoma* 25, 92
alfredae, *Bryconamericus* 17, 90
aliata, *Stewardia* 82, 92
aliceae, *Argentina* 11, 87
alienus, *Cypselurus* 31, 95
alleni, *Tetragonopterus* 83, 92
alpenae, *Leucichthys* 52, 101
alpha, *Bryconamericus* 17, 90
alta, *Chromis* 24, 101
 Crenicichla 28, 92
alternatum, *Pygidium* 72, 103
alternus, *Leporinus* 51, 86
altipinna, *Pseudorasbora* 71, 94
 Rhoadsia 75, 91
altispinosa, *Crenicara* 28, 92
altum, *Sicydium* 79, 96
altus, *Acestorhynchus* 5, 89
alvarezii, *Gambusia* 36, 100
amabilis, *Chonerhinus* 24, 103
amates, *Poecilia* 67, 101
amaurus, *Bunocephalus* 19, 87
amazonae, *Bleptonema* 16, 90
amblops, *Felichthys* 36, 87
amphiloxus, *Hemicetopsis* 44, 89
 Nematobrycon 60, 91
anale, *Dorosoma* 33, 93
analisis australis, *Clinocottus* 27, 93
 Hemigrammus 44, 91
 Piabina 65, 91
anchovia, *Agoniatas* 6, 89
andersoni, *Corymbophanes* 28, 98
angelica, *Emblemaria* 34, 89
anguillicauda, *Liobagrus* 52, 88
anguillicaudatus tungting, *Misgurnus* 58, 93
anisitsi, *Aphyocharax* 9, 89
 Pterygoplichthys 72, 98
annae, *Cheirodon* 23, 90
annectens hesperis, *Priapichthys* 69, 101
anoplus, *Barbus* 15, 93
anpinensis, *Mugil* 58, 98
antillarum, *Ancylopsetta* 8, 88
antoncichi, *Cypselurus* 31, 95
anuli, *Pimephales* 66, 94
aokii, *Culter* 29, 94
aphanea, *Algansea* 7, 93
aphelochelil, *Protomyzon* 70, 97
aporia, *Gobiopsis* 41, 96
aporrhox, *Lepophidium* 51, 99
aquilus, *Lophiobagrus* 53, 88
archipelagicus, *Hemiramphus* 45, 96
arcus, *Leporinus* 51, 86
ardens, *Franzia* 36, 102
ardesiaca, *Nansenia* 59, 88
arenicola, *Anchovia* 8, 95
arenidens, *Solivomer* 80, 99
argentata, *Ilisha* 49, 93
argenteus, *Brycon* 19, 90
 Trachurus 84, 89
arge, *Upeneus* 84, 99
argi, *Centropyge* 21, 101
argyromus, *Myripristis* 58, 97
argyrophanus Paratrachichthys 64, 103
arleoi, *Anduzedoras* 8, 95
armatus australis, *Leptocottus* 52, 93
asper earli, *Exerpes* 36, 97
aspersus, *Prionotus* 69, 104
asterias, *Spheroides* 80, 103
astrinus, *Calliurichthys* 20, 88
atcafalayae vanhyningi, *Signalosa* 79, 93
ateleostus, *Nemaclinus* 60, 97
aterrimus, *Ctenogobius* 29, 96
 Hymenocephalus 47, 98
atherodon, *Optonurus* 63, 98
athi, *Barbus* 15, 93
atlanticus, *Kaupichthys* 50, 104
atracaudata, *Deuterodon* 31, 90
atratoensis, *Curimatus* 30, 93
atra, *Phrynelox* 65, 86
atrilatus, *Pseudaspius* 70, 94
atrimentatus, *Teuthis* 83, 86
atripectoralis, *Chromis* 24, 101
attenuatum, *Chiostoma* 23, 87
aubynei, *Pristella* 69, 91
aurata vallahica, *Cobitis* 27, 93
auratus, *Poecilobrycon* 68, 97
aurea marshallensis, *Pseudochromis* 71, 101
aurocaudatus, *Astyanax* 12, 89
aurocingulus, *Aparrius* 9, 95
aurora, *Pikea* 65, 102
australe, *Corydoras* 27, 88
 Pyrrhulina 74, 97
australis, *Ameiurus* 7, 97
avanhandavae, *Pimelodella* 65, 100
awani, *Aequidens* 5, 92
azonus, *Pleurogrammus* 67, 96
aztecus, *Notropis* 60, 94
azurifer, *Aequidens* 5, 92
azygia, *Farlowella* 36, 98
badia, *Macgregorella* 54, 96
badiorufus, *Sebastapistes* 78, 102
baldwini, *Hemipteronotus* 45, 97
 Prionodes 69, 102
 Stegastes 81, 101
balsanus, *Melaniris* 55, 87
balsas, *Poeciliopsis* 68, 101
balteata, *Vaimosa* 84, 96
banneai, *Pygidium* 72, 103
bantolanensis, *Barbus* 15, 93
barbata, *Rhamdia* 74, 100
barbatus, *Acheilognathus* 5, 93
 Platysilurus 67, 100
barbouri, *Pygidium* 72, 103
barnardi, *Petersius* 64, 91
barnesi, *Hypoatherina* 48, 87
bathoecetor, *Salmo* 77, 102
bathoiketes, *Porichthys* 69, 88
bathytatos, *Sciaena* 78, 102
bayeri, *Gymnothorax* 42, 99
beani, *Heros* 46, 92
beauforti, *Apogon* 10, 86
beebe, *Scopelarchus* 78, 102
behrae, *Brycon* 19, 90
belizensis, *Anchoviella* 8, 95
bellus, *Brachirus* 16, 102
beni, *Creagrutus* 28, 90
 Phenacogaster 64, 91
benjamini, *Entomocorus* 34, 87
bergi, *Pseudaspius* 70, 94
berlangai, *Opisthonema* 62, 93
bermudae, *Branchiostoma* 17, 88
bermudensis, *Antennarius* 9, 86
 Diplodus 32, 103
 Hemiramphus 45, 96
 Rhegma 75, 96
berndti, *Myripristis* 58, 97
beroe, *Lophiodes* 53, 98
berryi, *Symphysanodon* 82, 102
beta, *Bryconamericus* 17, 90
biaggi, *Heterenchelys* 46, 96
bifasciatus, *Cyprinodon* 31, 94
 Hyphessobrycon 48, 91
 Parodon 64, 99
bifrenatus, *Barbus* 15, 93
bifurcus, *Acanthophaelus* 4, 100
bigelowi, *Bathypterois* 15, 92
bikiniensis, *Gymnothorax* 42, 99
bilinearis, *Callechelys* 19, 99
bilineata, *Hasemania* 43, 91
bilineatus, *Pygocentrus* 73, 91

- bilunulata*, *Kuhlia* 50, 97
bimaculata, *Iridio* 49, 97
Nannacara 59, 92
bimaculatus novae, *Astyanax* 12, 89
paraguayensis, *Astyanax* 12, 89
binghami, *Ceratobranchia* 21, 90
biniplicata, *Myxine* 59, 99
binotatus, *Synodus* 82, 103
birtwistlei, *Gobiella* 41, 96
blennioides, *Characidium* 22, 90
blochi, *Corydoras* 27, 88
blohmi, *Epapterus* 34, 88
boehlkei, *Barbuligobius* 15, 95
Rhinodoras 75, 95
bogotense, *Pygidium* 72, 103
bolivarensis, *Hildadoras* 46, 95
boliviana, *Pimelodella* 65, 100
boliviense, *Cichlasoma* 25, 92
bonanox, *Coryphaenoides* 28, 98
bondi coppenamensis, *Corydoras* 27, 88
boninius, *Cephalopholis* 21, 102
bonita, *Gambusia* 37, 100
boquiae, *Hemibrycon* 43, 91
boquillae, *Cetopsorhamdia* 21, 100
borneensis, *Neostethus* 60, 99
bosquianus longimaxilla, *Chasmodes* 23, 88
bovalii, *Poecilocharax* 68, 91
bowditchae, *Myripristis* 58, 97
brachychirus, *Kaupichthys* 50, 104
branchiomelas, *Stolephorus* 82, 95
branneri voulezi, *Rhamdia* 74, 100
Rhamdia 74, 100
brasiliensis iporangensis, *Geophagus* 38, 92
itapicuruensis, *Geophagus* 38, 92
brevicaudum, *Siphostoma* 79, 103
breviceps, *Rivulus* 76, 86
brevior, *Auchenipterus* 14, 87
Chasmocranus 22, 100
brevipinnis, *Creagrutus* 28, 90
brevirostra, *Anchovia* 8, 95
brevis, *Hemigrammus* 44, 91
Thoracocharax 83, 95
brevispinis, *Lasiancistrus* 51, 98
Lophiobagrus 53, 88
breviventralis, *Alectis* 6, 89
broccae, *Spintherobolus* 80, 92
browni, *Moenkhausia* 57, 91
brucei, *Argentina* 11, 87
brunneolus, *Pseudolithus* 71, 102
brunneus, *Ectenias* 33, 93
Hippocampus 46, 103
Leporinus 51, 86
Rivulus 76, 86
bucca, *Varicus* 85, 96
bullisi, *Grimatocetes* 42, 86
Holocentrus 47, 97
burkei, *Careproctus* 20, 93
cabrerae, *Rhamdia* 74, 100
caetei, *Apistogramma* 10, 92
caliense, *Pygidium* 72, 103
caliensis, *Gambusia* 37, 100
californiensis elongatus, *Eucinos-tomus* 35, 95
calobrense, *Cichlasoma* 25, 92
cana, *Gambusia* 37, 100
candalarius, *Profundulus* 70, 94
canensis, *Acanthicus* 4, 98
capetensis, *Loricaria* 53, 98
capito, *Centrus* 21, 89
captiva, *Goodea* 42, 96
carens, *Barbus* 15, 93
caribbaea, *Trichopsetta* 84, 88
carletoni, *Leuciscus* 52, 94
carnegiei, *Cnesterodon* 27, 100
Pseudancistrus 70, 98
carpintis, *Neotroplus* 59, 92
carri, *Microdesmus* 56, 98
Neoharriotta 60, 101
carychroa, *Enchelycore* 34, 99
cascajalensis, *Bryconamericus* 17, 90
Gambusia 37, 101
catalepta, *Dermatocheir* 31, 90
catenatum, *Characidium* 22, 90
catocala, *Scorpaenopsis* 78, 102
cauae, *Roeboides* 76, 91
caucanum, *Characidium* 22, 90
caucanus, *Bryconamericus* 17, 90
Creagrutus 28, 90
Gephyrocharax 38, 90
Lasiancistrus 51, 98
caudata, *Poecilia* 67, 101
centrale, *Cichlasoma* 25, 92
cephalotaenia, *Plesiops* 67, 100
cerasina, *Curimata* 30, 93
chagresi, *Menidia* 56, 87
chamaizelus, *Bunocephalus* 19, 87
chancellori, *Odontorhamphus* 61, 96
chapmani, *Cyclopium* 30, 87
Pygidium 72, 103
chauliodus, *Aplatophis* 10, 99
chavesi, *Dorosoma* 33, 93
chiuhuahua, *Notropis* 60, 94
chimantanus, *Aequidens* 5, 92
Chasmocranus 22, 100
chinssuensis, *Pseudogobio* 71, 94
chionaraia, *Serranus* 79, 102
chobensis, *Aplocheilus* 10, 86
chocoensis, *Bryconamericus* 17, 90
Diphyacantha 32, 100
Gephyrocharax 39, 90
chosenicus, *Rhodeus* 75, 94
chryseres, *Myripristis* 58, 97
chrysonemus, *Pseudupeneus* 71, 99
chrysus, *Eupomacentrus* 35, 101
cismontanus, *Bryconamericus* 18, 90
citurensis, *Oxyloricaria* 63, 98
clarias coprophagus, *Pimelodus* 66, 100
clarkae, *Adontosternarchus* 5, 87
clarkei, *Scopelogadus* 78, 99
clarki declivifrons, *Salmo* 77, 102
jordani, *Salmo* 77, 102
Euleptoeleotris 35, 95
clavilatus, *Sebastodes* 78, 102
claytonii, *Gobius* 41, 96
cliffordpopei, *Gobius* 42, 96
coccogenis, *Serrasalmus* 79, 92
colombianus, *Bunocephalus* 19, 87
Hemibrycon 43, 91
Heterandria 46, 101
colombiensis, *Thyrina* 83, 87
colonensis, *Stellifer* 81, 102
combata, *Antigonia* 9, 89
compressus, *Hemigrammus* 44, 91
Odontognathus 61, 93
Rivulus 76, 86
conchos, *Catostomus* 21, 89
conklini, *Sphagebranchius* 80, 99
conradi, *Pygidium* 72, 103
conserialis, *Petersius* 64, 91
consocium, *Chirostoma* 24, 87
conspicuum, *Anthias* 9, 102
conventus, *Bryconamericus* 18, 90
copei, *Anduzedoras* 9, 95
corethromycter, *Luciobrotula* 54, 99
coriparoides, *Gobio* 41, 94
corlettei, *Gnatholepis* 40, 96
corniger, *Ogcocephalus* 61, 99
corumbae, *Heterogramma* 46, 92
costaricensis, *Gobiesox* 41, 95
cranei, *Pomacentrus* 68, 101
Thalassoma 83, 97
crassiceps, *Cryptotomus* 29, 102
crenatus, *Stethaprion* 82, 92
crenolepis, *Allanetta* 7, 87
cubensis, *Fundulus* 36, 94
Heterandria 46, 101
cumberlandensis, *Phoxinus* 65, 94
cuprea, *Siphamia* 79, 87
curviventris, *Psectrogaster* 70, 93
cyaneus, *Mugilops* 58, 87
cycloidea, *Ancylapsetta* 8, 88
cyclolepis, *Parecbasis* 64, 91
cyclopus santanderensis, *Astroblepus* 11, 87
cylindricus, *Hemigrammus* 44, 91
cyri orientalis, *Chondrostoma* 24, 94
daguae, *Astyanax* 12, 89
Gobius 42, 96
Hemiancistrus 43, 98
Joturus 50, 98
dariensis, *Gambusia* 37, 101
dariensis, *Hemibrycon* 43, 91
Oxyloricaria 63, 98
Parodon 64, 99
Sternopygus 81, 103
davisi, *Pygidium* 72, 103
dawsoni, *Echiodon* 33, 89
dayi, *Decapterus* 31, 89
deani, *Sebastodes* 78, 102
dearborni, *Cyprinodon* 31, 94
Rhigophila 75, 104
decemlineatus, *Awaous* 14, 95
declivirostris, *Ogcocephalus* 62, 99
decoratus, *Iridio* 49, 97
decurrens, *Bryconamericus* 18, 90
delicata, *Tagusa* 82, 88
demerarae, *Auchenipterus* 14, 87
dentatus, *Bryconamericus* 18, 90

- depressirostris*, *Pseudorasbora* 71, 94
depressus, *Bunocephalus* 19, 87
derbyi, *Plecostomus* 67, 98
deuterodonoides caudovittatus, *Bryconamericus* 18, 90
Bryconamericus 18, 90
devenanzii, *Adontosternarchus* 5, 87
diaboli, *Dionda* 32, 94
dialepturus, *Cheirodon* 23, 90
diazii, *Zoogoneticus* 86, 96
dilatus, *Spirinchus* 81, 99
diquensis, *Bryconamericus* 18, 90
discus, *Acestridium* 4, 98
dispar, *Pempheris* 64, 99
distethommatum, *Callionymus* 19, 88
dorsatum, *Cichlasoma* 25, 92
dorsocellata, *Crenicichla* 28, 92
dorsostriatum, *Pygidium* 72, 103
dubius, *Bathytrois* 15, 92
duopunctata, *Aequidens* 5, 92
duragenys, *Hyphessobrycon* 48, 91
eбелиngi, *Gonostoma* 42, 93
ebisui, *Malacotenus* 54, 97
ectenurus, *Leptocephalus* 52, 93
ecuadoriensis, *Hyphessobrycon* 48, 91
Leporinus 51, 86
Parodon 64, 99
edwardraneyi, *Notropis* 60, 94
efatensis, *Gymnothorax* 42, 99
eidolon, *Psychichthys* 71, 92
eigenmannia, *Anchovia* 8, 95
eigenmanni, *Chaenothorax* 22, 88
Cheirodon 23, 90
Cichlasoma 25, 92
Fitzroyia 36, 97
Parapelecus 63, 94
Phalloptychus 64, 101
Pimelodella 66, 100
Siphostoma 79, 103
elcapitanense, *Siphostoma* 80, 103
elegans erloli, *Gibbonsia* 39, 93
montereyensis, *Gibbonsia* 39, 93
velifera, *Gibbonsia* 39, 93
Ammocryptocharax 7, 89
Iridio 49, 97
Neoheterandria 60, 101
elerae, *Corythoichthys* 28, 103
elliotti, *Pterois* 71, 102
Thorichthys 83, 92
ellisae, *Corydoras* 27, 88
elongata, *Anchovia* 8, 95
elongatus, *Bramocharax* 17, 90
Carpiodes 20, 89
Epinephelus 34, 102
elymus aphrozoicus, *Atherion* 14, 87
freyi, *Atherion* 14, 87
emoryi, *Epinephelus* 34, 102
emphaeus, *Sebastodes* 78, 102
engraulis, *Hemiculterella* 44, 94
ensifer, *Gladiogobius* 39, 96
Holocentrus 47, 97
eos, *Hyphessobrycon* 48, 91
eques, *Cheirocerus* 23, 100
erimyzonops, *Dionda* 32, 94
erionna, *Scolopsis* 78, 99
eristigma, *Belligobio* 16, 94
erythrozonus, *Hemigrammus* 44, 91
erythrurus, *Aphyocharax* 9, 89
Poecilobrycon 68, 97
essequibensis, *Asiyanax* 12, 89
Cynopotamus 31, 90
Leptoglanis 52, 100
ethelwynnae, *Gobiocichla* 41, 92
eugenius, *Quisquilius* 74, 96
evermanni, *Atherina* 14, 87
Cichlasoma 25, 92
Diapterus 32, 95
Ophichthys 62, 99
evides, *Encaetra* 34, 95
Hemipteronotus 45, 97
Thysanichthys 83, 102
evurus, *Etelis* 35, 98
exilis, *Trachelyichthys* 83, 88
falcatus, *Glaridichthys* 40, 101
fasciata, *Gambusia* 37, 101
fasciatus, *Gastromyzon* 38, 97
fasciata, *Crossostoma* 29, 93
fasciolatus, *Phallotorynus* 64, 101
ferdebueni, *Poblana* 67, 87
fijiensis, *Halichoeres* 43, 97
filamentosa latiuscula, *Loricaria* 53, 98
seminuda, *Loricaria* 53, 98
filamentosus, *Macropodus* 54, 88
fimbriata, *Loricaria* 53, 98
finis, *Soleonassus* 80, 102
fischeri, *Pseudoraja* 71, 101
fisheri, *Hippocampus* 46, 103
Labeo 50, 94
Trachycorystes 84, 88
fitzsimonsi, *Barbus* 15, 93
flabelliferus, *Ochmacanthus* 61, 103
flammeus, *Mulloidops* 58, 98
flavater, *Noturus* 61, 97
floridae, *Branchiostoma* 17, 88
fluviatilis, *Achirus* 5, 102
Disparichthys 32, 89
Gobiesox 41, 95
Leptophilypnus 52, 95
Neetroplus 59, 92
focaliger, *Menticirrhus* 56, 102
fontinalis, *Cyprinodon* 31, 94
forlonensis, *Nototropis* 60, 94
formosa, *Rineloricaria* 76, 98
formosae, *Liza* 53, 98
formosanus, *Caranx* 20, 89
Rhinogobius 75, 96
Scomberoides 78, 89
fosteri, *Priapichthys* 69, 101
fowleri, *Cephalosilurus* 21, 100
fragilis, *Scepterius* 78, 87
franciscensis, *Fowlerina* 36, 90
franciscoensis, *Phenacogaster* 65, 91
freemani, *Pristipomoides* 69, 98
frenata, *Brachyglanis* 17, 100
frenatus, *Astroblepus* 11, 87
Rivulus 76, 86
fria, *Poecilia* 67, 101
frontale, *Cichlasoma* 25, 92
frontatus, *Microspathodon* 57, 101
fukuzakii, *Trachipterus* 83, 103
fulvus, *Gobiesox* 41, 95
fundicola, *Suruga* 82, 96
furcatus, *Oreias* 63, 93
fuscoguttatus, *Uropterygius* 84, 99
fuscolineata, *Siphamia* 79, 87
fuscomaculata, *Rabula* 74, 99
gabonensis, *Grasseichthys* 42, 97
gabunensis, *Ischnomembras* 50, 87
galusdae, *Cheirodon* 23, 90
garmani, *Girardinus* 39, 101
garrupellus, *Plectranthias* 67, 102
geniornatus, *Cryptocentrus* 29, 96
georgei, *Argentina* 11, 87
gigas, *Diaphus* 32, 99
Malacotenus 54, 97
gilberti, *Careproctus* 20, 93
Cololrhynchus 27, 98
Formosania 36, 94
Peristedion 64, 104
gillii, *Curimatus* 30, 93
gimbeli, *Porotergus* 69, 87
glandulifer, *Diaphus* 32, 99
glaucum, *Stizostedion* 82, 100
gloverensis, *Triathalassothia* 84, 88
gracilicauda, *Gymnapogon* 42, 86
gracilior, *Pygidium* 72, 103
gracilis, *Acheilognathus* 5, 93
Araiphos 11, 103
Psilorhynchus 71, 101
Tomeurus 83, 101
grammica, *Aphyodite* 10, 89
grammodes, *Cichlasoma* 25, 92
granadense, *Cichlasoma* 25, 92
grandis, *Astyanax* 12, 89
Pomadasys 68, 96
grandisquamis, *Amblypharyngodon* 7, 93
granti, *Leporinus* 51, 86
greenei, *Notropis* 61, 94
greenfieldi, *Acanthemblemaria* 4, 89
greyae, *Bathylagus* 15, 88
Gillellus 39, 94
Peristedion 64, 104
greyi, *Bembrops* 16, 100
griffini, *Pimelodella* 66, 100
grimaldii, *Hymenocephalus* 47, 98
grinnelli, *Physiculus* 65, 98
griseus, *Loricaria* 53, 98
grixalvii microscens, *Astroblepus* 11, 87
grosvenori, *Bryconamericus* 18, 90
guaporensis, *Aequidens* 5, 92
Astyanax 12, 89
Metynnis 56, 91
Microschemobrycon 57, 91
guianensis, *Acestrorhynchus* 5, 89
Ageneiosus 6, 86
Astyanax 12, 89
Brachyhalcinus 17, 90
Pygidium 73, 103
Stolephorus 82, 95
guuja, *Thyrina* 83, 87

- gymnocephalus*, *Dormitator* 33, 95
gymnodontus, *Psolidodon* 70, 91
gymnogaster lagoichthys, *Loricaria* 53, 98
Loricaria 53, 98
gymnogenys, *Asryanax* 13, 89
gymnotus, *Porotergus* 69, 87

habbema, *Terapon* 83, 103
hankinsoni, *Hybognathus* 47, 94
harringtonensis, *Leptocephalus* 52, 93
harrisoni, *Poecilobrycon* 68, 97
hartwegi, *Cichlasoma* 25, 92
hasemani, *Apareiodon* 9, 99
Asryanax 13, 89
Farlowella 36, 98
Heterandria 46, 101
Pimelodella 65, 100
Pygidium 73, 103
Sternarchus 81, 87
Vandellia 84, 104
hastatus, *Odontostilbe* 61, 91
hedricki, *Cichlasoma* 25, 92
helleri, *Anchoiella* 8, 95
Barbus 15, 94
Hemibrycon 43, 91
hemigrammus, *Aphyocheirodon* 10, 89
Asiphonichthys 11, 89
hemimurus, *Plecostomus* 67, 98
henni, *Brycon* 19, 90
Bryconamericus 18, 90
hepsetus colonensis, *Anchoa* 7, 95
heraldi, *Centropyge* 21, 101
herwerdenii, *Eleotris* 33, 95
heterandria, *Pseudocorynopoma* 71, 91
heterochir, *Gambusia* 37, 101
heterodon, *Eleotris* 33, 95
Holesthes 47, 91
heterolepis, *Cynoglossus* 30, 93
heteromorphus, *Corydoras* 27, 88
heteropleurus, *Pimelodus* 66, 100
heterostomus, *Probolodus* 70, 91
heterura, *Compsura* 27, 90
heterurus, *Asryanax* 13, 89
hiatti, *Acanthoplesiops* 4, 86
higmani, *Lonchopisthus* 53, 99
hildebrandi, *Hyporhamphus* 49, 96
Roeboides 76, 91
Sicydium 79, 96
hingi, *Homaloptera* 47, 97
hoignei, *Pterolebias* 71, 86
hollandi, *Allolepis* 7, 104
Areliscus 11, 93
Imparfinis 49, 100
Limia 52, 101
Myctophum 58, 99
Raja 74, 101
Serrasalmo 79, 92
Spinibarbus 80, 94
holmiae, *Rivulus* 76, 86
hondae, *Pseudoperilampus* 71, 94
horiae, *Vaimosa* 84, 96
housei, *Glyptothorax* 40, 102

hubbsi hubbsi, *Malacoctenus* 55, 97
polyporosus, *Malacoctenus* 54, 97
Novumbra 61, 104
hudsoni, *Callionymus* 20, 88
humeralis gracilior, *Serrasalmo* 79, 92
hurtadoi, *Gambusia* 37, 101
hyltoni, *Emblemaria* 34, 89
hypesson, *Bryconamericus* 18, 90
hyposagmatus, *Ophichthus* 62, 99
hypostomus, *Gymnorhamphichthys* 42, 101

iguassuensis, *Crenicichla* 29, 92
iharae, *Sciaena* 78, 102
iheringi, *Microglanis* 57, 100
Pygidium 73, 103
ijimae, *Gnathopogon* 40, 94
ijimae, *Sebastodes* 79, 102
Synchiropus 82, 89
ijimai, *Astronesthes* 12, 87
illecebrosus, *Alburnops* 6, 93
illustris, *Eumegistus* 35, 88
imbrius, *Salvelinus* 77, 102
imswe, *Varicus* 85, 96
incompta, *Curimata* 30, 93
inconspicua, *Apistogramma* 10, 92
inequalis, *Glandulocauda* 39, 91
inermis, *Platophrys* 66, 88
infans, *Gambusia* 37, 101
insularum cedroscensis, *Atherinops* 14, 87
guadalupae, *Atherinops* 14, 87
Atherina 14, 87
integripinnis zacae, *Paraclinus* 63, 97
intermedius, *Gephyrocharax* 39, 90
Squalogadus 81, 98
interruptus ibicuhiensis, *Cheirodon* 23, 90
intonus, *Ernstichthys* 35, 87
ionthas, *Epinephelus* 34, 102
iota, *Hemigrammus* 44, 91
irinae, *Pseudanos* 70, 86
ischanus, *Icticus* 49, 99
ischyrus, *Chelidonichthys* 23, 104
ishikawae, *Gnathopogon* 40, 94
Oncorhynchus 62, 102
ishikawai, *Caranx* 20, 89
issororoensis, *Curimatus* 30, 93
isthmensis, *Eleotris* 33, 95
itapicuruensis, *Apareiodon* 9, 99
Pimelodella 65, 100

jaguarensis, *Crenicichla* 29, 92
jalapae, *Xiphophorus* 85, 101
japonica, *Eviota* 36, 96
jauruensis, *Farlowella* 36, 98
jeb, *Calamopteryx* 19, 99
jenkinsi, *Pomacentrus* 68, 101
jimi, *Parotocinclus* 64, 98
joaquinae, *Puntius* 72, 94
johnsoni, *Cirriacaula* 26, 99
jordani, *Chirostoma* 24, 87
Conger 27, 93
Labeo 50, 94

jordani, *Lampanyctus* 50, 99
Pantosteus 63, 89
Trachinotus 83, 89
jusanensis, *Achahara* 4, 93

kabia, *Melanotaenia* 56, 98
kabiliensis, *Brachygnathus* 17, 95
kanazawai, *Dicrolene* 32, 99
kanganamanensis, *Arius* 11, 87
kasawae, *Iburiella* 49, 86
katsukii, *Stellistius* 81, 96
kawamurae, *Oncorhynchus* 62, 102
keithi, *Mastacembelus* 55, 98
kendalli, *Sphoeroides* 80, 103
kennedyi, *Hemigrammus* 45, 91
kesleri banaticus, *Gobio* 41, 94
kikuchii, *Phoxinus* 65, 94
kiyi, *Leucichthys* 52, 101
knightii, *Gnatholepis* 40, 96
koelzi, *Crossochir* 29, 95
komp, *Asryanax* 13, 89
koragensis, *Glossogobius* 40, 96
kretamensis, *Tetraodon* 83, 103
kulambangrae, *Petrosirtus* 64, 88
kumae, *Monomitopus* 58, 99
kumperae, *Ancylorsetta* 8, 88
kuroiwa, *Tridentiger* 84, 96
kurumeus, *Rhodeus* 75, 94
kynosemeum, *Porichthys* 69, 88

labarcae, *Chirostoma* 24, 87
labarum, *Diplectrum* 32, 102
labiosus, *Carpiodes* 20, 89
lacortei, *Nematobrycon* 60, 91
lafa, *Ambassis* 7, 89
lamprotaenia, *Anchoa* 8, 95
lanceolatus, *Rivulus* 76, 86
landoni, *Pterobrycon* 71, 91
langi, *Alestes* 6, 89
laterale, *Characidium* 22, 90
laticauda typhla, *Rhamdia* 74, 100
laticeps australis, *Pimelodella* 66, 100
Pimelodella 66, 100
Schizodontopsis 78, 86
latidens, *Astroblepus* 11, 87
Landonia 50, 91
latifasciatus, *Eleotris* 34, 95
latipunctata, *Gambusia* 37, 101
Poecilia 67, 101
latistriatum, *Pygidium* 73, 103
latus, *Diaphus* 32, 99
lauia, *Scarus* 78, 102
leavellii, *Ctenogobius* 29, 96
lentinus, *Doras* 33, 95
Labrisomus 50, 97
lepidogaster, *Gastromyzon* 38, 97
lepidura hasemani, *Moenkhausia* 57, 91
leporhinus, *Hemidoras* 44, 95
leptocaulus, *Apogon* 10, 86
leptorhynchus, *Sternarchus* 81, 87
lermae, *Skiffia* 80, 96
letonai, *Priapichthys* 69, 101
leucomelas, *Chaetostomus* 22, 98
leucostigma, *Gymnothorax* 43, 99

- levis*, *Hemieleotris* 44, 95
lignus, *Hypsoblennius* 49, 88
limbaughi, *Chromis* 24, 101
lineata, *Ypsigrama* 85, 102
lini, *Otolithes* 63, 102
linnelli, *Leptodoras* 52, 95
liolepis, *Euctenogobius* 35, 96
lisa, *Menidia* 56, 87
lithoides, *Lithoxus* 53, 98
lithurgicus, *Ancistrus* 8, 98
longifilis, *Gnathopogon* 40, 94
longior, *Chasmocranus* 23, 100
longirostris, *Lumpenus* 54, 103
lorentzi, *Helotes* 43, 103
lorenzi, *Abudefduf* 4, 101
luigiana, *Rhamdia* 74, 100
lunatus, *Holacanthus* 46, 101
lutzi, *Heterandria* 46, 101
luzonensis, *Aplocheilichthys* 10, 86
lyolepis, *Stolephorus* 82, 95
lyriformis, *Agmus* 6, 87

mabahiss, *Vinciguerra* 85, 100
macilentus, *Hemicetopsis* 44, 89
mackayi nesiotes, *Syngnathus* 82, 103
macmasteri, *Apistogramma* 10, 92
macrocephalus catherinae, *Mniertes* 57, 97
macrolepidota sabana, *Hampala* 43, 94
macrolepis, *Eleotris* 34, 95
Heterocharax 46, 91
Roeboides 76, 91
macrosquamis, *Sargocentron* 77, 97
macturki, *Pimelodella* 66, 100
maculatus, *Wallago* 85, 102
maculipinnis, *Pterolebias* 72, 86
maculocephala, *Cascadura* 21, 88
maculosus, *Aspidoras* 11, 88
Tylobronchus 84, 92
madeirae, *Cheirodon* 23, 90
madeirensis, *Ageneiosus* 6, 86
maderensis, *Diplogonurus* 32, 95
magdalenae, *Astyanax* 13, 90
Creagrutus 28, 90
Hoplosternum 47, 88
Rivulus 76, 86
Roeboides 77, 91
Thoracocharax 83, 95
Xiliphenis 85, 87
magdalenensis, *Bryconamericus* 18, 90
majimae, *Gnathopogon* 40, 94
majua majua, *Jenkinsia* 50, 93
woodsii, *Jenkinsia* 50, 93
malaris, *Clarias* 26, 93
malayana, *Antigonia* 9, 89
malayanus, *Myxus* 59, 98
malekulae, *Ctenogobius* 29, 96
malua, *Boroda* 16, 95
margaritae mexicanus, *Malacotenus* 55, 97
marginatus, *Hemigrammus* 45, 91
Nannostomus 59, 97
mariae, *Acanthopthalmus* 4, 93

mariae, *Aequidens* 5, 92
Sternarchus 81, 87
marijeanae, *Gobiesox* 41, 95
marionae, *Astyanax* 13, 90
Bathytyphlops 16, 92
maripicru, *Prochilodus* 70, 101
marmoratus, *Ageneiosus* 6, 86
marquesensis, *Sardinella* 77, 93
marshallensis, *Rabula* 74, 99
martini, *Salarias* 77, 88
maxillaris, *Hasemanian* 43, 91
mayoloi, *Hemiancistrus* 43, 98
maytagi, *Serranus* 79, 102
medirastre, *Opisthonema* 62, 93
meeki, *Cichlasoma* 26, 92
Cyprinodon 31, 94
Holocentrus 47, 97
Roeboides 77, 91
Thyrina 83, 87
megalopsis, *Gastromyzon* 38, 97
Monolene 58, 88
megalophtalma, *Sonoda* 80, 103
megalops, *Pimelodella* 66, 100
megalopter, *Megalampodus* 55, 91
megalostictus, *Phenacogaster* 65, 91
megalostomus, *Hysteronotus* 49, 91
melampodus, *Apogonichthys* 10, 86
melandetus, *Odontostilbe* 61, 91
melanocheir, *Gephyrocharax* 39, 90
Stellifer 81, 102
melanogenys, *Glandulocauda* 39, 91
melanopleura cooki, *Doryrhamphus* 33, 103
Glandulocauda 39, 91
melanopleurus, *Hyphessobrycon* 45, 91
melanopter, *Hemigrammus* 48, 91
Hyporhamphus 49, 96
melanopus, *Leiogaster* 51, 88
melanostigma, *Epinephelus* 35, 102
melanotus, *Aphyocharax* 10, 89
melanura, *Hasemanian* 43, 91
melanus, *Centropomus* 21, 102
melanzonus, *Acanthophtalmus* 4, 100
Creagrutus 28, 90
melas, *Ancistrus* 8, 98
Brachyglanis 17, 100
meridiana, *Chromis* 24, 101
merretti, *Polyacanthonotus* 68, 99
metae, *Astyanax* 13, 90
Charax 22, 90
Curimatus 30, 93
Moenkhausia 57, 91
Pimelodella 66, 100
metzi, *Gibbonsia* 39, 93
mexicana campi, *Signalosa* 79, 93
Aztecula 14, 93
mexicanus atlanticus, *Pikea* 65, 102
mexicanus, *Pikea* 65, 102
Ameiurus 7, 97
Gobiesox 41, 95
meyeri, *Iridio* 50, 97
mezquital, *Chirostoma* 24, 87

microctenus, *Ancylosetta* 8, 88
microdon, *Cheirodon* 23, 90
Loricaria 53, 98
microlepis, *Acanthocharax* 4, 89
Astyanax 13, 90
micropoetus, *Hemidoras* 44, 95
microps, *Imparfinis* 49, 100
Pogoneleotris 68, 95
Rhamdia 74, 100
micropterus, *Megalampodus* 55, 91
micropunctatus, *Lampanyctus* 50, 99
microstictus, *Phenacogaster* 65, 91
microstomus, *Carpoides* 21, 89
Hemidoras 44, 95
Iridio 50, 97
Oligobrycon 62, 91
milleri, *Cichlasoma* 26, 92
minacae, *Gila* 39, 94
mindii, *Microeleotris* 56, 95
Stellifer 81, 102
miniatus, *Zoogoneticus* 86, 96
minimus, *Hyphessobrycon* 48, 91
Nannostomus 59, 97
minor, *Hyphessobrycon* 48, 91
minutulus, *Synchiropus* 82, 89
minutus, *Archicheir* 11, 97
Evorthodus 36, 96
Hemicetopsis 44, 89
Hemigrammus 45, 91
Microbrycon 56, 91
Microgenys 56, 91
mirini, *Imparfinis* 49, 100
misakius, *Chasmias* 22, 95
mittelli diaphana, *Anchoa* 8, 95
miyakamii, *Caranx* 20, 89
mizolepis fukien, *Misgurnus* 57, 93
grangeri, *Misgurnus* 57, 93
modestus, *Pseudaspis* 70, 94
moenkhausii, *Poecilurichthys* 68, 91
mohoity yunnan, *Misgurnus* 57, 93
mojarra, *Cichlasoma* 26, 92
mola, *Metynnus* 56, 91
mollispinus mollispinus, *Lepidomeda* 51, 94
montana, *Rhamdella* 74, 100
montanum, *Sicydium* 79, 96
monticola, *Algansea* 7, 93
montium, *Rivulus* 76, 86
mooreanus, *Anthias* 9, 102
Myripristis 58, 97
morawhannae, *Curimatus* 30, 93
moreirai, *Rhamdopsis* 75, 100
morii, *Sarcocheilichthys* 77, 94
mossambicus, *Chromis* 27, 92
Clarias 26, 93
moibrayii, *Grammonus* 42, 99
Liopropoma 52, 102
Rhinogobius 75, 96
mucronatus, *Astyanax* 13, 90
multiporosus, *Labrisomus* 50, 97
multiradiatus, *Characodon* 22, 96
multispinus, *Ctenopoma* 29, 86
mundus, *Orestias* 63, 94
munitus, *Noturus* 61, 97
muskoka, *Notropis* 61, 94

- mutator*, *Astyanax* 13, 90
myersi, *Amblygobius* 7, 95
Ekemblemaria 33, 89
myriaster, *Porichthys* 69, 88
myxodermus, *Ctenogobius* 29, 95
- nantoenis*, *Liobagrus* 52, 88
nasus, *Cetopsorhamdia* 21, 100
nasuta, *Rhamdia* 74, 100
nasutus, *Acestrorhynchus* 5, 89
Astyanax 13, 90
nazas, *Nototropis* 60, 94
nefastus, *Chonerhinus* 24, 103
nemacheir, *Nannorhamdia* 59, 100
neumanni, *Fundulus* 36, 94
nicholsi, *Blennius* 16, 88
nigra, *Emblemaria* 34, 89
nigrescens, *Pseudancistrus* 70, 98
nigripinnis woodsi, *Cotylis* 28, 95
nigritum, *Cichlasoma* 26, 92
nigrofasciatus, *Apogon* 10, 86
nigrolineatus, *Pseudauchenipterus* 70, 88
nigroventralis, *Gambusia* 37, 101
niveilatus, *Xyrichtys* 85, 97
norae, *Gibbonsia* 39, 93
notemigonoides, *Astyanax* 13, 90
notomelas, *Cheirodon* 23, 90
Pimelodella 66, 100
notosipilus, *Hemiodoras* 44, 95
notropoides, *Creagrutus* 28, 90
nouhuysi, *Atherinichthys* 14, 87
novae, *Bryconamericus* 18, 90
novaeaguineae, *Copidoglanis* 27, 100
novemmaculosus, *Alticus* 7, 88
nox, *Arius* 11, 87
nudus, *Gerhardinus* 39, 96
nuttingii, *Polyipnus* 68, 103
- oatesii*, *Barbus* 15, 94
Mastacembelus 55, 98
oaxacae, *Fundulus* 36, 94
Rhamdia 74, 100
obeyense, *Etheostoma* 35, 100
obliquus, *Lamprologus* 50, 92
obscurus, *Brycon* 19, 90
occidental, *Hoplostethus* 47, 103
Lepomis 51, 89
Roeboides 77, 92
Starksia 81, 97
ocellatum, *Mylossoma* 58, 91
ocellatus, *Callogobius* 20, 95
Poecilobrycon 68, 97
octoactinus, *Symphysanodon* 82, 102
oculomaculatus, *Gnathodentex* 40, 98
ohiensis, *Alosa* 7, 93
oligolinx, *Rhizoprionodon* 75, 89
olomina, *Priapichthys* 69, 100
opalescens, *Ctenogobius* 29, 96
operculatum, *Scleronema* 78, 103
orbiculatus, *Astronotus* 12, 92
orcuttii, *Phoxinus* 65, 94
oregona, *Draconetta* 33, 95
- oregoni*, *Synaphobranchus* 82, 103
orientalis americana, *Epinnula* 35, 95
pacifica, *Epinnula* 35, 95
Dasyscopelus 31, 99
ornatum gephyrum, *Cichlasoma* 26, 92
ortholepis, *Bryconamericus* 18, 90
orthonops, *Hemiodus* 45, 96
orthonotus, *Cyprinodon* 31, 94
orthus, *Hemigrammus* 45, 91
ortmanni, *Heterogramma* 46, 92
Pimelodus 66, 100
osakae, *Netuma* 60, 87
osgoodi, *Bryconamericus* 18, 90
Copeina 27, 97
Vaimosa 84, 96
oshimai, *Caranx* 20, 89
othonops, *Hemicetopsis* 44, 89
ovalaua, *Atherina* 14, 87
oviceps, *Archosargus* 11, 103
owstoni, *Alepocephalus* 6, 86
Sebastodes 79, 102
oxyrhynchus desotoi, *Acipenser* 5, 86
- pachacuti*, *Bryconamericus* 18, 90
pacificum, *Diplectrum* 32, 102
pacificus, *Mionorus* 57, 87
paegnius, *Solenostomus* 80, 103
palustris, *Beirabarbatus* 16, 94
paramensis daguae, *Hyphessobrycon* 48, 91
Eucenogobius 35, 96
Microeleotris 56, 95
Neotroplus 60, 92
Priapichthys 69, 101
panarcys, *Notropis* 61, 94
paranostictus, *Ogcocephalus* 62, 99
panuco, *Gambusia* 37, 101
paolence, *Pygidium* 73, 103
papillatus, *Hoplomyzon* 47, 87
papillifer, *Hemipimelodus* 45, 87
papilliferus, *Spintherobolus* 81, 92
papuensis, *Plotosus* 67, 100
paradiseus, *Gobiesox* 41, 95
paraguayensis, *Aphyocharax* 10, 89
Bleptonema 16, 90
Odontostilbe 61, 91
parahybae, *Cheirodon* 23, 90
paranaense, *Cichlasoma* 26, 92
paranahybae, *Astyanax* 13, 90
parismina, *Gambusia* 37, 101
parnaguae, *Hemiodus* 45, 96
parva, *Anchovia* 8, 95
Liza 53, 98
Pseudotatia 71, 88
parvanalis, *Kryptopterus* 50, 102
parvulus, *Hyphessobrycon* 48, 91
parvus, *Glossogobius* 40, 96
Gobius 42, 96
Sphoeroides 80, 103
pascual, *Entonanthias* 34, 102
patiae, *Curimatus* 30, 93
patzcuaro, *Chiostoma* 24, 87
- pauciradiatus*, *Antennarius* 9, 86
Diretmus 32, 94
Gastromyzon 38, 97
pectoralis, *Bathypterois* 15, 92
pediculatus, *Pseudancistrus* 70, 98
pejerato, *Sternopygus* 82, 103
pellucidum, *Characidium* 22, 90
peninsulae atrimentis, *Menidia* 56, 87
peruanus ricae, *Bryconamericus* 18, 90
Curimatus 30, 93
Pneumatophorus 67, 102
pescadorensis, *Liza* 53, 98
petrosus, *Brycon* 19, 90
phaeonota, *Moenkhausia* 57, 91
phalacra, *Brachyglanis* 17, 100
phantasia, *Cetopsorhamdia* 21, 100
phenax, *Plagiotremus* 66, 88
pheromystax, *Lepophidium* 51, 99
philippinus, *Paralepis* 63, 99
pholeter, *Astroblepus* 12, 87
phoxocephalum, *Characidium* 22, 90
- pileatus*, *Diceratias* 32, 94
pinnatus, *Deuterodon* 31, 90
piperata, *Tympanopleura* 84, 86
pirrense, *Cyclopius* 30, 87
pittieri, *Poecilia* 67, 101
plagiozonatus, *Aequidens* 6, 92
planiceps, *Ancistrus* 8, 98
planifrons, *Netuma* 60, 87
platycephalum, *Megalonema* 55, 100
plicatus, *Anostomus* 9, 86
poco, *Adioryx* 5, 97
poeciloides, *Hyphessobrycon* 48, 91
poecilus, *Microglanis* 57, 100
polynotatus, *Corythoichthys* 28, 103
polytaenia, *Leucogobio* 52, 94
potamius, *Gobiesox* 41, 95
potaroensis, *Aequidens* 6, 92
Astyanax 13, 90
Deuterodon 31, 90
Myoglanis 58, 100
pricei, *Emblemariopsis* 34, 89
profunda, *Moenkhausia* 58, 91
proops parahybae, *Pygidium* 73, 103
proteus, *Blenniulus* 16, 88
Hyphessobrycon 48, 91
protractila, *Bivibranchia* 16, 96
pseudaplites, *Micropterus* 57, 89
pteroides, *Characidium* 22, 90
puertoricensis, *Malacoctenus* 55, 97
pulvereus, *Zygonectes* 86, 94
pumilio, *Serraniculus* 79, 102
pumilus, *Melamphaes* 55, 98
Ogcocephalus 62, 99
punctatissimus, *Ophioscion* 62, 102
punctatum, *Cichlasoma* 26, 92
Hoplosternum 47, 88
Megalonema 55, 100
punctipectophorus, *Coryphopterus* 28, 96
punctulatum piurae, *Pygidium* 73, 103

- punctulatus*, *Enneapterygius* 34, 104
Gastromyzon 38, 97
purpurascens, *Nesiotes* 60, 101
purpurea, *Botia* 16, 93
purpureus, *archolepis*, *Anoplar-chus* 9, 103
pygmaeus, *Glossanodon* 40, 87
pyropunctata, *Rachovia* 74, 86
- quadrifasciatus*, *Hemiodus* 45, 96
quadrizonatus, *Thalassophryne* 83, 88
quercinum japonicum, *Macrostoma* 54, 99
- radcliffei*, *Spectrunculus* 80, 99
randalli, *Acanthurus* 4, 86
Chromis 25, 101
Entomacrodus 34, 88
Myripristis 59, 97
rastrous, *Caranx* 20, 89
regani, *Astyanax* 13, 90
Chirostoma 24, 87
Gambusia 37, 101
Rhamdia 74, 100
reidi, *Pomacentrus* 68, 101
reighardi, *Leucichthys* 52, 101
reinhardti, *Pygidium* 73, 103
remotus, *Chonerhinus* 24, 103
reticulatus, *Hyphessobrycon* 48, 91
Lophiodes 53, 98
retropinnis, *Leporinodus* 51, 86
rhodurus, *Oncorhynchus* 62, 102
rhomaleum, *Sparisoma* 80, 102
rhomboides, *Hemicaranx* 44, 89
ribeirae, *Astyanax* 13, 90
ribeiroi, *Bellator* 16, 104
Glanidium 39, 88
richardi, *Macrurus* 54, 98
richardsoni, *Ulua* 84, 89
riddlei, *Hemigrammus* 45, 91
Holopristis 47, 91
Toxus 83, 101
ridens, *Gastromyzon* 38, 97
ritense, *Heterogramma* 46, 92
rivasi, *Apuredoras* 10, 95
Limia 52, 101
rivuloides, *Gobio* 41, 94
robertsi, *Zenarchopterus* 85, 96
robusta, *Rhaphiobellone* 75, 88
robustum, *Megalonema* 55, 100
robustus, *Astyanax* 14, 90
Leiocassis 51, 88
Notropis 61, 94
Varicorhinus 85, 94
rocasensis, *Eupomacentrus* 35, 101
rodwayi, *Hemigrammus* 45, 91
roosevelti, *Metynnis* 56, 91
Salmo 77, 102
rosacea, *Melanotaenia* 56, 98
rosaceus, *Hyphessobrycon* 48, 91
rosae, *Chasmocranus* 23, 100
rosei, *Astroblepus* 12, 87
roseola, *Chaenopsis* 21, 89
rostellum, *Ogcocephalus* 62, 99
rostratus, *Sternarchus* 81, 87
- ruberrimus*, *Astyanax* 14, 90
rubescens, *Algansea* 7, 93
rubriques, *Moxostoma* 58, 89
rupununi, *Charax* 22, 90
russula, *Sagamia* 77, 96
- sabana*, *Dangila* 31, 94
sabanus, *Brachyogobius* 17, 95
Mystus 59, 88
Ompok 62, 102
sadina, *Parastremma* 63, 91
saepepallens, *Gobionellus* 41, 96
sagamiensis, *Diaphus* 32, 99
sagax, *Scopelarchus* 78, 102
saladonis, *Notropis* 61, 94
saludae, *Hololepis* 47, 100
salvadoris, *Roeboides* 77, 92
sandakanensis, *Acanthopthalmus* 4, 93
Glossogobius 40, 96
sanguinea, *Vandellia* 84, 104
santa, *Macgregorella* 54, 96
santaanae, *Pantosteus* 63, 89
santaeritae, *Pygidium* 73, 103
santaremensis, *Crenicichla* 29, 92
santarosaliae, *Notropis* 61, 94
sardina, *Melaniris* 56, 87
sargenti, *Pseudoplesiops* 71, 101
sarmenii, *Squaliolus* 81, 103
sasakii, *Liopempheris* 52, 99
satsumae, *Zaescopus* 85, 104
sauteri, *Pristiurus* 69, 102
scabripinnis paranae, *Astyanax* 14, 90
schmidtii, *Otophidium* 63, 99
Parahollandia 63, 103
schroederii, *Bathyclupea* 15, 88
schwartzii, *surinamensis*, *Corydoras* 27, 88
scierus apristis, *Hadropterus* 43, 100
Astyanax 14, 90
sciistius, *Xyrichtys* 85, 97
scopiferus guaytarae, *Bryconamer-icus* 19, 90
Bryconamericus 19, 90
scovelli makaxi, *Syngnathus* 82, 103
scythrops, *Flammeo* 36, 97
secundus, *Microglanis* 57, 100
sentaniensis, *Rhombattractus* 75, 98
sepikensis, *Rhombosoma* 75, 98
Zenarchopterus 85, 96
septentrionalis, *Pygidium* 73, 103
septentrionalis, *Coripareius* 27, 94
serrata, *Pimelodella* 66, 103
Rhinosardinia 75, 93
shideleri, *Moenkhausia* 58, 91
shropshirei, *Euleptoeleotris* 35, 95
sialis, *Amia* 7, 86
sicaria, *Evermannella* 35, 95
siebethalae, *Brycon* 19, 90
silus, *Chonerhinus* 24, 103
Porogadus 69, 99
simoni, *Crenicichla* 29, 92
simplex, *Nannostomus* 59, 97
simus, *Creagrutus* 28, 90
sinusarabici, *Dollfusichthys* 32, 93
- snyderi*, *Apogon* 10, 86
Hyporhamphus 49, 96
Puntius 72, 94
Taenioides 82, 96
soldatovi minulus, *Gobio* 41, 94
solidus, *Arius* 11, 87
sonorensis, *Catostomus* 21, 89
sparsidens, *Chromis* 25, 92
spilotoceps, *Epinephelus* 35, 102
pilotum, *Cichlasoma* 26, 92
spiniger, *Spinoblennius* 80, 88
spinus, *Ancistrus* 8, 98
stagnatus, *Rivulus* 76, 86
stanleyi, *Coregonus* 27, 101
starksi, *Menidia* 56, 87
starri, *Pomadasys* 68, 96
staurophor, *Lepophidium* 51, 99
stellatum, *Pygidium* 73, 103
stellatus, *Bodianus* 16, 102
stellifer, *Rivulus* 76, 86
stenodon, *Cheirodon* 23, 90
stewarti, *Argentina* 11, 87
Heptapterus 46, 100
Loricaria 54, 98
steyermarki, *Leporinus* 52, 86
stictus, *Hyphessobrycon* 48, 91
stigma, *Pomacentrus* 68, 101
stigmaturus, *Distichodus* 32, 93
straminium, *Pygidium* 73, 103
streber nerensis, *Zingel* 85, 100
striatum, *Pygidium* 73, 103
strophodes, *Lepidaplois* 51, 97
stroudi, *Gelanoglanis* 38, 88
submarginatus, *Loricaria* 54, 98
suensis, *Micrognathus* 57, 103
suwae, *Gnathopogon* 40, 94
svetovidovi, *Belone* 16, 88
symmetricus, *Myripristis* 59, 97
- tabira*, *Acheilognathus* 5, 93
tabl, *Decapterus* 31, 89
taenia melanoleuca, *Cobitis* 27, 93
taeniatum pertense, *Heterogramma* 46, 92
taiwanensis, *Pseudobagrus* 70, 88
taiwanus, *Rhinogobius* 75, 96
tamuisiensis, *Scaphesthes* 78, 94
tanakae, *Lycodes* 54, 104
Zesticthys 85, 104
tanakai, *Caranx* 20, 89
tarpon, *Genycharax* 38, 90
taurocephalus, *Hyphessobrycon* 48, 91
tegatus, *Vesicatrax* 85, 92
temminckii, *Alectis* 6, 89
templei, *Pomadasys* 69, 96
tenuis, *Cichlasoma* 26, 92
tenuis, *Callionymus* 20, 88
Cynodonichthys 30, 86
Pellonula 64, 93
Poecilia 68, 101
terminalis, *Parodon* 64, 99
tetranema, *Bathybagrus* 15, 88
thamalakanensis, *Alestes* 6, 89
thomersoni, *Rhinodoras* 75, 95
thompsoni, *Sebastodes* 79, 102

- thurni*, *Roeboides* 77, 92
tigris, *Alepideleotris* 6, 95
tiki, *Myripristis* 59, 97
timagamiensis, *Salvelinus* 77, 102
tlahuacensis, *Evarra* 35, 94
tolimae, *Bryconamericus* 19, 90
toni posteroventralis, *Barbatula* 15, 93
tosae, *Zalescopus* 85, 104
toweri, *Goodea* 42, 96
townsendi, *Parascloopsis* 63, 99
transvestitus, *Nanochromis* 59, 92
tremantinae, *Odontostilbe* 61, 91
trewavasae, *Sciaena* 78, 102
triangulatus, *Malacoctenus* 55, 97
tricuspidata, *Atuona* 14, 95
trifasciatum maciliense, *Heterogramma* 46, 92
Cyclopium 30, 87
triguttatum, *Pygidium* 73, 103
tristegus marquesensis, *Acanthurus* 4, 86
tropicus, *Platyopocilus* 67, 101
troscheli, *Glyphisodon* 40, 101
tsotsorogensis, *Barbus* 15, 94
tsuchigae, *Gnathopogon* 41, 94
tuba, *Cichlasoma* 26, 92
tubbi, *Pangasius* 63, 99
tuckeri, *Monacanthus* 58, 88
tumacoensis, *Umbrina* 84, 102
turrubarensis, *Gambusia* 37, 101
tutuila, *Centropyge* 21, 101
tuyra, *Arius* 11, 87
tuyrense, *Cichlasoma* 26, 92
tuyrensis, *Loricaria* 54, 98

uii, *Caranx* 20, 89
ulae, *Synodus* 82, 103
umalii, *Puntius* 72, 94
umbratilis, *Gambusia* 38, 101
umbriceps, *Alepocephalus* 6, 86
umbriferum, *Cichlasoma* 26, 92
undecimalis, *Myripristis* 59, 97
unicolor, *Echidna* 33, 99
unicuspis, *Ichthyomyzon* 49, 100
unidorsalis, *Helogenes* 43, 96
unifasciatum, *Cyclopium* 30, 87

uracantha rupestre, *Loricaria* 54, 98
urophthalmus mayorum, *Cichlasoma* 26, 92
urosilotus, *Gymnapogon* 42, 87
uruguayanae, *Macropsobrycon* 54, 91
ushiei, *Dasyatis* 31, 94
uxorius, *Bombonia* 16, 103

vaivasensis, *Ambassis* 7, 89
vanceae, *Cyclopium* 30, 87
variabilis, *Starksia* 81, 97
variatus, *Platyopocilus* 67, 101
variegata venezuelae, *Loricaria* 54, 98
Skiffia 80, 96
variegatus, *Microglanis* 57, 100
velutinus, *Hemipimelodus* 45, 87
venezuelae, *Ophioscion* 62, 102
ventrale, *Cyclopium* 30, 87
verecundus, *Spirinchus* 81, 99
vermiculatum, *Pygidium* 73, 103
verrucosus, *Antennarius* 9, 86
villadolidi, *Ctenogobius* 29, 96
Chromis 25, 101
villiersi, *Physalia* 65, 102
villosus, *Lithogenes* 53, 98
Pseudopimelodus 71, 100
vineta, *Echidna* 33, 99
vintoni, *Characidium* 22, 90
virgata, *Albula* 6, 86
virgatulus, *Ctenogobius* 29, 96
viridesens, *Gunnellichthys* 42, 98
vittata, *Gambusia* 38, 101
volcanus, *Rivulus* 76, 86

waigiensis, *Ctenogobius* 29, 96
Enneapterygius 34, 104
waimacui, *Rivulus* 76, 86
wakiyae, *Malakichthys* 55, 100
walensis, *Salarias* 77, 88
waltersi, *Batrachoides* 15, 88
weberi, *Chromis* 25, 101
weitzmani, *Corydoras* 28, 88
whaleri, *Gephyrocharax* 39, 90
whitei, *Goodea* 42, 96
wichmanni, *Apogon* 10, 87

wilburi, *Callionymus* 20, 88
Macrodonogobius 54, 96
williamsi, *Anisotremus* 9, 96
wilsoni, *Hemiancistrus* 43, 98
Phenagoniates 65, 91
wonderi, *Gobiocichla* 41, 92
woodsii, *Anthias* 9, 102
Chromis 25, 101
Gobiopsis 41, 96
Microdesmus 56, 98
Myripristis 59, 97
Rhynchodoras 75, 95

x-punctata trautmani, *Hybopsis* 47, 94
x-punctata, *Hybopsis* 47, 94
xantherythrus, *Holocentrus* 47, 97
xanthomelas, *Brachygobius* 17, 95
xanthosoma, *Gambusia* 38, 101
xanthum, *Megalonema* 55, 100
xenandrus, *Engyproson* 34, 88
xenicus, *Calymnichthys* 20, 89

y-lineata, *Starksia* 81, 97
yamamotis, *Moroco* 58, 94
yarkandensis sellaefer, *Barbatula* 15, 93
yarrowi, *Agosia* 6, 93
yoshiae, *Anarchias* 7, 99
yucata australis, *Gambusia* 38, 101

zaca, *Malacoctenus* 55, 97
zanectes, *Calliurichthys* 20, 88
zebra, *Characidium* 22, 90
zebrinus, *Phrynelox* 65, 86
zeteki, *Bryconamericus* 19, 90
zirahuen, *Chirostoma* 24, 87
zonatum, *Cichlasoma* 26, 92
Pygidium 73, 103
zonatus, *Bostriechus* 16, 95
zonifer sudensis, *Malacoctenus* 55, 97
zonophaea, *Echidna* 33, 99
zuloagai, *Doraops* 32, 95